



PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/19005>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.

IMMUNOTOXINS IN HEMATOLOGY

Ypke van Oosterhout



IMMUNOTOXINS IN HEMATOLOGY

Ypke van Oosterhout

Van Oosterhout, Y.V.J.M.	Immunotoxins in Hematology
ISBN	90-9014622-9
NUGI	743
Printing:	Ponsen & Looijen B.V., Wageningen
Graphic Design:	Marie-Louise Dusée
Picture used for cover:	The procession of the Trojan Horse into Troy by Giovanni Domenico Tiepolo ©National Gallery Picture Library, London

No part of this book may be reproduced by any mechanical, photographic or electronic process, nor may it be stored in a retrieval-system, transmitted, or otherwise copied for public or private use, without the written permission of the author.

IMMUNOTOXINS IN HEMATOLOGY

Een wetenschappelijke proeve op het gebied van de Medische Wetenschappen

PROEFSCHRIFT

ter verkrijging van de graad van doctor aan de Katholieke Universiteit Nijmegen
volgens besluit van het College van Decanen
in het openbaar te verdedigen op
vrijdag 20 april 2001 des namiddags om 1.30 uur precies.

door

Ypke Vincentius Johannes Maria van Oosterhout

Geboren 30 mei 1964 te Nijmegen

Promotor:

Prof. dr. T.J.M. de Witte

Co-promotores:

Dr. F.W.M.B. Preijers

Dr. A.V.M.B. Schattenberg

Manuscriptcommissie:

Prof. dr. P.H.M. de Mulder (voorzitter)

Prof. dr. L.A. Monnens

Prof. dr. W.E. Fibbe (LUMC)

*Aan Joyce, Jelle, Youp & Merel,
aan mijn ouders*

Financial support for publication of this thesis was kindly provided by:

Amersham Pharmacia Biotech

Amgen B.V.

Ernst & Young Accountants

Farma Research B.V.

Immunotoko B.V.

Medarex, Inc.

OctoPlus B.V.

Sigma-Tau Ethifarma B.V.

Synthon B.V.

VEREENIGDE

Novartis Pharma B.V.

Roche Nederland B.V.

Shering-Plough B.V.

Inveresk Research Ltd.

IQ products B.V.

SANGSTAT

TABLE OF CONTENTS

	Synopsis	9
Chapter 1	Immunotoxins, general introduction	13
Chapter 2	Immunotoxins in Hematology	47
Chapter 3	Cytotoxicity of CD3-ricin A chain immunotoxins in relation to cellular uptake and degradation kinetics <i>Cancer Res</i> 1992; 52: 5921-5925	65
Chapter 4	Effect of isotype on internalization and cytotoxicity of CD19-ricin A-immunotoxins <i>Cancer Res</i> 1994; 54: 3527-3532	79
Chapter 5	Flow cytometric evaluation of immunotoxin induced cell kill <i>Prog Clin Biol Res</i> 1994; 389: 89-95 & <i>Leukemia</i> 1999; 13: 241-249	97
Chapter 6	Laboratory production of anti-CD3 and Anti-CD7 ricin A-immunotoxins for a clinical pilot study <i>Int J Pharm</i> , in press	109
Chapter 7	A combination of anti-CD3 and anti-CD7 ricin A-immunotoxins for the in vivo treatment of acute graft-versus-host disease <i>Blood</i> 2000; 95: 3693-3701	127
Chapter 8	Summary and discussion	151
	Samenvatting	163
	List of abbreviations	173
	List of publications	177
	Dankwoord	181
	Curriculum vitae	187

SYNOPSIS

Already hundred years ago, Paul Ehrlich envisioned the use of antibodies for the specific targeting of cancer cells without causing harm to healthy tissue. With the invention of the hybridoma technique, Ehrlich's magic bullet concept seemed to come within practical reach. This technique, as described by Kohler and Milstein in 1975, enabled the production of antibodies of defined specificity in sufficient quantities for clinical therapy. As a consequence, a number of clinical trials were initiated, using various monoclonal antibodies (MoAbs) to target diffuse and solid tumors and varying immune diseases.

OKT3 was the first therapeutic MoAb to be approved for marketing by the US Food and Drug Administration (FDA) in 1986. This murine MoAb demonstrated activity in reversing acute organ transplant rejection. MoAb 17-1A, registered in 1994 in Germany, was the first MoAb for the treatment of cancer to win approval in any country. This murine MoAb significantly reduced post-operative mortality in colon cancer patients when applied as adjuvant therapy. Unfortunately, these two cases appeared to be exceptions rather than the forebode of a new class of broadly effective therapeutics. In general, responses to MoAb therapy were poor, if any, and their administration resulted frequently in allergic type reactions. Excuses were found in the immunogenicity of the xenogenic MoAbs, the heterogenic expression or modulation of antigen, insufficient accessibility of target cells, the presence of MoAb-capturing circulating free antigen, and limited interaction of the (mostly murine) MoAbs with the host immune effector mechanisms.

Though far from satisfying, these initial experiences formed the base for the development of improved antibody strategies. These include the (partial) substitution of the constant regions of the murine MoAbs with their human counterparts to render them less immunogenic and to improve their interaction with the host immune system. More recently, different techniques have become available which enable the production of fully human MoAbs or fragments thereof. Another strategy forms the generation of bispecific antibodies with affinity for both the target cell and an immune effector cell. By bridging these cells, bispecific antibodies aim at the maximal exploitation of the patient's immune system. As an alternative, MoAbs have been conjugated to effector molecules, like radioisotopes or plant/bacterial derived toxins, in an attempt to improve on their intrinsic, host independent, cytotoxicity.

The implication of these technological and conceptual advances strongly boosted the scientific as well as pharmaceutical interest. In line with this MoAb-revival, the FDA approved its first therapeutic antibody for treatment of cancer in 1997. This humanized CD20 MoAb (Rituximab) demonstrated efficacy in the treatment of certain forms of B-

cell Non-Hodgkin's lymphoma. Since then, an additional five (partly) humanized MoAbs have been approved by the FDA for the treatment of certain forms of cancer and immune disorders. Moreover, with about 60 MoAb-based agents currently being tested in the clinic, MoAb represent the second largest category of biotechnology products in clinical trials.¹

The present thesis focuses on the approach of antibody-toxin conjugates, the so called immunotoxins (IT). More specifically, the potential pitfalls, as well as the drawbacks and therapeutic advantages of these agents when applied in the field of hematology will be highlighted.

REFERENCE

1. Walsh G: Biopharmaceutical benchmarks. *Nat Biotechnol* 18:831-833, 2000

IMMUNOTOXINS, GENERAL INTRODUCTION



CONTENTS

- 1.1 General principle
- 1.2 Monoclonal antibody part
- 1.3 Toxins
- 1.4 Linking MoAb and toxin
- 1.5 Non-toxin based effector mechanisms
- 1.6 Lessons from clinical trials

1.1 GENERAL PRINCIPLE

Conventional ITs consist of MoAbs conjugated to a potent toxin, mostly of bacterial or plant origin. Their mechanism of action is depicted in Figure 1. The MoAb-moiety first binds specifically to its antigen expressed on target cells, following which the entire anti-gen-IT complex is internalized. Once inside the cell, the bond between the MoAb and toxin is broken, thereby releasing free toxin into the cytoplasm. The toxin then irreversibly inhibits protein synthesis by means of a catalytic reaction culminating in cell death. Because the toxin-moiety is unable to enter the cell autonomously, and is inactive outside the cell, ITs are only hazardous to cells capable of binding and internalizing the MoAb.

Instead of MoAbs, or fragments thereof, cells can also be targeted by cytokines which receptor is (over)expressed on the target cell. As an alternative to chemical linkage, MoAb-fragments or cytokines can also be genetically spliced to a toxin, generating a recombinant IT.



Figure 1 *Mechanism of action*

1.2 MONOCLONAL ANTIBODY PART

Antibody structure

Antibodies can be considered as bifunctional molecules, comprising a highly variable antigen-binding domain (V domain) on one site and a constant effector domain (C domain) on the other site (Figure 2). Based on their physical properties, antibodies can be divided in IgM, IgG, IgA, IgE, and IgD classes, each having its own function in the host defense system. IgG is the antibody class most widely exploited for clinical use. The IgG molecule comprises two heavy chains of the structure $VH-CH_1-CH_2-CH_3$ and two light chains of the structure $VL-CL$. The heavy and light chain are linked to each other by a disulphide bond between CH_1 and CL . Additional disulphide bonds formed in the so called hinge region of the two heavy chains result in the formation of homodimers, containing two identical antigen binding-sites. The actual antigen-recognition site is provided by the complementarity-determining regions (CDR), formed by three hypervariable loops of each the VH and VL domain ($CDRH1-3$ and $CDRL1-3$, respectively). This antigen-combining region is immunologically defined as the antibody's idiotype. The IgG effector functions are located in the Fc-part in the form of an $C1q$ complement-binding site and Fc-receptor binding site.

When considering MoAbs solely as a toxin transporter, it makes sense to use only those fragments containing the binding function of the CDR's, such as the 110 kDa bivalent $F(ab')_2$ fragment, or the monovalent Fab' or Fab fragments (Figure 2). The even smaller recombinant single-chain Fv fragments (sFv) contain only the VL and VH domain connected to each other through a short flexible peptide spacer. In disulfide stabilized Fv fragments (dsFv), the peptide linker is replaced by a more stable disulfide bond introduced by genetically engineering cysteine residues in VL and VH at positions allowing post-translational disulfide bond formation. The Fv-fragments are only 26 kDa in size, less than 20% of the 155 kDa IgG.




Figure 2 Whole IgG and IgG-derived fragments used in Immunotoxins

Removing the Fc portion of a 155 kDa IgG molecule results in $F(ab')_2$ fragments of 110 kDa. These can be converted to Fab' fragments by reduction, proteolytic treatment or by recombinant means. Single-chain Fv (scFv) fragments consists of the variable domain of a MoAb joined by a linker (26 kDa). In a dsFv the linker is replaced by a more stable disulfide bond.

Of note, in particular cases, the MoAb Fc-part can be used as the targeting moiety. An example forms the treatment of allergic disorders by the elimination of high-affinity Fc-epsilon-receptor I (Fc ϵ RI) expressing mast cells and basophils by toxin-conjugated IgE Fc-fragments.¹⁻³

Specificity and distribution of antigen

The MoAb's binding specificity is the major property determining its suitability as a toxin transporter. Ideally, the MoAb binds an antigen that is expressed solely on those cells that are to be eliminated. The Ig-idiotypic expressed by malignant B cells and myeloma cells, for instance, represents a tumor-clone specific marker.⁴⁻⁹ A more general tumor specific marker is formed by the mutant epidermal growth factor receptor (EGFRvIII), expressed in glioblastoma, and carcinomas of breast and lung. EGFRvIII contains an in-frame deletion resulting in the surface expression of a tumor-specific extracellular sequence. ITs directed against this tumor-specific target demonstrated

clear cytotoxicity to EGFRvIII transfected cells, while showing little or no efficacy towards cells expressing high levels of normal EGFR.^{10,11}

In practice, however, tumor restricted antigens are rare and differences in expression relative to non-target tissue are generally quantitative rather than qualitative (*e.g.* tissue-associated rather than tumor specific). About 25-30% of patients with breast and ovarian cancer shows a 10 to 100-fold overexpression of the human epidermal growth factor receptor 2 (HER-2, *erbB2*). This phenotype, caused by a mutation in the HER-2/*neu* gene, is associated with a rapid disease progression and a poor prognosis.¹² Treatment of this particular patient group with anti-HER-2 MoAb resulted in objective responses.¹³ Several groups are now working on anti-HER-2 ITs in an attempt to further improve on the clinical efficacy.¹⁴⁻²⁰ Other tumor-associated markers which have been clinically exploited as target for ITs include lewis^x (Le^x) expressed on a variety of solid tumors²¹, high-molecular-weight melanoma associated antigen (HMW-MAA) expressed on melanoma²²⁻²⁶, CD56 (neural cell adhesion molecule, NCAM) associated with small cell lung cancer²⁷⁻²⁹, and carcinoembryonic antigen (CEA) overexpressed on adenocarcinomas³⁰. The clinical use of ITs directed against Le^x clearly illustrated the potential limitations of targeting tumor-associated antigens.²¹ The Le^x-expression on normal vascular endothelial cells resulted in serious vascular leak syndrome (VLS) preventing the administration of sufficient IT to achieve therapeutic blood levels.³¹ Similarly, the HER-2 expression on normal liver cells caused hepatotoxicity in all six patients treated with an anti-HER-2 IT.²⁰ Generally, though, it has been demonstrated that ITs can be delivered safely to tumor-associated antigens without severe irreversible toxicities occurring. Moreover, target-antigen expression on healthy bystander tissue is less critical in case of strongly regenerative tissue. An illustrative example from the hematopoietic lineage markers.³² Though an effective therapy is expected to affect healthy lineage cells as well, early progenitor cells will rapidly restore the hematopoietic compartment. At current, hematological disorders represent the group most effectively targeted with immuntoxins (see also Chapter 2).

The opposite concern of cross-reactivity is the occurrence of antigen negative (or non-overexpressing) target cells. This can be at the population level, *e.g.* less than 30% of patients with breast/ovarian cancer over-expresses HER-2, but may also be at the patient level, due to heterogeneous antigen expression within the tumor.³³⁻³⁵ In practice, an arbitrary threshold is often set defining the minimal percentage of antigen-positive cells required for therapy. In certain cases, co-administration of cytokines may increase the expression of tumor-associated antigens. An example forms the IFN- γ induced upregulation of the tumor-associated antigen 72 (TAG-72) and CEA on colon carcinoma.³⁶⁻³⁹ Moreover, the use of an appropriate combination of immunotoxins may circumvent the heterogeneity observed in the expression of individual antigens.⁴⁰⁻⁴⁴ Additional obstacles preventing an optimal targeting are a concentration of natural ligand,

competing with the MoAb for binding, or a high amount of circulating free antigen or antigen expressing normal cells, which may both act as an antigen sink.^{45,46}

As an attractive alternative to direct tumor-targeting, Burrow *et al.* introduced the concept of targeting the tumor-nursing vascular bed.⁴⁷⁻⁴⁹ This approach affords several important advantages: (1) the pivotal problem of poor tumor penetration is circumvented, (2) even incomplete destruction of the vascular bed results in massive tumor destruction, and (3) a single immunotoxin can be used for the treatment of a variety of tumors of different origin. The proof of principle was generated in nude mice grafted with IFN- γ transfected tumor cells. These cells induced an artificial expression of major histocompatibility complex (MHC) Class II antigens on the surrounding tumor vascular endothelium. The subsequent administration of anti-Class II directed IT caused extensive thrombosis of the tumor vasculature, inducing permanent complete remissions in over half of the animals. In later trials, comparable results could be obtained using ITs reactive with endoglin, a natural proliferation-associated antigen of endothelial cells.⁵⁰⁻⁵² More recently, 'clotting initiating human tissue factor' proved in this setting to be a useful alternative for the conventional plant or bacterial derived (immunogenic) toxin-moiety.⁵³⁻⁵⁵

Internalization of MoAb-antigen complex

Following antigen-binding, the toxin-conjugated MoAbs needs to be internalized to deliver the toxin to the cytoplasm. Preijers *et al.* demonstrated for several anti-T-cell ITs that the absolute amount of internalized molecules rather than the antigen surface expression determines the efficacy.⁵⁶ The amount and kinetics of internalization itself are determined by the combination of antigen, MoAb and target cell.⁵⁶⁻⁶⁷

Surface antigens differ widely regarding their intrinsic internalization capacity. Some appear to be good internalizers, like T-cell antigens CD3 and CD7 and B-cell antigen CD22, while others hardly internalize, like T-cell antigen CD8 or B-cell antigen CD20.^{56,68-71} The use of MoAb-fragments instead of whole IgG generally reduces the efficacy of the respective IT, either due to a decrease in antigen-binding or in the subsequent internalization.^{66,72-75} Anti-CD5 ITs form an exception and are more effective with F(ab')₂ or Fab'-fragments than with whole IgG.^{65,67,76} For CD19-ITs it has been demonstrated that the MoAb isotype may affect the internalization and, thereby, the efficacy of the IT. This phenomenon could be attributed to an interaction of the Fc-part of murine IgG1 MoAb (and not IgG2a MoAb) to the Fc-gamma-receptor IIa (Fc γ RIIa). Analogous to the mechanism as described by Kurlander, this facilitated the internalization of antigen-bound IT.⁷⁷

Once internalized, the intracellular routing determines the ultimate efficacy of the immunotoxin.^{59,69,78,79} The IT needs to be transported to an appropriate subcellular compartment to facilitate the translocation of the toxin-moiety to the cytoplasm. As a result, ITs with comparable internalization rates may differ greatly in efficacy. Anti-CD19

ITs, for example, are generally less effective than anti-CD22 ITs, despite similar or higher internalization rates.⁸⁰⁻⁸² For T-cell antigen CD2, and the B-cell antigen surface-IgD, it has been demonstrated that MoAb binding epitopes closer to the cell surface make superior ITs. This was also attributed to a favorable intracellular handling.^{59,62}

The above picture becomes even more complex considering that plant and bacterial toxins prefer different routes of internalization for optimal efficacy (see Paragraph 1.3).^{58,83} Therefore, an indirect sandwich cytotoxicity assay has been developed which facilitates the *in vitro* screening of particular MoAb-toxin conjugates.⁸⁴ In this assay, dilutions of MoAbs are incubated with the target cells, followed by the binding of a second (isotype-specific) antibody-fragment conjugated to a toxin of choice. This technique allows the quick screening of a panel of MoAbs in combination with different toxins.⁸⁴⁻⁸⁷ The only concern is that the indirect approach might influence the internalization and/or intracellular handling of the particular toxin. Using this assay, we dramatically underscored the potential of a CD7 MoAb that proved highly effective when conjugated directly to the toxin applied (WT1-dgA, unpublished observations).

1.3 TOXINS

The toxins most commonly applied for the construction of ITs are derived from either bacteria or plants and have in common that they catalytically and irreversibly disrupt protein synthesis, typically at picomolar concentration (Figure 3). The bacterial toxins most extensively studied so far are *Pseudomonas* exotoxin (PE) and diphtheria toxin




Figure 3 Schematic representation of toxin molecular structure.

PE, *Pseudomonas* exotoxin: Ia, Ib - binding domain, II - translocation domain, III - catalytic domain. DT, diphtheria toxin: A, catalytic fragment, B, binding fragment. The shaded segments mark the position of the protease-sensitive loop region. Ricin: A, active chain, B, binding chain.

Figure adapted from E.J.Wawrzyniczak.²⁷⁷

(DT). The plant toxins, also referred to as ribosome inhibitory proteins (RIP), can be divided into holotoxins (type II RIP), including abrin, mistletoe lectin, modeccin, and ricin, and hemitoxins (type I RIP), such as gelonin, pokeweed antiviral protein (PAP), and saporin.

Pseudomonas exotoxin

PE contains three functional domains, responsible for binding (domain Ia), translocation (domain II), and protein synthesis inhibition (domain III). Domain Ib, of yet unknown function, joins domain II and the C-terminal domain III.⁸⁸⁻⁹⁰ Native PE is excreted by the bacterium as a 66 kDa single-chain protein. The N-terminal domain Ia binds the α_2 -macroglobulin receptor present on the surface of all eukaryotic cell types. Following endocytosis, domain II is proteolytically cleaved at a low pH by cellular protease furine, resulting in two disulfide-bonded fragments of 28 and 37 kDa.^{91,92} The 37 kDa fragment contains part of the translocation domain II and the active enzyme domain III. The latter contains a REDLK sequence which facilitates the transport of the 37 kDa fragment from the transreticular Golgi apparatus to the endoplasmic reticulum (ER).⁹³ There, the remaining domain II fragment mediates the translocation of domain III to the cytoplasm. In the cytoplasm, domain III inactivates elongation factor 2 (EF-2) by its ADP ribosylating activity resulting in the cell's death. Current PE-based IT use truncated forms of PE, in which the entire Ia domain (PE40)⁹⁴⁻⁹⁷ and frequently also part of the Ib domain (PE38, PE35)⁹⁸⁻¹⁰⁵ have been deleted. Cell targeting ligands are chemically or genetically linked to the N-terminus. In some cases, the C-terminal REDLK sequence is altered to a KDEL sequence.¹⁰⁶⁻¹⁰⁹ This sequence has a higher affinity for the ER KDEL receptor and, therefore, results in an increased efficacy of the respective IT.

Diphtheria toxin

DT compromises an A and B-domain, which are responsible for cell binding and inhibition of proteins synthesis, respectively.⁸⁸ The crystal structure reveals a third domain, the translocation domain (T), located within the B-domain in the center of the molecule.¹¹⁰ Native DT is released in the bacterial medium as a single polypeptide chain of 535 amino acids of approximately 62 kDa. Outside the cell, DT is proteolytically cleaved between the C-terminal A (24 kDa) and N-terminal B-domain (38 kDa), resulting in a disulfide bonded two-chain form. The C-terminal B-domain binds to a complex of heparin-binding EGF-like growth factor precursor and CD9. After endocytosis, the disulfide bond is broken and the low endosomal pH promotes a conformational change of the T domain resulting in the formation of a transmembrane channel through which the enzymatic A domain can enter the cytoplasm.^{111,112} In the cytosol, the A domain disrupts cellular protein synthesis by ADP ribosylation of EF-2. For the construction of IT, the C-terminus of either a binding site mutant (a.o. CRM107, CRM9)^{113,114} or C-terminal truncation mutant (DT388, DT390, DAB486, DAB389)¹¹⁵⁻¹¹⁸ is linked to a cell binding ligand. Because most people are vaccinated against DT, C-terminal truncation

mutants are to be preferred as pre-existing anti-DT antibodies may primarily be directed against the cell binding B-domain.¹¹⁸

Plant holotoxins toxins

Plant holotoxins compromise two disulfide-bonded chains: a protein synthesis inhibiting A-chain, and a cell binding and translocation facilitating B-chain.¹¹⁹ Ricin, a natural product derived from the castor bean (*Ricinus Communis*), is the plant holotoxin most extensively tested *in vivo*. Native ricin is synthesized as a 64 kDa single chain pro-enzyme containing a 12 residue linking peptide joining the 32 kDa A- and B-chain. Following post-translational proteolytic cleavage, an interchain disulfide bond links the two chains. Cellular intoxication starts by binding of cell surface galactose residues by the three lectin domains within the B-chain.¹²⁰ After endocytosis, a considerable portion of the internalized ricin is transported and degraded in the lysosomes.¹²¹ A fraction of the internalized toxin is delivered to the trans-Golgi network. The further routing appears to be dependent on receptors recycling between the terminal compartments of the Golgi stack and the ER.¹²²⁻¹²⁴ The B chain dissociates from the A chain somewhere during the transport to the ER. In the ER, the A chain is translocated to the cytoplasm and inhibits protein synthesis through N-glycosidase activity removing the base of A⁴³²⁴ in 28S rRNA, thus preventing the association of EF-1 and EF-2.¹²⁵

For clinical use, the binding capacity of the B-chain has to be eliminated either by addition of excess galactose or lactose¹²⁶ (only for *ex vivo* applications), chemical blocking of the lectin-sites e.g. with galactose-based affinity labels (blocked ricin)¹²⁷⁻¹³¹, or omitting the entire B-chain¹³². An additional concern forms the high mannose and fucose carbohydrate groups present on A and B-chain, which are bound by receptors on liver Kupfer cells leading to rapid clearance and hepatotoxicity.¹³³ Using either chemically deglycosylated ricin A (dgA) or non-glycosylated recombinant ricin A (rA) has successfully circumvented this problem.¹³⁴⁻¹³⁷ Notably, deglycosylation is not an option for blocked ricin due to the galactose-based blocking of lectin activity.

The cytotoxicity of ricin A-based ITs is substantially lower than that of blocked ricin-based ITs.^{128,138} This can largely be ascribed to the membrane translocation enhancing capability of the B-chain.^{129,131,139} Moreover, the B-chain seems to protect the A-chain from intracellular degradation.¹⁴⁰ Whereas PE and DT are dependent on proteolytic cleavage for enzymatic activity, the ricin A chain is largely inactivated by endosomal or lysosomal proteases. Compounds that neutralize the low pH of intracellular compartments, or interfere with the routing to the lysosomes, may strongly enhance the cytotoxicity of ricin A-based IT.¹⁴¹⁻¹⁴³ Most commonly used are lysosomotropic agents, like NH₄Cl and chloroquine, and carboxylic ionophores, like monensin and nigericin.^{68,144-150} Though biologically active concentrations are usually not achievable *in vivo*, these agents have proven to be useful for *ex vivo* applications like the purging of hematopoietic stem cell grafts.¹⁵¹⁻¹⁵³

Plant Hemitoxins

Plant RIP I Hemitoxins, like saporin, gelonin, and pokeweed antiviral protein (PAP), contain no cell binding domain and resemble the A-chain of RIP II in size and in function.¹¹⁹ For the construction of ITs, RIP I are typically conjugated directly to a cell targeting ligand without further modifications.¹⁵⁴⁻¹⁵⁸

Alternative toxins

Li and Ramakrishnan constructed a recombinant hybrid plant-bacterial toxin by fusing the coding region of ricin A-chain to the A-domain of DT. The rationale is that since bacterial and plant toxin act at different steps in protein translocation, a combination of their activities could be more effective. In fact, ITs containing these chimeric toxins were *in vitro* 100-1000 fold more effective than ITs prepared with either toxin alone.¹⁵⁹

Though extremely potent, the major drawback of the bacterial and plant derived toxins is their proven immunogenicity. Especially now human targeting ligands (antibodies and cytokines) are readily available, it makes sense to search for human toxin equivalents as well. So far, several human Rnases, Dnases, and human TNF α have been used to construct fully human ITs.¹⁶⁰⁻¹⁷² Though not as toxic as their plant/bacterial counterparts, the initial *in vitro* results appear to be encouraging. The major benefit of human ITs is to be expected in the treatment of diseases which ask for successive administrations (e.g. autoimmune disorders). Since they have not been clinically tested yet, it remains an open question whether the junction of human toxin and cell targeting ligand does not form a new antigenic determinant.

As an alternative to the large catalytic toxins (typically ≥ 30 kDa), several synthetic chemotherapeutic molecules have also been coupled to MoAbs. These include vinblastine, doxorubicin, daunomycin, methotrexate, adriamycin, maytansinoids and calicheamicin.¹⁷³⁻¹⁸² Due to their smaller size (< 1 kDa), these agents may improve tumor tissue distribution and will probably be less immunogenic. As compared to their unconjugated forms, these agents proved more effective, and cause less side effects, following conjugation to the MoAb. Their intracellular mechanism of action, however, is identical. As such, they lack the advantage of plant or bacterial derived toxins which, by there alternative mode of cytotoxicity, may overcome acquired drug resistance to conventional therapy.¹⁸³⁻¹⁸⁹

1.4 LINKING MONOCLONAL ANTIBODY AND TOXIN

Chemical conjugates

Conventional ITs are prepared by linking a toxin fragment and cell binding ligand using a chemical crosslinker. Early studies demonstrated that plant toxin-based ITs should be conjugated by a disulfide-bond containing linker to enable the essential intracellular dissociation of ligand and toxin.^{126,190-192} For maximal stability outside the cell, crosslink-

ers have been developed that contain a benzene ring and methyl group around the disulfide bond to protect it from reduction by thiols in the blood and tissues.^{193,194} When using blocked ricin, a nonreducible thioether bond is preferred as the natural occurring interchain disulfide bond is already present. The same holds true for truncated bacterial toxins, still containing their native protease-sensitive sequence and interchain disulfide bond.

The chemical conjugation procedure has some potential disadvantages. The ligand conjugation is based on the derivatization of lysine residues, some of which may be located within the antigen binding domain. The same holds true for derivatization of the toxin, which may interfere with its catalytic activity.¹⁹⁵ In general, chemically constructed ITs form a heterogeneous mixture with respect to derivatization sites and the number of toxins per IT molecule. Recently, production methods have been developed resulting in an increased product uniformity.¹⁹⁶ There is no clear consensus, though, regarding the optimal number of conjugated toxins per ligand.^{81,197-200} ITs should be individually tested to determine whether the addition of two versus one toxin improves the *in vivo* cytotoxicity. If a ligand-toxin combination forms an effective conjugate, the conventional chemical production method has proven to be straightforward with good production yields, especially for whole IgG ITs. Moreover, the chemical linkage allows a great range of available toxins to choose from.

Recombinant fusion toxins

Recombinant fusion toxins are prepared by genetically splicing a toxin encoding DNA to a gene encoding for a MoAb-fragment or other cell binding ligand, and expressing the resulting construct as a single chain fusion protein. This approach is especially suited for the truncated bacterial toxins, which contain a native protease-sensitive sequence required for intracellular dissociation. Plant toxin-based ITs, in contrast, require post-translational removal of the interchain peptide sequence to facilitate intracellular ligand dissociation. Ricin A-based fusion toxins compromising an artificial protease sensitive linker, have thus far demonstrated only modest cytotoxicity.²⁰¹

MoAb-based recombinant immunotoxins typically contain monovalent scFv or dsFv fragments (see Figure 2). This is not only for technical reasons (bacterial expression systems not allowing expression of whole IgG), but also since the small size is considered an important advantage for the penetration of solid tumors.^{202,203} Some potential disadvantages should be noted as well. The absence of IgG Fc-parts prevents potentially beneficial interactions with the patient's immune system (see Paragraph 1.5). Moreover, the smaller size results in an accelerated clearance that, together with the monovalent binding, may negatively affect tumor saturation. The limited choice of toxins may also be a disadvantage, as the intracellular routing can be suboptimal for the available toxins. Another concern is that the reduced size may result in an increased renal excretion associated with profound renal toxicity.²⁰⁴

The ongoing developments in genetic engineering are likely to solve most of these problems. An example forms the fusion, by various techniques, of Fv fragments into di- or multivalent molecules.²⁰⁵ Depending on the Fv fragments applied, these can be mono, bi or multi-specific. The resulting molecules often demonstrate improved pharmacokinetics and have affinities equalizing or surpassing that of whole IgG. The additional incorporation of constant parts of the IgG heavy chain results in so called minibodies possessing natural IgG effector functions. The phage display technology is another recent development having a major impact on the production of recombinant ITs. The expression of scFv or Fab' fragments on the surface of filamentous phage enables the isolation of specific MoAb fragments from huge phage-display libraries, thus circumventing the conventional immunization and hybridoma technology. Libraries derived from human donor tissue (*e.g.* PBLs, spleen, tonsil or bone marrow) provide fully human antibody fragments of the desired specificity. These fragments can readily be engineered into the preferred form and spliced to an appropriate toxin-encoding gene. Considering the rapid developments within this area, recombinant production of ITs will undoubtedly become the future standard, surpassing conventional biochemical techniques in rapidly generating homogeneous batches with great flexibility regarding the form of ligand and toxin. Moreover, the greatest promise is still to be fulfilled, the generation of entirely human constructs comprising a human ligand fused to a human enzyme equalizing the conventional toxins in efficacy.

1.5 NON-TOXIN-BASED EFFECTOR MECHANISMS

Apart from its role as toxin transporter, the MoAb moiety contains intrinsic effector mechanisms that may significantly contribute to the overall therapeutic efficacy. These include the following:

The MoAb Fc-part

The MoAb Fc-part may fix complement C1q or bind to Fc receptors expressed on various immune cells.²⁰⁶ Triggering the complement cascade results in direct complement-dependent cytotoxicity (CDC), as well as in the activation of several immune effector cells. Binding of Fc receptors initiates a diversity of biological responses as well, including antibody dependent cellular cytotoxicity (ADCC), phagocytosis, endocytosis, release of inflammatory mediators, and enhancement of antigen presentation.

Though the potential anti-tumor effect of these mechanisms has been clearly demonstrated in xenografted tumor mouse models, the full clinical relevance has to be determined yet.²⁰⁷⁻²⁰⁹ Several studies report of complement deposition or mononuclear infiltrates in post-treatment biopsies of cancer patients treated with unconjugated (murine) MoAbs.^{210,211} In a comparative clinical trial of two CAMPATH (CDw52) rat isotype-switch variants, the most effective target cell depletion was observed with the

ADCC-inducing IgG2b MoAb.²¹² Moreover, CDC and ADCC are both considered *in vivo* effector mechanisms of chimeric anti-CD20 MoAbs in the treatment of B cell malignancies.²¹³⁻²¹⁸

Interference with cellular functions

The binding of the MoAb itself may interfere with cellular functions. The MoAb may block or mimic the binding of natural ligand. This may directly induce growth arrest or apoptosis, or the cell may be deprived of a critical survival/proliferation signal.

Members of the type I growth factor receptor family, including epidermal growth factor (EGF) receptor (EGFR, erbB1) and HER-2 (erbB2), have been implicated in the cellular transformation of several human malignancies.²¹⁹⁻²²³ The antiproliferative effect of EGFR and HER-2 blocking MoAbs has been demonstrated in several preclinical studies, including xenografted mouse tumor models.²²⁴⁻²²⁸ Moreover, a humanized version of a HER-2 blocking MoAb induced objective clinical responses in patients with breast cancer.¹³ Currently, combinations of anti-EGFR and anti-HER-2 MoAbs with other therapeutic agents are clinically evaluated as accumulating data suggest these MoAbs enhance the response to certain chemo- and hormone therapies.²²⁹⁻²³¹

Humanized anti-CD20 MoAb Rituximab forms an example of a therapeutic MoAb which clinical efficacy might partly be mediated by apoptosis induction. CD20 is a signaling protein involved in the regulation of B-cell proliferation, and is expressed on nearly all normal and malignant mature B cells. Ligation of CD20 by several MoAbs induces cell death by apoptosis.²³²⁻²³⁶ In addition to ADCC and CDC, this may contribute to the impressive responses observed in the treatment of CD20-positive neoplasms with unconjugated CD20 MoAbs.²³⁷⁻²⁴⁰

Anti-idiotypic responses

The MoAb's immunogenicity is a major obstacle potentially preventing multiple cycles of therapy. The reverse side is that the humoral responses mounted by the patient may elicit host anti-target cell antibodies as well. According to the 'idiotypic-network', the administered MoAb (Ab1) may induce host anti-idiotypic antibodies (Ab2), which in turn may provoke host anti-anti-idiotypic antibodies (Ab3), which in part bind the antigen of the Ab1 MoAb.²⁴¹ Indirect support for this concept was provided by the treatment of colon carcinoma with murine MoAb 17-1A. The presence of Ab3 antibodies appeared to be a positive prognostic factor for prolonged survival.²⁴²⁻²⁴⁸ Other studies have been entirely based on exploiting the anti-idiotypic strategy. In these studies, patients were administered Ab2 antibodies, mounted in animals immunized with a tumor-specific Ab1 MoAb (e.g. CO17-1A, HMW-MAA, CEA). Together with the formation of Ab3, several clinical responses could be observed, providing the proof of principle.²⁴⁹⁻²⁵⁴

Relevance of MoAb-based effector mechanisms for IT efficacy

There are several reports which confirm that the above effector mechanisms may contribute to the IT efficacy as well.

Hertler *et al.*, for instance, observed that an anti-CD5 IT (T101-ricin A) induced a rapid but transient fall in white blood cells of patients with chronic lymphocytic leukemia.²⁵⁵ This response was attributed to an interaction of the MoAb-moiety with the reticuloendothelial system.^{143,255} Yokota *et al.* demonstrated in a mouse tumor-xenograft model that the interaction with macrophages enhanced the anti-tumor efficacy of an IT-combination.^{42,256} More recently, Flavell *et al.* demonstrated in a mouse model that part of the anti-leukemia effect of a CD7-saporin IT was delivered by ADCC through recruitment of NK cells.²⁵⁷ Vitteta *et al.* provided an example of the MoAb-binding contributing to the IT efficacy. They demonstrated that the cytotoxicity of an anti-CD19 IT was partly based on the growth inhibitory signal induced by binding of the CD19 antigen.^{258,259} This binding also impairs the activity of the drug efflux pump P-glycoprotein (P-gp), leaving target cells more susceptible for chemotherapy.²⁶⁰ For anti-CD3 ITs, it has been demonstrated that binding of the CD3/T cell receptor complex (CD3/TCR) attributes to the IT efficacy by inducing activation induced cell death (AICD).²⁶¹⁻²⁶³ Moreover, when treating immune disorder, the binding of the CD3/TCR may block antigen stimulation.²⁶³ Similarly, targeting activated T-cells with anti-CD25 ITs may decrease ongoing activation by blocking the IL-2 signaling.

1.6 LESSONS FROM CLINICAL TRIALS

A variety of ITs has been tested in the clinic for diverse indications. Both recombinant and chemical conjugated ITs have been tested, administered by bolus as well as continuous infusion schedules. Most of these studies have been Phase I/II trials, designed to determine the dose limiting toxicity, maximal tolerable dose, and to gather information regarding biological and clinical efficacy. Due to the nature of these studies, it is too early to draw definite conclusions regarding the administration schedule or type of IT which are eventually to be preferred for *in vivo* use. Nevertheless, some interesting trends have been observed allowing the following general conclusions.

The observed toxicities can roughly be divided in 'general toxicities' directly mediated by the toxin moiety, and 'specific toxicities' due to binding of the cell targeting ligand to bystander tissue. An example of a general toxicity is vascular leak syndrome (VLS), characterized by hypoalbuminaemia, weight gain, peripheral oedema, pleural effusion and, occasionally, life threatening pulmonary oedema. Though observed with all applied bacterial and plant toxins, dose limiting severe VLS appears to be predominantly associated with ricin (A)-based ITs. Generally, VLS is transient and ceases shortly after discontinuation of IT administration. The second dose limiting toxicity reported for dgA-based

ITs is myalgia, rarely associated with rhabdomyolysis. Reversible elevations of serum liver enzymes have frequently been reported for ITs constructed with DT, PE, type I RIP, and blocked ricin.²⁶⁴⁻²⁶⁶ Hepatotoxicity has not been associated with dgA-ITs.

An example of specific toxicity has been reported by Pai *et al.* They observed an unexpected and severe central nervous system toxicity when clinically evaluating an IT (OVB₃-PE) reactive with human ovarian cancer. The underlying cross-reactivity with normal human brain tissue was not detected during preclinical screening. Similarly, an IT reactive with breast cancer (260F₉-RTA) caused severe sensorimotor neuropathy due to unexpected reactivity with Schwann cells. To predict and to anticipate on specific toxicities, ITs are to be extensively screened pre-clinically using a broad panel of human tissues from multiple donors. Nevertheless, MoAb-antigen interactions might be missed due to tissue fixation altering the antigenicity or destroying certain epitopes. The same holds true for the opposite, epitopes may be exposed that are hidden, or only marginally accessible, under normal physical conditions (*e.g.* intracellular antigens or antigens protected by the blood-brain barrier). Cross-reactivity observed *in vitro* should, therefore, not automatically eliminate a MoAb from consideration.^{27,267} In general, cross-reactivity observed during preclinical screening asks for additional studies in relevant test animals.

A common finding among IT trials is the occurrence of humoral responses to either the cell targeting ligand or the toxin component. For patients with hematological disorders the incidence of humoral anti-IT responses is typically in the order of 20-30%.^{99,263,268-273} The immunogenicity rate seen in ITs trials against solid tumors, in contrast, is often 90 to 100%.^{21,29} Many of these responses can be detected within 10 days of the first administration. Though hypersensitivity reactions are rare, repeated dosages may be less effective due to accelerated plasma clearance or the formation of inactive immune-complexes. Notably, extensive clinical responses have been seen even in the face of anti-IT antibodies.^{25,274,275}

The accessibility of target cells is a major factor determining the IT-efficacy. So far, solid tumors are treated with only limited success. This may not only be due to heterogeneous blood supply, but also to tight junctions between tumor cells and a high interstitial pressure. Though tumor penetration of small recombinant toxins is more effective, this appears to be counterbalanced by their rapid clearance by the kidney. The $T_{1/2}$ of the recombinant conjugates is often less than 1 hour, as compared to 6 or more hours for whole IgG-containing ITs. The treatment of diffuse hematological and immunological disorders has proved more successful, with overall clinical response rates ranging from 12-75%.²⁷⁶ Though many of these responses were partial and often transient, this is not such a bad score considering the early nature of most studies (mostly Phase I or II trials). So far, one immunotoxin has actually passed the registration process. The FDA approved recombinant toxin DAB₃89IL₂ (Ontak) for the treat-

ment of patients with advanced or recurrent cutaneous T-cell lymphoma.

Though responses have been obtained at safe dose levels, some important issues have to be addressed in order for ITs to become broadly effective therapeutics. Major challenges include the prevention of severe toxin-induced vascular leakage, the reduction of immunogenicity, and improving on the penetration of solid tumors. The optimal clinical benefit is likely to be obtained in an adjuvant setting, following conventional 'target-cell-debulking' therapy. Simultaneous application of multiple ITs may be necessary to solve the problem of heterogeneous antigen expression.

REFERENCES

1. Boltansky H, Slater J, Youle R, Isersky C, Kaliner M: IgE-immunotoxins. II. IgE-ricin A-chain. *Immunopharmacology* 14:47-62, 1987
2. Fishman A, Lorberboum-Galski H: Targeted elimination of cells expressing the high-affinity receptor for IgE (Fc epsilon RI) by a *Pseudomonas* exotoxin-based chimeric protein. *Eur J Immunol* 27:486-94, 1997
3. Fishman A, Prus D, Belostotsky R, Lorberboum-Galski H: Targeted Fc2'-3-PE40 chimeric protein abolishes passive cutaneous anaphylaxis in mice. *Clin Exp Immunol* 119:398-403, 2000
4. Miller RA, Maloney DG, Warnke R, Levy R: Treatment of B-cell lymphoma with monoclonal anti-idiotypic antibody. *N Engl J Med* 306:517-22, 1982
5. Garcia CF, Lowder J, Meeker TC, Bindl J, Levy R, Warnke RA: Differences in "host infiltrates" among lymphoma patients treated with anti-idiotypic antibodies: correlation with treatment response. *J Immunol* 135:4252-60, 1985
6. Meeker TC, Lowder J, Maloney DG, Miller RA, Thielemans K, Warnke R, Levy R: A clinical trial of anti-idiotypic therapy for B cell malignancy. *Blood* 65:1349-63, 1985
7. Lowder JN, Meeker TC, Campbell M, Garcia CF, Gralow J, Miller RA, Warnke R, Levy R: Studies on B lymphoid tumors treated with monoclonal anti-idiotypic antibodies: correlation with clinical responses. *Blood* 69:199-210, 1987
8. Allebes WA, Preijers FW, Haanen C, Capel PJ: The development of non-responsiveness to immunotherapy with monoclonal anti-idiotypic antibodies in a patient with B-CLL. *Br J Haematol* 70:295-300, 1988
9. Davis TA, Maloney DG, Czerwinski DK, Liles TM, Levy R: Anti-idiotypic antibodies can induce long-term complete remissions in non-Hodgkin's lymphoma without eradicating the malignant clone. *Blood* 92:1184-90, 1998
10. Lorimer IA, Wikstrand CJ, Batra SK, Bigner DD, Pastan I: Immunotoxins that target an oncogenic mutant epidermal growth factor receptor expressed in human tumors. *Clin Cancer Res* 1:859-64, 1995
11. Archer GE, Sampson JH, Lorimer IA, McLendon RE, Kuan CT, Friedman AH, Friedman HS, Pastan IH, Bigner DD: Regional treatment of epidermal growth factor receptor VIII-expressing neoplastic meningitis with a single-chain immunotoxin, MR-1. *Clin Cancer Res* 5:2646-52, 1999
12. Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL: Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science* 235:177-82, 1987
13. Baselga J, Tripathy D, Mendelsohn J, Baughman S, Benz CC, Dantis L, Sklarin NT, Seidman AD, Hudis CA, Moore J, Rosen PP, Twaddell T, Henderson IC, Norton L: Phase II study of weekly intravenous recombinant humanized anti-p185HER2 monoclonal antibody in patients with HER2/neu-overexpressing metastatic breast cancer. *J Clin Oncol* 14:737-44, 1996
14. Wels W, Harwerth IM, Mueller M, Groner B, Hynes NE: Selective inhibition of tumor cell growth by a recombinant single-chain antibody-toxin specific for the erbB-2 receptor. *Cancer Res* 52:6310-7, 1992
15. Di Lazzaro C, Digiesi G, Tecce R, Lotti LV, Torrisi MR, Natali PG: Immunotoxins to the HER-2 oncogene product: functional and ultrastructural analysis of their cytotoxic activity. *Cancer Immunol Immunother* 39:318-24, 1994
16. Xu FJ, Boyer CM, Bae DS, Wu S, Greenwald M, O'Brian K, Yu YH, Mills GB, Bast RC, Jr.: The tyrosine kinase activity of the C-erbB-2 gene product (p185) is required for growth inhibition by anti-p185 antibodies but not for the cytotoxicity of an anti-p185-ricin-A chain immunotoxin. *Int J Cancer* 59:242-7, 1994

17. Dean GS, Puzstai L, Xu FJ, O'Briant K, DeSombre K, Conaway M, Boyer CM, Mendelsohn J, Bast RC, Jr.: Cell surface density of p185(c-erbB-2) determines susceptibility to anti-p185(c-erbB-2)-ricin A chain (RTA) immunotoxin therapy alone and in combination with anti-p170(EGFR)-RTA in ovarian cancer cells. *Clin Cancer Res* 4:2545-50, 1998
18. Boyer CM, Puzstai L, Wiener JR, Xu FJ, Dean GS, Bast BS, O'Briant KC, Greenwald M, DeSombre KA, Bast RC, Jr.: Relative cytotoxic activity of immunotoxins reactive with different epitopes on the extracellular domain of the c-erbB-2 (HER-2/neu) gene product p185. *Int J Cancer* 82:525-31, 1999
19. Rosenblum MG, Shawver LK, Marks JW, Brink J, Cheung L, Langton-Webster B: Recombinant immunotoxins directed against the c-erb-2/HER2/neu oncogene product: in vitro cytotoxicity, pharmacokinetics, and in vivo efficacy studies in xenograft models. *Clin Cancer Res* 5:865-74, 1999
20. Pai-Scherf LH, Villa J, Pearson D, Watson T, Liu E, Willingham MC, Pastan I: Hepatotoxicity in cancer patients receiving erb-38, a recombinant immunotoxin that targets the erbB2 receptor. *Clin Cancer Res* 5:2311-5, 1999
21. Pai LH, Wittes R, Setser A, Willingham MC, Pastan I: Treatment of advanced solid tumors with immunotoxin LMB-1: an antibody linked to *Pseudomonas* exotoxin. *Nat Med* 2:350-3, 1996
22. Hertler AA, Spitler LE, Frankel AE: Humoral immune response to a ricin A chain immunotoxin in patients with metastatic melanoma. *Cancer Drug Deliv* 4:245-53, 1987
23. Spitler LE, del Rio M, Khentigan A, Wedel NI, Brophy NA, Miller LL, Harkonen WS, Rosendorf LL, Lee HM, Mischak RP, et al.: Therapy of patients with malignant melanoma using a monoclonal antimelanoma antibody-ricin A chain immunotoxin. *Cancer Res* 47:1717-23, 1987
24. Spitler LE, Mischak R, Scannon P: Therapy of metastatic malignant melanoma using Xomazyme Mel, a murine monoclonal anti-melanoma ricin A chain immunotoxin. *Int J Rad Appl Instrum [B]* 16:625-7, 1989
25. Oratz R, Speyer JL, Wernz JC, Hochster H, Meyers M, Mischak R, Spitler LE: Antimelanoma monoclonal antibody-ricin A chain immunoconjugate (XMMME-001-RTA) plus cyclophosphamide in the treatment of metastatic malignant melanoma: results of a phase II trial. *J Biol Response Mod* 9:345-54, 1990
26. Selvaggi K, Saria EA, Schwartz R, Vlock DR, Ackerman S, Wedel N, Kirkwood JM, Jones H, Ernstoff MS: Phase I/II study of murine monoclonal antibody-ricin A chain (XOMAZYME-Mel) immunoconjugate plus cyclosporine A in patients with metastatic melanoma. *J Immunother* 13:201-7, 1993
27. Lynch TJ, Jr.: Immunotoxin therapy of small-cell lung cancer. N901-blocked ricin for relapsed small-cell lung cancer. *Chest* 103:436S-439S, 1993
28. Epstein C, Lynch T, Shefner J, Wen P, Maxted D, Braman V, Ariniello P, Coral F, Ritz J: Use of the immunotoxin N901-blocked ricin in patients with small-cell lung cancer. *Int J Cancer Suppl* 8:57-9, 1994
29. Lynch TJ, Jr., Lambert JM, Coral F, Shefner J, Wen P, Blattler WA, Collinson AR, Ariniello PD, Braman G, Cook S, Esseltine D, Elias A, Skarin A, Ritz J: Immunotoxin therapy of small-cell lung cancer: a phase I study of N901-blocked ricin. *J Clin Oncol* 15:723-34, 1997
30. Zalberg JR, Pietersz G, Toohey B, Laird J, Huggins R, Zimet AS, Hennessy O, McKenzie A, McKenzie IF: A phase I/II study of the intralesional injection of ricin-monoclonal antibody conjugates in patients with hepatic metastases. *Eur J Cancer* 9:1227-31, 1994
31. Kuan CT, Pai LH, Pastan I: Immunotoxins containing *Pseudomonas* exotoxin that target LeY damage human endothelial cells in an antibody-specific mode: relevance to vascular leak syndrome. *Clin Cancer Res* 1:1589-94, 1995
32. Multani PS, Grossbard ML: Monoclonal antibody-based therapies for hematologic malignancies. *J Clin Oncol* 16:3691-710, 1998
33. Culig Z, Hittmair A, Hobisch A, Bartsch G, Klocker H, Pai LH, Pastan I: Expression of Lewis carbohydrate antigens in metastatic lesions from human prostatic carcinoma. *Prostate* 36:162-7, 1998

34. Starling JJ, Maciak RS, Law KL, Hinson NA, Briggs SL, Laguzza BC, Johnson DA: In vivo antitumor activity of a monoclonal antibody-Vinca alkaloid immunoconjugate directed against a solid tumor membrane antigen characterized by heterogeneous expression and noninternalization of antibody-antigen complexes. *Cancer Res* 51:2965-72, 1991
35. Colombatti M, Bisconti M, Lorenzi P, Stevanoni G, Dipasquale B, Gerosa M, Tridente G: Human glioma cell lines: tumour associated antigens distribution and sensitivity to antibody-toxin or ligand-toxin conjugates. A preliminary report. *Acta Neurochir Suppl* 43:121-5, 1988
36. Greiner JW, Ullmann CD, Nieroda C, Qi CF, Eggensperger D, Shimada S, Steinberg SM, Schlom J: Improved radioimmunotherapeutic efficacy of an anticarcinoma monoclonal antibody (131I-CC49) when given in combination with gamma-interferon. *Cancer Res* 53:600-8, 1993
37. Matsumoto H, Takahashi T, Mitsuhashi N, Higuchi K, Niibe H: Modification of tumor-associated antigen (CEA) expression of human lung cancer cells by irradiation, either alone or in combination with interferon-gamma. *Anticancer Res* 19:307-11, 1999
38. Meredith RF, Khazaeli MB, Macey DJ, Grizzle WE, Mayo M, Schlom J, Russell CD, LoBuglio AF: Phase II study of interferon-enhanced 131I-labeled high affinity CC49 monoclonal antibody therapy in patients with metastatic prostate cancer. *Clin Cancer Res* 5:3254s-3258s, 1999
39. Roselli M, Guadagni F, Buonomo O, Belardi A, Vittorini V, Mariani-Costantini R, Greiner JW, Casciani CU, Schlom J: Systemic administration of recombinant interferon alfa in carcinoma patients upregulates the expression of the carcinoma-associated antigens tumor-associated glycoprotein-72 and carcinoembryonic antigen. *J Clin Oncol* 14:2031-42, 1996
40. Yu YH, Crews JR, Cooper K, Ramakrishnan S, Houston LL, Leslie DS, George SL, Lidor Y, Boyer CM, Ring DB, et al.: Use of immunotoxins in combination to inhibit clonogenic growth of human breast carcinoma cells. *Cancer Res* 50:3231-8, 1990
41. Crews JR, Maier LA, Yu YH, Hester S, O'Brian K, Leslie DS, DeSombre K, George SL, Boyer CM, Argon Y, et al.: A combination of two immunotoxins exerts synergistic cytotoxic activity against human breast-cancer cell lines. *Int J Cancer* 51:772-9, 1992
42. Yokota S, Okazaki M, Yoshida M, Seon BK: Biodistribution and in vivo antitumor efficacy of the systemically administered anti-human T-leukemia immunotoxins and potentiation of their efficacy by alpha-interferon. *Leuk Res* 17:69-79, 1993
43. Engert A, Gottstein C, Bohlen H, Winkler U, Schon G, Manske O, Schnell R, Diehl V, Thorpe P: Cocktails of ricin A-chain immunotoxins against different antigens on Hodgkin and Sternberg-Reed cells have superior anti-tumor effects against H-RS cells in vitro and solid Hodgkin tumors in mice. *Int J Cancer* 63:304-9, 1995
44. Flavell DJ, Noss A, Pulford KA, Ling N, Flavell SU: Systemic therapy with 3BIT, a triple combination cocktail of anti-CD19, -CD22, and -CD38-saporin immunotoxins, is curative of human B-cell lymphoma in severe combined immunodeficient mice. *Cancer Res* 57:4824-9, 1997
45. Laurent G, Pris J, Farcet JP, Carayon P, Blythman H, Casellas P, Poncelet P, Jansen FK: Effects of therapy with T101 ricin A-chain immunotoxin in two leukemia patients. *Blood* 67:1680-7, 1986
46. Junghans RP, Carrasquillo JA, Waldmann TA: Impact of antigenemia on the bioactivity of infused anti-Tac antibody: implications for dose selection in antibody immunotherapies. *Proc Natl Acad Sci U S A* 95:1752-7, 1998
47. Burrows FJ, Thorpe PE: Eradication of large solid tumors in mice with an immunotoxin directed against tumor vasculature. *Proc Natl Acad Sci U S A* 90:8996-9000, 1993
48. Burrows FJ, Thorpe PE: Vascular targeting—a new approach to the therapy of solid tumors. *Pharmacol Ther* 64:155-74, 1994
49. Thorpe PE, Burrows FJ: Antibody-directed targeting of the vasculature of solid tumors [see comments]. *Breast Cancer Res Treat* 36:237-51, 1995

50. Burrows FJ, Derbyshire EJ, Tazzari PL, Amlot P, Gazdar AF, King SW, Letarte M, Vitetta ES, Thorpe PE: Up-regulation of endoglin on vascular endothelial cells in human solid tumors: implications for diagnosis and therapy. *Clin Cancer Res* 1:1623-34, 1995
51. Seon BK, Matsuno F, Haruta Y, Kondo M, Barcos M: Long-lasting complete inhibition of human solid tumors in SCID mice by targeting endothelial cells of tumor vasculature with antihuman endoglin immunotoxin. *Clin Cancer Res* 3:1031-44, 1997
52. Matsuno F, Haruta Y, Kondo M, Tsai H, Barcos M, Seon BK: Induction of lasting complete regression of preformed distinct solid tumors by targeting the tumor vasculature using two new anti-endoglin monoclonal antibodies. *Clin Cancer Res* 5:371-82, 1999
53. Huang X, Molema G, King S, Watkins L, Edgington TS, Thorpe PE: Tumor infarction in mice by antibody-directed targeting of tissue factor to tumor vasculature [see comments]. *Science* 275:547-50, 1997
54. Brekken RA, Huang X, King SW, Thorpe PE: Vascular endothelial growth factor as a marker of tumor endothelium. *Cancer Res* 58:1952-9, 1998
55. Ran S, Gao B, Duffy S, Watkins L, Rote N, Thorpe PE: Infarction of solid Hodgkin's tumors in mice by antibody-directed targeting of tissue factor to tumor vasculature. *Cancer Res* 58:4646-53, 1998
56. Preijers FW, Tax WJ, De Witte T, Janssen A, v.d. Heijden H, Vidal H, Wessels JM, Capel PJ: Relationship between internalization and cytotoxicity of ricin A-chain immunotoxins. *Br J Haematol* 70:289-94, 1988
57. Carriere D, Arcier JM, Derocq JM, Fontaine C, Richer G: Antigenic modulation induced by four monoclonal antibodies adsorbed on gold particles (specificity anti-CD4, anti-CD5, anti-CD7, and anti-150- kDa antigen): relationship between modulation and cytotoxic activity of immunotoxins. *Exp Cell Res* 182:114-28, 1989
58. Chignola R, Colombatti M, Dell'Arciprete L, Candiani C, Tridente G: Distribution of endocytosed molecules to intracellular acidic environments correlates with immunotoxin activity. *Int J Cancer* 46:117-23, 1990
59. May RD, Wheeler HT, Finkelman FD, Uhr JW, Vitetta ES: Intracellular routing rather than cross-linking or rate of internalization determines the potency of immunotoxins directed against different epitopes of sIgD on murine B cells. *Cell Immunol* 135:490-500, 1991
60. Press OW, Vitetta ES, Farr AG, Hansen JA, Martin PJ: Evaluation of ricin A-chain immunotoxins directed against human T cells. *Cell Immunol* 102:10-20, 1986
61. Embleton MJ, Byers VS, Lee HM, Scannon P, Blackhall NW, Baldwin RW: Sensitivity and selectivity of ricin toxin A chain-monoclonal antibody 791T/36 conjugates against human tumor cell lines. *Cancer Res* 46:524-8, 1986
62. Press OW, Martin PJ, Thorpe PE, Vitetta ES: Ricin A-chain containing immunotoxins directed against different epitopes on the CD2 molecule differ in their ability to kill normal and malignant T cells. *J Immunol* 141:4410-7, 1988
63. Luo Y, Seon BK: Marked difference in the in vivo antitumor efficacy between two immunotoxins targeted to different epitopes of common acute lymphoblastic leukemia antigen (CD10). Mechanisms involved in the differential activities of immunotoxins. *J Immunol* 145:1974-82, 1990
64. May RD, Finkelman FD, Wheeler HT, Uhr JW, Vitetta ES: Evaluation of ricin A chain-containing immunotoxins directed against different epitopes on the delta-chain of cell surface-associated IgD on murine B cells. *J Immunol* 144:3637-42, 1990
65. Derocq JM, Casellas P, Laurent G, Ravel S, Vidal H, Jansen F: Comparison of the cytotoxic potency of Tr101 Fab, F(ab')₂ and whole IgG immunotoxins. *J Immunol* 141:2837-43, 1988

66. Ghetie MA, Richardson J, Tucker T, Jones D, Uhr JW, Vitetta ES: Antitumor activity of Fab' and IgG-anti-CD22 immunotoxins in disseminated human B lymphoma grown in mice with severe combined immunodeficiency disease: effect on tumor cells in extranodal sites. *Cancer Res* 51:5876-80, 1991
67. Better M, Bernhard SL, Lei SP, Fishwild DM, Lane JA, Carroll SF, Horwitz AH: Potent anti-CD5 ricin A chain immunoconjugates from bacterially produced Fab' and F(ab')₂. *Proc Natl Acad Sci U S A* 90:457-61, 1993
68. van Oosterhout YV, Preijers FW, Wessels HM, de Witte T: Cytotoxicity of CD3-ricin A chain immunotoxins in relation to cellular uptake and degradation kinetics. *Cancer Res* 52:5921-5, 1992
69. van Horssen PJ, van Oosterhout YV, de Witte T, Preijers FW: Cytotoxic potency of CD22-ricin A depends on intracellular routing rather than on the number of internalized molecules. *Scand J Immunol* 41:563-9, 1995
70. Shan D, Press OW: Constitutive endocytosis and degradation of CD22 by human B cells. *J Immunol* 154:4466-75, 1995
71. Press OW, Howell-Clark J, Anderson S, Bernstein I: Retention of B-cell-specific monoclonal antibodies by human lymphoma cells. *Blood* 83:1390-7, 1994
72. Fulton RJ, Uhr JW, Vitetta ES: The effect of antibody valency and lysosomotropic amines on the synergy between ricin A chain- and ricin B chain-containing immunotoxins. *J Immunol* 136:3103-9, 1986
73. Siegall CB, Gawlak SL, Chin JJ, Zoeckler ME, Kadow KF, Brown JP, Braslawsky GR: Cytotoxicity of chimeric (human-murine) monoclonal antibody BR96 IgG, F(ab')₂, and Fab' conjugated to *Pseudomonas* exotoxin. *Bioconjug Chem* 3:302-7, 1992
74. Faguet GB, Agee JF: Four ricin chain A-based immunotoxins directed against the common chronic lymphocytic leukemia antigen: in vitro characterization. *Blood* 82:536-43, 1993
75. Woo BH, Lee JT, Lee KC: Ricin A immunotoxins of IgG and Fab of anti-CALLA monoclonal antibody: effect of water soluble long-chain SPDP on conjugate yield, immunoselectivity and cytotoxicity. *Arch Pharm Res* 17:452-7, 1994
76. Chiron M, Derocq JM, Kuhlein E, Roubinet F, Demur C, Grandjean H, Huguet F, Jaffrezou JP, Jansen F, Laurent G: Sensitivity of fresh leukemic cells to T101 ricin A-chain immunotoxin: a comparative study between Fab fragment and whole Ig conjugates. *Leuk Res* 13:491-9, 1989
77. van Oosterhout YV, van den Herik-Oudijk IE, Wessels HM, de Witte T, van de Winkel JG, Preijers FW: Effect of isotype on internalization and cytotoxicity of CD19-ricin A immunotoxins. *Cancer Res* 54:3527-32, 1994
78. Yazdi PT, Wenning LA, Murphy RM: Influence of cellular trafficking on protein synthesis inhibition of immunotoxins directed against the transferrin receptor. *Cancer Res* 55:3763-71, 1995
79. Battelli MG, Bolognesi A, Olivieri F, Polito L, Stirpe F: Different sensitivity of CD30+ cell lines to Ber-H2/saporin-S6 immunotoxin. *J Drug Target* 5:181-91, 1998
80. Ghetie MA, May RD, Till M, Uhr JW, Ghetie V, Knowles PP, Relf M, Brown A, Wallace PM, Janossy G, et al.: Evaluation of ricin A chain-containing immunotoxins directed against CD19 and CD22 antigens on normal and malignant human B-cells as potential reagents for in vivo therapy. *Cancer Res* 48:2610-7, 1988
81. Ghetie V, Engert A, Schnell R, Vitetta ES: The in vivo anti-tumor activity of immunotoxins containing two versus one deglycosylated ricin A chains. *Cancer Lett* 98:97-101, 1995
82. Bonardi MA, French RR, Amlot P, Gromo G, Modena D, Glennie MJ: Delivery of saporin to human B-cell lymphoma using bispecific antibody: targeting via CD22 but not CD19, CD37, or immunoglobulin results in efficient killing. *Cancer Res* 53:3015-21, 1993
83. Kreitman RJ, Pastan I: Recombinant toxins containing human granulocyte-macrophage colony-stimulating factor and either *pseudomonas* exotoxin or diphtheria toxin kill gastrointestinal cancer and leukemia cells. *Blood* 90:252-9, 1997

84. Till M, May RD, Uhr JW, Thorpe PE, Vitetta ES: An assay that predicts the ability of monoclonal antibodies to form potent ricin A chain-containing immunotoxins. *Cancer Res* 48:1119-23, 1988
85. Weltman JK, Pedroso P, Johnson SA, Davignon D, Fast LD, Leone LA: Rapid screening with indirect immunotoxin for monoclonal antibodies against human small cell lung cancer. *Cancer Res* 47:5552-6, 1987
86. Derbyshire EJ, Wawrzynczak EJ: Monoclonal antibodies recognising the cluster 2 antigen associated with human small cell lung cancer mediate the toxic effects of ricin A chain in an indirect assay of immunotoxin cytotoxicity. *Br J Cancer Suppl* 14:74-7, 1991
87. Derbyshire EJ, de Leij L, Wawrzynczak EJ: Refinement of an indirect immunotoxin assay of monoclonal antibodies recognising the human small cell lung cancer cluster 2 antigen. *Br J Cancer* 67:1242-7, 1993
88. Wilson BA, Collier RJ: Diphtheria toxin and *Pseudomonas aeruginosa* exotoxin A: active-site structure and enzymic mechanism. *Curr Top Microbiol Immunol* 175:27-41, 1992
89. Collier RJ: Structure-activity relationships in diphtheria toxin and *Pseudomonas aeruginosa* exotoxin A. *Cancer Treat Res* 37:25-35, 1988
90. Fitzgerald D, Pastan I: *Pseudomonas* exotoxin and recombinant immunotoxins derived from it. *Ann N Y Acad Sci* 685:740-5, 1993
91. Ogata M, Chaudhary VK, Pastan I, Fitzgerald DJ: Processing of *Pseudomonas* exotoxin by a cellular protease results in the generation of a 37,000-Da toxin fragment that is translocated to the cytosol. *J Biol Chem* 265:20678-85, 1990
92. Chiron MF, Fryling CM, Fitzgerald DJ: Cleavage of *pseudomonas* exotoxin and diphtheria toxin by a furin-like enzyme prepared from beef liver. *J Biol Chem* 269:18167-76, 1994
93. Chaudhary VK, Jinno Y, Fitzgerald D, Pastan I: *Pseudomonas* exotoxin contains a specific sequence at the carboxyl terminus that is required for cytotoxicity. *Proc Natl Acad Sci U S A* 87:308-12, 1990
94. Kondo T, Fitzgerald D, Chaudhary VK, Adhya S, Pastan I: Activity of immunotoxins constructed with modified *Pseudomonas* exotoxin A lacking the cell recognition domain. *J Biol Chem* 263:9470-5, 1988
95. Batra JK, Fitzgerald D, Gately M, Chaudhary VK, Pastan I: Anti-Tac(Fv)-PE₄₀, a single chain antibody *Pseudomonas* fusion protein directed at interleukin 2 receptor bearing cells. *J Biol Chem* 265:15198-202, 1990
96. Batra JK, Fitzgerald DJ, Chaudhary VK, Pastan I: Single-chain immunotoxins directed at the human transferrin receptor containing *Pseudomonas* exotoxin A or diphtheria toxin: anti-TFR(Fv)-PE₄₀ and DT388-anti-TFR(Fv). *Mol Cell Biol* 11:2200-5, 1991
97. Haggerty HG, Warner WA, Comerkeski CR, Peden WM, Mezza LE, Damle BD, Siegall CB, Davidson TJ: BR96 sFv-PE₄₀ immunotoxin: nonclinical safety assessment. *Toxicol Pathol* 27:87-94, 1999
98. Benhar I, Pastan I: Characterization of B1(Fv)PE₃₈ and B1(dsFv)PE₃₈: single-chain and disulfide-stabilized Fv immunotoxins with increased activity that cause complete remissions of established human carcinoma xenografts in nude mice. *Clin Cancer Res* 1:1023-9, 1995
99. Kreitman RJ, Wilson WH, White JD, Stetler-Stevenson M, Jaffe ES, Giardina S, Waldmann TA, Pastan I: Phase I trial of recombinant immunotoxin anti-Tac(Fv)-PE₃₈ (LMB-2) in patients with hematologic malignancies. *J Clin Oncol* 18:1622-36, 2000
100. Brinkmann U, Pai LH, Fitzgerald DJ, Willingham M, Pastan I: B3(Fv)-PE₃₈KDEL, a single-chain immunotoxin that causes complete regression of a human carcinoma in mice. *Proc Natl Acad Sci U S A* 88:8616-20, 1991
101. Kreitman RJ, Margulies I, Stetler-Stevenson M, Wang QC, Fitzgerald DJ, Pastan I: Cytotoxic activity of disulfide-stabilized recombinant immunotoxin RFB4(dsFv)-PE₃₈ (BL22) toward fresh malignant cells from patients with B-cell leukemias. *Clin Cancer Res* 6:1476-87, 2000

102. Robbins DH, Margulies I, Stetler-Stevenson M, Kreitman RJ: Hairy cell leukemia, a B-cell neoplasm that is particularly sensitive to the cytotoxic effect of anti-Tac(Fv)-PE38 (LMB-2). *Clin Cancer Res* 6:693-700, 2000
103. Kreitman RJ, Wilson WH, Robbins D, Margulies I, Stetler-Stevenson M, Waldmann TA, Pastan I: Responses in refractory hairy cell leukemia to a recombinant immunotoxin. *Blood* 94:3340-8, 1999
104. Essand M, Pastan I: Anti-prostate immunotoxins: cytotoxicity of E4 antibody-Pseudomonas exotoxin constructs. *Int J Cancer* 77:123-7, 1998
105. Mansfield E, Pastan I, FitzGerald DJ: Characterization of RFB4-Pseudomonas exotoxin A immunotoxins targeted to CD22 on B-cell malignancies. *Bioconjug Chem* 7:557-63, 1996
106. Seetharam S, Chaudhary VK, FitzGerald D, Pastan I: Increased cytotoxic activity of Pseudomonas exotoxin and two chimeric toxins ending in KDEL. *J Biol Chem* 266:17376-81, 1991
107. Kreitman RJ, Pastan I: Importance of the glutamate residue of KDEL in increasing the cytotoxicity of Pseudomonas exotoxin derivatives and for increased binding to the KDEL receptor. *Biochem J* 307:29-37, 1995
108. Proby CM, Ota T, Suzuki H, Koyasu S, Gamou S, Shimizu N, Wahl JK, Wheelock MJ, Nishikawa T, Amagai M: Development of chimeric molecules for recognition and targeting of antigen-specific B cells in pemphigus vulgaris [see comments]. *Br J Dermatol* 142:321-30, 2000
109. Kreitman RJ, Chaudhary VK, Kozak RW, FitzGerald DJ, Waldman TA, Pastan I: Recombinant toxins containing the variable domains of the anti-Tac monoclonal antibody to the interleukin-2 receptor kill malignant cells from patients with chronic lymphocytic leukemia. *Blood* 80:2344-52, 1992
110. Choe S, Bennett MJ, Fujii G, Curmi PM, Kantardjieff KA, Collier RJ, Eisenberg D: The crystal structure of diphtheria toxin. *Nature* 357:216-22, 1992
111. O'Keefe DO, Cabiaux V, Choe S, Eisenberg D, Collier RJ: pH-dependent insertion of proteins into membranes: B-chain mutation of diphtheria toxin that inhibits membrane translocation, Glu-349—Lys. *Proc Natl Acad Sci U S A* 89:6202-6, 1992
112. vanderSpek J, Cassidy D, Genbauffe F, Huynh PD, Murphy JR: An intact transmembrane helix 9 is essential for the efficient delivery of the diphtheria toxin catalytic domain to the cytosol of target cells. *J Biol Chem* 269:21455-9, 1994
113. Greenfield L, Johnson VG, Youle RJ: Mutations in diphtheria toxin separate binding from entry and amplify immunotoxin selectivity. *Science* 238:536-9, 1987
114. Neville DM, Jr., Srinivasachar K, Stone R, Scharff J: Enhancement of immunotoxin efficacy by acid-cleavable cross-linking agents utilizing diphtheria toxin and toxin mutants. *J Biol Chem* 264:14653-61, 1989
115. Lakkis F, Steele A, Pacheco-Silva A, Rubin-Kelley V, Strom TB, Murphy JR: Interleukin 4 receptor targeted cytotoxicity: genetic construction and in vivo immunosuppressive activity of a diphtheria toxin-related murine interleukin 4 fusion protein. *Eur J Immunol* 21:2253-8, 1991
116. Williams DP, Snider CE, Strom TB, Murphy JR: Structure/function analysis of interleukin-2-toxin (DAB486-IL-2). Fragment B sequences required for the delivery of fragment A to the cytosol of target cells. *J Biol Chem* 265:11885-9, 1990
117. Chaudhary VK, Gallo MG, FitzGerald DJ, Pastan I: A recombinant single-chain immunotoxin composed of anti-Tac variable regions and a truncated diphtheria toxin. *Proc Natl Acad Sci U S A* 87:9491-4, 1990
118. Thompson J, Hu H, Scharff J, Neville DM, Jr.: An anti-CD3 single-chain immunotoxin with a truncated diphtheria toxin avoids inhibition by pre-existing antibodies in human blood. *J Biol Chem* 270:28037-41, 1995
119. Barbieri L, Battelli MG, Stirpe F: Ribosome-inactivating proteins from plants. *Biochim Biophys Acta* 1154:237-82, 1993

120. Fu T, Burbage C, Tagge EP, Brothers T, Willingham MC, Frankel AE: Ricin toxin contains three lectin sites which contribute to its *in vivo* toxicity. *Int J Immunopharmacol* 18:685-92, 1996
121. Sandvig K, van Deurs B: Endocytosis, intracellular transport, and cytotoxic action of Shiga toxin and ricin. *Physiol Rev* 76:949-66, 1996
122. Manske JM, Buchsbaum DJ, Valleria DA: The role of ricin B chain in the intracellular trafficking of anti-CD5 immunotoxins. *J Immunol* 142:1755-66, 1989
123. Hudson TH, Grillo FG: Brefeldin-A enhancement of ricin A-chain immunotoxins and blockade of intact ricin, modeccin, and abrin. *J Biol Chem* 266:18586-92, 1991
124. Simpson JC, Dascher C, Roberts LM, Lord JM, Balch WE: Ricin cytotoxicity is sensitive to recycling between the endoplasmic reticulum and the Golgi complex. *J Biol Chem* 270:20078-83, 1995
125. Endo Y, Tsurugi K: The RNA N-glycosidase activity of ricin A-chain. The characteristics of the enzymatic activity of ricin A-chain with ribosomes and with rRNA. *J Biol Chem* 263:8735-9, 1988
126. Stong RC, Youle RJ, Valleria DA: Elimination of clonogenic T-leukemic cells from human bone marrow using anti-Mr 65,000 protein immunotoxins. *Cancer Res* 44:3000-6, 1984
127. Kaminski NE, Roberts JF, Guthrie FE: Target ricin by coupling to an anti-macrophage monoclonal antibody. *J Immunopharmacol* 8:15-37, 1986
128. Cattel L, Delprino L, Brusa P, Dosio F, Comoglio PM, Prat M: Comparison of blocked and non-blocked ricin-antibody immunotoxins against human gastric carcinoma and colorectal adenocarcinoma cell lines. *Cancer Immunol Immunother* 27:233-40, 1988
129. Goldmacher VS, Lambert JM, Blattler WA: The specific cytotoxicity of immunoconjugates containing blocked ricin is dependent on the residual binding capacity of blocked ricin: evidence that the membrane binding and A-chain translocation activities of ricin cannot be separated. *Biochem Biophys Res Commun* 183:758-66, 1992
130. Lambert JM, McIntyre G, Gauthier MN, Zullo D, Rao V, Steeves RM, Goldmacher VS, Blattler WA: The galactose-binding sites of the cytotoxic lectin ricin can be chemically blocked in high yield with reactive ligands prepared by chemical modification of glycopeptides containing triantennary N-linked oligosaccharides. *Biochemistry* 30:2324-47, 1991
131. Lambert JM, Goldmacher VS, Collinson AR, Nadler LM, Blattler WA: An immunotoxin prepared with blocked ricin: a natural plant toxin adapted for therapeutic use. *Cancer Res* 51:6236-42, 1991
132. Blythman HE, Casellas P, Gros O, Gros P, Jansen FK, Paolucci F, Pau B, Vidal H: Immunotoxins: hybrid molecules of monoclonal antibodies and a toxin subunit specifically kill tumour cells. *Nature* 290:145-6, 1981
133. Bourrie BJ, Casellas P, Blythman HE, Jansen FK: Study of the plasma clearance of antibody—ricin-A-chain immunotoxins. Evidence for specific recognition sites on the A chain that mediate rapid clearance of the immunotoxin. *Eur J Biochem* 155:1-10, 1986
134. Fulton RJ, Tucker TF, Vitetta ES, Uhr JW: Pharmacokinetics of tumor-reactive immunotoxins in tumor-bearing mice: effect of antibody valency and deglycosylation of the ricin A chain on clearance and tumor localization. *Cancer Res* 48:2618-25, 1988
135. Blakey DC, Watson GJ, Knowles PP, Thorpe PE: Effect of chemical deglycosylation of ricin A chain on the *in vivo* fate and cytotoxic activity of an immunotoxin composed of ricin A chain and anti-Thy 1.1 antibody. *Cancer Res* 47:947-52, 1987
136. Wawrzynczak EJ, Cumber AJ, Henry RV, Parnell GD: Comparative biochemical, cytotoxic and pharmacokinetic properties of immunotoxins made with native ricin A chain, ricin A1 chain and recombinant ricin A chain. *Int J Cancer* 47:130-5, 1991
137. Shire D, Bourrie BJ, Carillon C, Derocq JM, Dousset P, Dumont X, Jansen FK, Kaghad M, Legoux R, Lelong P, et al.: Biologically active A-chain of the plant toxin ricin expressed from a synthetic gene in *Escherichia coli*. *Gene* 93:183-8, 1990

138. Wawrzynczak EJ, Watson GJ, Cumber AJ, Henry RV, Parnell GD, Rieber EP, Thorpe PE: Blocked and non-blocked ricin immunotoxins against the CD4 antigen exhibit higher cytotoxic potency than a ricin A chain immunotoxin potentiated with ricin B chain or with a ricin B chain immunotoxin. *Cancer Immunol Immunother* 32:289-95, 1991
139. Timar J, McIntosh DP, Henry R, Cumber AJ, Parnell GD, Davies AJ: The effect of ricin B chain on the intracellular trafficking of an A chain immunotoxin. *Br J Cancer* 64:655-62, 1991
140. Bilge A, Howell-Clark J, Ramakrishnan S, Press OW: Degradation of ricin A chain by endosomal and lysosomal enzymes—the protective role of ricin B chain. *Ther Immunol* 1:197-204, 1994
141. Casellas P, Brown JP, Gros O, Gros P, Hellstrom I, Jansen FK, Poncelet P, Roncucci R, Vidal H, Hellstrom KE: Human melanoma cells can be killed in vitro by an immunotoxin specific for melanoma-associated antigen p97. *Int J Cancer* 30:437-43, 1982
142. Raso V, Lawrence J: Carboxylic ionophores enhance the cytotoxic potency of ligand- and antibody-delivered ricin A chain. *J Exp Med* 160:1234-40, 1984
143. Casellas P, Bourrie BJ, Gros P, Jansen FK: Kinetics of cytotoxicity induced by immunotoxins. Enhancement by lysosomotropic amines and carboxylic ionophores. *J Biol Chem* 259:9359-64, 1984
144. Casellas P, Jansen FK: Immunotoxin enhancers. *Cancer Treat Res* 37:351-69, 1988
145. Casellas P, Ravel S, Bourrie BJ, Derocq JM, Jansen FK, Laurent G, Gros P: T-lymphocyte killing by T101-ricin A-chain immunotoxin: pH-dependent potentiation with lysosomotropic amines. *Blood* 72:1197-202, 1988
146. Preijers FW, Tax WJ, Wessels JM, Capel PJ, De Witte T, Haanen C: Different susceptibilities of normal T cells and T cell lines to immunotoxins. *Scand J Immunol* 27:533-40, 1988
147. Press OW, DeSantes K, Anderson SK, Geissler F: Inhibition of catabolism of radiolabeled antibodies by tumor cells using lysosomotropic amines and carboxylic ionophores. *Cancer Res* 50:1243-50, 1990
148. Ravel S, Colombatti M, Casellas P: Internalization and intracellular fate of anti-CD5 monoclonal antibody and anti-CD5 ricin A-chain immunotoxin in human leukemic T cells. *Blood* 79:1511-7, 1992
149. van Horssen PJ, van Oosterhout YV, Evers S, Backus HH, van Oijen MG, Bongaerts R, de Witte T, Preijers FW: Influence of cytotoxicity enhancers in combination with human serum on the activity of CD22-recombinant ricin A against B cell lines, chronic and acute lymphocytic leukemia cells. *Leukemia* 13:241-9, 1999
150. Wu M: Enhancement of immunotoxin activity using chemical and biological reagents. *Br J Cancer* 75:1347-55, 1997
151. Casellas P, Canat X, Fauser AA, Gros O, Laurent G, Poncelet P, Jansen FK: Optimal elimination of leukemic T cells from human bone marrow with T101-ricin A-chain immunotoxin. *Blood* 65:289-97, 1985
152. Preijers FW, De Witte T, Wessels JM, De Gast GC, Van Leeuwen E, Capel PJ, Haanen C: Autologous transplantation of bone marrow purged in vitro with anti-CD7- (WT1-) ricin A immunotoxin in T-cell lymphoblastic leukemia and lymphoma. *Blood* 74:1152-8, 1989
153. Preijers FW, De Witte T, Wessels JM, Meyerink JP, Haanen C, Capel PJ: Cytotoxic potential of anti-CD7 immunotoxin (WT1-ricin A) to purge ex vivo malignant T cells in bone marrow. *Br J Haematol* 71:195-201, 1989
154. Myers DE, Jun X, Clementson D, Donelson R, Sicheneder A, Hoffman N, Bell K, Sarquis M, Langlie MC, Turner N, Uckun FM: Large scale manufacturing of TXU(anti-CD7)-pokeweed antiviral protein (PAP) immunoconjugate for clinical trials. *Leuk Lymphoma* 27:275-302, 1997

155. Myers DE, Irvin JD, Smith RS, Kuebelbeck VM, Uckun FM: Production of a pokeweed antiviral protein (PAP)-containing immunotoxin, B43-PAP, directed against the CD19 human B lineage lymphoid differentiation antigen in highly purified form for human clinical trials. *J Immunol Methods* 136:221-37, 1991
156. Irvin JD, Uckun FM: Pokeweed antiviral protein: ribosome inactivation and therapeutic applications. *Pharmacol Ther* 55:279-302, 1992
157. Thorpe PE, Brown AN, Bremner JA, Jr., Foxwell BM, Stirpe F: An immunotoxin composed of monoclonal anti-Thy 1.1 antibody and a ribosome-inactivating protein from *Saponaria officinalis*: potent antitumor effects in vitro and in vivo. *J Natl Cancer Inst* 75:151-9, 1985
158. Bolognesi A, Tazzari PL, Tassi C, Gromo G, Gobbi M, Stirpe F: A comparison of anti-lymphocyte immunotoxins containing different ribosome-inactivating proteins and antibodies. *Clin Exp Immunol* 89:341-6, 1992
159. Li BY, Ramakrishnan S: Recombinant hybrid toxin with dual enzymatic activities. Potential use in preparing highly effective immunotoxins. *J Biol Chem* 269:2652-8, 1994
160. Hoogenboom HR, Raus JC, Volckaert G: Targeting of tumor necrosis factor to tumor cells: secretion by myeloma cells of a genetically engineered antibody-tumor necrosis factor hybrid molecule. *Biochim Biophys Acta* 1096:345-54, 1991
161. Scherf U, Benhar I, Webber KO, Pastan I, Brinkmann U: Cytotoxic and antitumor activity of a recombinant tumor necrosis factor- B1(Fv) fusion protein on LeY antigen-expressing human cancer cells. *Clin Cancer Res* 2:1523-31, 1996
162. Gadina M, Newton DL, Rybak SM, Wu YN, Youle RJ: Humanized immunotoxins. *Ther Immunol* 1:59-64, 1994
163. Rosenblum MG, Cheung L, Mujoo K, Murray JL: An antimelanoma immunotoxin containing recombinant human tumor necrosis factor: tissue disposition, pharmacokinetic, and therapeutic studies in xenograft models. *Cancer Immunol Immunother* 40:322-8, 1995
164. Gho YS, Chae CB: Luteinizing hormone releasing hormone-RNase A conjugates specifically inhibit the proliferation of LHRH-receptor-positive human prostate and breast tumor cells. *Mol Cells* 9:31-6, 1999
165. Jinno H, Ueda M, Ozawa S, Kikuchi K, Ikeda T, Enomoto K, Kitajima M: Epidermal growth factor receptor-dependent cytotoxic effect by an EGF- ribonuclease conjugate on human cancer cell lines—a trial for less immunogenic chimeric toxin. *Cancer Chemother Pharmacol* 38:303-8, 1996
166. Psarras K, Ueda M, Yamamura T, Ozawa S, Kitajima M, Aiso S, Komatsu S, Seno M: Human pancreatic RNase-human epidermal growth factor fusion: an entirely human 'immunotoxin analog' with cytotoxic properties against squamous cell carcinomas. *Protein Eng* 11:1285-92, 1998
167. Yoon JM, Han SH, Kwon OB, Kim SH, Park MH, Kim BK: Cloning and cytotoxicity of fusion proteins of EGF and angiogenin. *Life Sci* 64:1435-45, 1999
168. Linardou H, Deonarain MP, Spooner RA, Epenetos AA: Deoxyribonuclease I (DNase I). A novel approach for targeted cancer therapy. *Cell Biophys* 25:243-8, 1994
169. Linardou H, Epenetos AA, Deonarain MP: A recombinant cytotoxic chimera based on mammalian deoxyribonuclease-I. *Int J Cancer* 86:561-9, 2000
170. Rybak SM, Saxena SK, Ackerman EJ, Youle RJ: Cytotoxic potential of ribonuclease and ribonuclease hybrid proteins. *J Biol Chem* 266:21202-7, 1991
171. Rybak SM, Hoogenboom HR, Newton DL, Raus JC, Youle RJ: Rational immunotherapy with ribonuclease chimeras. An approach toward humanizing immunotoxins. *Cell Biophys* 21:121-38, 1992
172. Rybak SM, Hoogenboom HR, Meade HM, Raus JC, Schwartz D, Youle RJ: Humanization of immunotoxins. *Proc Natl Acad Sci U S A* 89:3165-9, 1992

173. Apelgren LD, Bailey DL, Briggs SL, Barton RL, Guttman-Carlisle D, Koppel GA, Nichols CL, Scott WL, Lindstrom TD, Baker AL, et al.: Chemoimmunoconjugate development for ovarian carcinoma therapy: preclinical studies with vinca alkaloid-monoclonal antibody constructs. *Bioconjug Chem* 4:121-6, 1993
174. Suzuki S, Uno S, Fukuda Y, Aoki Y, Masuko T, Hashimoto Y: Cytotoxicity of anti-c-erbB-2 immunoliposomes containing doxorubicin on human cancer cells. *Br J Cancer* 72:663-8, 1995
175. Ding L, Samuel J, MacLean GD, Noujaim AA, Diener E, Longenecker BM: Effective drug-antibody targeting using a novel monoclonal antibody against the proliferative compartment of mammalian squamous carcinomas. *Cancer Immunol Immunother* 32:105-9, 1990
176. Biddle WC, Haruta Y, Seon BK, Henderson ES, Sarcione EJ: In vitro and in vivo cytotoxic activity of anti-human leukemia monoclonal antibodies SN5c and SN6 daunorubicin conjugates. *Leuk Res* 13:699-707, 1989
177. Dillman RO, Johnson DE, Shawler DL, Koziol JA: Superiority of an acid-labile daunorubicin-mono-clonal antibody immunoconjugate compared to free drug. *Cancer Res* 48:6097-102, 1988
178. Shawler DL, Johnson DE, Sweet MD, Myers LJ, Tudor SD, Beidler DE, Koziol JA, Dillman RO: Pre-clinical trials using an immunoconjugate of T101 and methotrexate in an athymic mouse/human T-cell tumor model. *J Biol Response Mod* 7:608-18, 1988
179. Adler R, Hurwitz E, Wands JR, Sela M, Shouval D: Specific targeting of adriamycin conjugates with monoclonal antibodies to hepatoma associated antigens to intrahepatic tumors in athymic mice. *Hepatology* 22:1482-7, 1995
180. Zhu Z, Kralovec J, Ghose T, Mammen M: Inhibition of Epstein-Barr-virus-transformed human chronic lymphocytic leukaemic B cells with monoclonal-antibody-adriamycin (doxorubicin) conju-gates. *Cancer Immunol Immunother* 40:257-67, 1995
181. Liu C, Tadayoni BM, Bourret LA, Mattocks KM, Derr SM, Widdison WC, Kedersha NL, Ariniello PD, Goldmacher VS, Lambert JM, Blattler WA, Chari RV: Eradication of large colon tumor xenografts by targeted delivery of maytansinoids. *Proc Natl Acad Sci U S A* 93:8618-23, 1996
182. Appelbaum FR: Antibody-targeted therapy for myeloid leukemia. *Semin Hematol* 36:2-8, 1999
183. Weil-Hillman G, Uckun FM, Manske JM, Vallera DA: Combined immunochemotherapy of human solid tumors in nude mice. *Cancer Res* 47:579-85, 1987
184. Pearson JW, Sivam G, Manger R, Wiltrout RH, Morgan AC, Jr., Longo DL: Enhanced therapeutic efficacy of an immunotoxin in combination with chemotherapy against an intraperitoneal human tumor xenograft in athymic mice. *Cancer Res* 49:4990-5, 1989
185. Griffin TW, Stocl M, Collins J, Fernandes J, Maher VE: Combined antitumor therapy with the chemotherapeutic drug doxorubicin and an anti-transferrin receptor immunotoxin: in vitro and in vivo studies. *J Immunother* 11:12-8, 1992
186. Ghetie MA, Tucker K, Richardson J, Uhr JW, Vitetta ES: Eradication of minimal disease in severe combined immunodeficient mice with disseminated Daudi lymphoma using chemotherapy and an immunotoxin cocktail. *Blood* 84:702-7, 1994
187. Scadden DT, Schenkein DP, Bernstein Z, Luskey B, Doweiko J, Tulpule A, Levine AM: Immuno-toxin combined with chemotherapy for patients with AIDS-related non-Hodgkin's lymphoma. *Cancer* 83:2580-7, 1998
188. Ek O, Gaynon P, Zeren T, Chelstrom LM, Myers DE, Uckun FM: Treatment of human B-cell pre-cursor leukemia in SCID mice by using a combination of the anti-CD19 immunotoxin B43-PAP with the standard chemotherapeutic drugs vincristine, methylprednisolone, and L- asparaginase. *Leuk Lymphoma* 31:143-9, 1998
189. Post J, Vooijs WC, Bast BJ, De Gast GC: Efficacy of an anti-CD138 immunotoxin and doxorubicin on drug-resistant and drug-sensitive myeloma cells. *Int J Cancer* 83:571-576, 1999

190. Masuho Y, Kishida K, Saito M, Umemoto N, Hara T: Importance of the antigen-binding valency and the nature of the cross-linking bond in ricin A-chain conjugates with antibody. *J Biochem (Tokyo)* 91:1583-91, 1982
191. Ramakrishnan S, Bjorn MJ, Houston LL: Recombinant ricin A chain conjugated to monoclonal antibodies: improved tumor cell inhibition in the presence of lysosomotropic compounds. *Cancer Res* 49:613-7, 1989
192. Wawrzynczak EJ, Thorpe PE: Effect of chemical linkage upon the stability and cytotoxic activity of A chain immunotoxins. *Cancer Treat Res* 37:239-51, 1988
193. Thorpe PE, Wallace PM, Knowles PP, Relf MG, Brown AN, Watson GJ, Knyba RE, Wawrzynczak EJ, Blakey DC: New coupling agents for the synthesis of immunotoxins containing a hindered disulfide bond with improved stability in vivo. *Cancer Res* 47:5924-31, 1987
194. Thorpe PE, Wallace PM, Knowles PP, Relf MG, Brown AN, Watson GJ, Blakey DC, Newell DR: Improved antitumor effects of immunotoxins prepared with deglycosylated ricin A-chain and hindered disulfide linkages. *Cancer Res* 48:6396-403, 1988
195. Stirpe F, Barbieri L, Tazzari PL, Dinota A, Gobbi M: Ex vivo bone marrow purging with immunotoxins. *Eur J Haematol Suppl* 51:173-5, 1989
196. Ghetie V, Swindell E, Uhr JW, Vitetta ES: Purification and properties of immunotoxins containing one vs. two deglycosylated ricin A chains. *J Immunol Methods* 166:117-22, 1993
197. Myers DE, Yanishevski Y, Masson E, Irvin JD, Evans WE, Uckun FM: Favorable pharmacodynamic features and superior anti-leukemic activity of B43 (anti-CD19) immunotoxins containing two poke-weed antiviral protein molecules covalently linked to each monoclonal antibody molecule. *Leuk Lymphoma* 18:93-102, 1995
198. Vallera DA, Burns LJ, Frankel AE, Sicheneder AR, Gunther R, Gajl-Peczalska K, Pennell CA, Kersey JH: Laboratory preparation of a deglycosylated ricin toxin A chain containing immunotoxin directed against a CD7 T lineage differentiation antigen for phase I human clinical studies involving T cell malignancies. *J Immunol Methods* 197:69-83, 1996
199. Flavell DJ, Boehm DA, Noss A, Flavell SU: Comparison of the potency and therapeutic efficacy of the anti-CD7 immunotoxin HB2-saporin constructed with one or two saporin moieties per immunotoxin molecule. *Br J Cancer* 75:1035-43, 1997
200. Grossbard ML, Lambert JM, Goldmacher VS, Blattler WA, Nadler LM: Correlation between in vivo toxicity and preclinical in vitro parameters for the immunotoxin anti-B4-blocked ricin. *Cancer Res* 52:4200-7, 1992
201. Westby M, Argent RH, Pitcher C, Lord JM, Roberts LM: Preparation and characterization of recombinant proricin containing an alternative protease-sensitive linker sequence. *Bioconjug Chem* 3:375-81, 1992
202. Fujimori K, Covell DG, Fletcher JE, Weinstein JN: Modeling analysis of the global and microscopic distribution of immunoglobulin G, F(ab')₂, and Fab in tumors. *Cancer Res* 49:5656-63, 1989
203. Debinski W, Pastan I: Recombinant F(ab')₂ C242-Pseudomonas exotoxin, but not the whole antibody-based immunotoxin, causes regression of a human colorectal tumor xenograft. *Clin Cancer Res* 1:1015-22, 1995
204. Vallera DA, Panoskaltsis-Mortari A, Blazar BR: Renal dysfunction accounts for the dose limiting toxicity of DT390anti-CD3sFv, a potential new recombinant anti-GVHD immunotoxin. *Protein Eng* 10:1071-6, 1997
205. Vallera DA, Kuroki DW, Panoskaltsis-Mortari A, Buchsbaum DJ, Rogers BE, Blazar BR: Molecular modification of a recombinant anti-CD3epsilon-directed immunotoxin by inducing terminal cysteine bridging enhances anti-GVHD efficacy and reduces organ toxicity in a lethal murine model. *Blood* 96:1157-65, 2000
206. Dillman RO: Antibodies as cytotoxic therapy. *J Clin Oncol* 12:1497-515, 1994

207. Adams DO, Hall T, Steplewski Z, Koprowski H: Tumors undergoing rejection induced by monoclonal antibodies of the IgG2a isotype contain increased numbers of macrophages activated for a distinctive form of antibody-dependent cytotoxicity. *Proc Natl Acad Sci U S A* 81:3506-10, 1984
208. Takahashi H, Nakada T, Nakaki M, Wands JR: Inhibition of hepatic metastases of human colon cancer in nude mice by a chimeric SF-25 monoclonal antibody. *Gastroenterology* 108:172-82, 1995
209. Denkers EY, Badger CC, Ledbetter JA, Bernstein ID: Influence of antibody isotype on passive serotherapy of lymphoma. *J Immunol* 135:2183-6, 1985
210. Houghton AN, Mintzer D, Cordon-Cardo C, Welt S, Fliegel B, Vadhan S, Carswell E, Melamed MR, Oettgen HF, Old LJ: Mouse monoclonal IgG3 antibody detecting GD3 ganglioside: a phase I trial in patients with malignant melanoma. *Proc Natl Acad Sci U S A* 82:1242-6, 1985
211. Goodman GE, Hellstrom I, Brodzinsky L, Nicaise C, Kulander B, Hummel D, Hellstrom KE: Phase I trial of murine monoclonal antibody L6 in breast, colon, ovarian, and lung cancer. *J Clin Oncol* 8:1083-92, 1990
212. Dyer MJ, Hale G, Hayhoe FG, Waldmann H: Effects of CAMPATH-1 antibodies in vivo in patients with lymphoid malignancies: influence of antibody isotype. *Blood* 73:1431-9, 1989
213. Grillo-Lopez AJ, White CA, Varns C, Shen D, Wei A, McClure A, Dallaire BK: Overview of the clinical development of rituximab: first monoclonal antibody approved for the treatment of lymphoma. *Semin Oncol* 26:66-73, 1999
214. Yang H, Rosove MH, Figlin RA: Tumor lysis syndrome occurring after the administration of rituximab in lymphoproliferative disorders: high-grade non-Hodgkin's lymphoma and chronic lymphocytic leukemia. *Am J Hematol* 62:247-50, 1999
215. Jensen M, Winkler U, Manzke O, Diehl V, Engert A: Rapid tumor lysis in a patient with B-cell chronic lymphocytic leukemia and lymphocytosis treated with an anti-CD20 monoclonal antibody (IDEC- C2B8, rituximab). *Ann Hematol* 77:89-91, 1998
216. Idusogie EE, Presta LG, Gazzano-Santoro H, Totpal K, Wong PY, Ullsch M, Meng YG, Mulkerrin MG: Mapping of the C1q binding site on rituxan, a chimeric antibody with a human IgG1 Fc. *J Immunol* 164:4178-84, 2000
217. Harjunpaa A, Junnikkala S, Meri S: Rituximab (anti-CD20) therapy of B-cell lymphomas: direct complement killing is superior to cellular effector mechanisms. *Scand J Immunol* 51:634-41, 2000
218. Golay J, Zaffaroni L, Vaccari T, Lazzari M, Borleri GM, Bernasconi S, Tedesco F, Rambaldi A, Introna M: Biologic response of B lymphoma cells to anti-CD20 monoclonal antibody rituximab in vitro: CD55 and CD59 regulate complement-mediated cell lysis. *Blood* 95:3900-8, 2000
219. Hynes NE, Stern DF: The biology of erbB-2/neu/HER-2 and its role in cancer. *Biochim Biophys Acta* 1198:165-84, 1994
220. Dougall WC, Qian X, Peterson NC, Miller MJ, Samanta A, Greene MI: The neu-oncogene: signal transduction pathways, transformation mechanisms and evolving therapies. *Oncogene* 9:2109-23, 1994
221. Di Fiore PP, Pierce JH, Fleming TP, Hazan R, Ullrich A, King CR, Schlessinger J, Aaronson SA: Overexpression of the human EGF receptor confers an EGF-dependent transformed phenotype to NIH 3T3 cells. *Cell* 51:1063-70, 1987
222. Wong ST, Winchell LF, McCune BK, Earp HS, Teixido J, Massague J, Herman B, Lee DC: The TGF- α precursor expressed on the cell surface binds to the EGF receptor on adjacent cells, leading to signal transduction. *Cell* 56:495-506, 1989
223. Velu TJ, Beguinot L, Vass WC, Willingham MC, Merlino GT, Pastan I, Lowy DR: Epidermal-growth-factor-dependent transformation by a human EGF receptor proto-oncogene. *Science* 238:1408-10, 1987
224. Kawamoto T, Sato JD, Le A, Polikoff J, Sato GH, Mendelsohn J: Growth stimulation of A431 cells by epidermal growth factor: identification of high-affinity receptors for epidermal growth factor by an anti-receptor monoclonal antibody. *Proc Natl Acad Sci U S A* 80:1337-41, 1983


225. Reiss M, Stash EB, Vellucci VF, Zhou ZL: Activation of the autocrine transforming growth factor alpha pathway in human squamous carcinoma cells. *Cancer Res* 51:6254-62, 1991
226. Fan Z, Masui H, Altas I, Mendelsohn J: Blockade of epidermal growth factor receptor function by bivalent and monovalent fragments of 225 anti-epidermal growth factor receptor monoclonal antibodies. *Cancer Res* 53:4322-8, 1993
227. Harwerth IM, Wels W, Schlegel J, Muller M, Hynes NE: Monoclonal antibodies directed to the erbB-2 receptor inhibit in vivo tumour cell growth. *Br J Cancer* 68:1140-5, 1993
228. Modjtahedi H, Eccles S, Box G, Styles J, Dean C: Immunotherapy of human tumour xenografts overexpressing the EGF receptor with rat antibodies that block growth factor-receptor interaction. *Br J Cancer* 67:254-61, 1993
229. Fan Z, Baselga J, Masui H, Mendelsohn J: Antitumor effect of anti-epidermal growth factor receptor monoclonal antibodies plus cis-diamminedichloroplatinum on well established A431 cell xenografts. *Cancer Res* 53:4637-42, 1993
230. Hancock MC, Langton BC, Chan T, Toy P, Monahan JJ, Mischak RP, Shawver LK: A monoclonal antibody against the c-erbB-2 protein enhances the cytotoxicity of cis-diamminedichloroplatinum against human breast and ovarian tumor cell lines. *Cancer Res* 51:4575-80, 1991
231. Yamauchi H, O'Neill A, Gelman R, Carney W, Tenney DY, Hosch S, Hayes DF: Prediction of response to antiestrogen therapy in advanced breast cancer patients by pretreatment circulating levels of extracellular domain of the HER-2/c-neu protein. *J Clin Oncol* 15:2518-25, 1997
232. Hofmeister JK, Cooney D, Coggeshall KM: Clustered CD20 Induced Apoptosis: Src-Family Kinase, the Proximal Regulator of Tyrosine Phosphorylation, Calcium Influx, and Caspase 3- Dependent Apoptosis. *Blood Cells Mol Dis* 26:133-143, 2000
233. Shan D, Ledbetter JA, Press OW: Apoptosis of malignant human B cells by ligation of CD20 with monoclonal antibodies. *Blood* 91:1644-52, 1998
234. Shan D, Ledbetter JA, Press OW: Signaling events involved in anti-CD20-induced apoptosis of malignant human B cells. *Cancer Immunol Immunother* 48:673-83, 2000
235. Winter JN: Prognostic indicators in diffuse, aggressive non-Hodgkin's lymphomas: regulators of growth fraction and programmed cell death. *Semin Oncol* 26:26-33, 1999
236. Taji H, Kagami Y, Okada Y, Andou M, Nishi Y, Saito H, Seto M, Morishima Y: Growth inhibition of CD20-positive B lymphoma cell lines by IDEC-C2B8 anti-CD20 monoclonal antibody. *Jpn J Cancer Res* 89:748-56, 1998
237. Feuring-Buske M, Buske C, Unterhalt M, Hiddemann W: Recent advances in antigen-targeted therapy in non-Hodgkin's lymphoma. *Ann Hematol* 79:167-74, 2000
238. Foran JM, Cunningham D, Coiffier B, Solal-Celigny P, Reyes F, Ghielmini M, Johnson PW, Gisselbrecht C, Bradburn M, Matthews J, Lister TA: Treatment of mantle-cell lymphoma with Rituximab (chimeric monoclonal anti-CD20 antibody): analysis of factors associated with response. *Ann Oncol* 11:117-21, 2000
239. Maloney DG: Preclinical and phase I and II trials of rituximab. *Semin Oncol* 26:74-8, 1999
240. McLaughlin P, Hagemeister FB, Grillo-Lopez AJ: Rituximab in indolent lymphoma: the single-agent pivotal trial. *Semin Oncol* 26:79-87, 1999
241. Jerne NK: Towards a network theory of the immune system. *Ann Immunol (Paris)* 125C:373-89, 1974
242. Frodin JE, Faxas ME, Hagstrom B, Lefvert AK, Masucci G, Nilsson B, Steinitz M, Unger P, Mellstedt H: Induction of anti-idiotypic (ab2) and anti-anti-idiotypic (ab3) antibodies in patients treated with the mouse monoclonal antibody 17-1A (ab1). Relation to the clinical outcome—an important antitumoral effector function? *Hybridoma* 10:421-31, 1991
243. Frodin JE, Lefvert AK, Mellstedt H: The clinical significance of HAMA in patients treated with mouse monoclonal antibodies. *Cell Biophys* 21:153-65, 1992

244. Gruber R, van Haarlem LJ, Warnaar SO, Holz E, Riethmuller G: The human antimouse immunoglobulin response and the anti-idiotypic network have no influence on clinical outcome in patients with minimal residual colorectal cancer treated with monoclonal antibody CO17-1A. *Cancer Res* 60:1921-6, 2000
245. Fagerberg J, Ragnhammar P, Liljefors M, Hjelm AL, Mellstedt H, Frodin JE: Humoral anti-idiotypic and anti-anti-idiotypic immune response in cancer patients treated with monoclonal antibody 17-1A. *Cancer Immunol Immunother* 42:81-7, 1996
246. Herlyn D, Harris D, Zaloudik J, Sperlagh M, Maruyama H, Jacob L, Kieny MP, Scheck S, Soma-sundaram R, Hart E, et al.: Immunomodulatory activity of monoclonal anti-idiotypic antibody to anti- colorectal carcinoma antibody CO17-1A in animals and patients. *J Immunother Emphasis Tumor Immunol* 15:303-11, 1994
247. Fagerberg J, Frodin JE, Ragnhammar P, Steinitz M, Wigzell H, Mellstedt H: Induction of an immune network cascade in cancer patients treated with monoclonal antibodies (ab1). II. Is induction of anti-idiotypic reactive T cells (T3) of importance for tumor response to mAb therapy? *Cancer Immunol Immunother* 38:149-59, 1994
248. Fagerberg J, Frodin JE, Wigzell H, Mellstedt H: Induction of an immune network cascade in cancer patients treated with monoclonal antibodies (ab1). I. May induction of ab1-reactive T cells and anti-anti-idiotypic antibodies (ab3) lead to tumor regression after mAb therapy? *Cancer Immunol Immunother* 37:264-70, 1993
249. Mittelman A, Chen ZJ, Kageshita T, Yang H, Yamada M, Baskind P, Goldberg N, Puccio C, Ahmed T, Arlin Z, et al.: Active specific immunotherapy in patients with melanoma. A clinical trial with mouse antiidiotypic monoclonal antibodies elicited with syngeneic anti-high-molecular-weight-melanoma-associated antigen monoclonal antibodies. *J Clin Invest* 86:2136-44, 1990
250. Mittelman A, Puccio C, Chun H, Ahmed T: Humoral antimelanoma immunity: induction by mouse antiidiotypic mAb, MK2-23 and association with survival of patients with advanced melanoma. *Mt Sinai J Med* 59:234-7, 1992
251. Mittelman A, Chen ZJ, Yang H, Wong GY, Ferrone S: Human high molecular weight melanoma-associated antigen (HMW-MAA) mimicry by mouse anti-idiotypic monoclonal antibody MK2-23: induction of humoral anti-HMW-MAA immunity and prolongation of survival in patients with stage IV melanoma. *Proc Natl Acad Sci U S A* 89:466-70, 1992
252. Herlyn D, Wettendorff M, Schmoll E, Iliopoulos D, Schedel I, Dreikhausen U, Raab R, Ross AH, Jaksche H, Scriba M, et al.: Anti-idiotypic immunization of cancer patients: modulation of the immune response. *Proc Natl Acad Sci U S A* 84:8055-9, 1987
253. Foon KA, Chakraborty M, John WJ, Sherratt A, Kohler H, Bhattacharya-Chatterjee M: Immune response to the carcinoembryonic antigen in patients treated with an anti-idiotypic antibody vaccine. *J Clin Invest* 96:334-42, 1995
254. Foon KA, John WJ, Chakraborty M, Sherratt A, Garrison J, Flett M, Bhattacharya-Chatterjee M: Clinical and immune responses in advanced colorectal cancer patients treated with anti-idiotypic monoclonal antibody vaccine that mimics the carcinoembryonic antigen. *Clin Cancer Res* 3:1267-76, 1997
255. Hertler AA, Schlossman DM, Borowitz MJ, Laurent G, Jansen FK, Schmidt C, Frankel AE: A phase I study of Tric-A chain immunotoxin in refractory chronic lymphocytic leukemia. *J Biol Response Mod* 7:97-113, 1988
256. Yokota S, Hara H, Luo Y, Seon BK: Synergistic potentiation of in vivo antitumor activity of anti-human T-leukemia immunotoxins by recombinant alpha-interferon and daunorubicin. *Cancer Res* 50:32-7, 1990
257. Flavell DJ, Warnes S, Noss A, Flavell SU: Host-mediated antibody-dependent cellular cytotoxicity contributes to the in vivo therapeutic efficacy of an anti-CD7-saporin immunotoxin in a severe combined immunodeficient mouse model of human T-cell acute lymphoblastic leukemia. *Cancer Res* 58:5787-94, 1998

258. Ghetie MA, Tucker K, Richardson J, Uhr JW, Vitetta ES: The antitumor activity of an anti-CD22 immunotoxin in SCID mice with disseminated Daudi lymphoma is enhanced by either an anti-CD19 antibody or an anti-CD19 immunotoxin. *Blood* 80:2315-20, 1992
259. Ghetie MA, Picker LJ, Richardson JA, Tucker K, Uhr JW, Vitetta ES: Anti-CD19 inhibits the growth of human B-cell tumor lines in vitro and of Daudi cells in SCID mice by inducing cell cycle arrest. *Blood* 83:1329-36, 1994
260. Ghetie MA, Ghetie V, Vitetta ES: Anti-CD19 antibodies inhibit the function of the P-gp pump in multidrug-resistant B lymphoma cells. *Clin Cancer Res* 5:3920-7, 1999
261. Hamawy MM, Tsuchina M, Manthei ER, Dong Y, Fechner JE, Knechtle JS: Activation of T lymphocytes for adhesion and cytokine expression by toxin-conjugated anti-CD3 monoclonal antibodies. *Transplantation* 68:693-8, 1999
262. Meng G, Jiang Y, Hubbard W, Thomas JM: The effect of anti-CD3-immunotoxin on T lymphocyte function in vitro. *Transpl Immunol* 6:53-9, 1998
263. van Oosterhout YV, van Emst L, Schattenberg AV, Tax WJ, Ruiter DJ, Spits H, Nagengast FM, Masereeuw R, Evers S, de Witte T, Preijers FW: A combination of anti-CD3 and anti-CD7 ricin A-immunotoxins for the in vivo treatment of acute graft-versus-host disease. *Blood* 95:3693-3701, 2000
264. Grossbard ML, Press OW, Appelbaum FR, Bernstein ID, Nadler LM: Monoclonal antibody-based therapies of leukemia and lymphoma. *Blood* 80:863-78, 1992
265. Falini B, Bolognesi A, Flenghi L, Tazzari PL, Broe MK, Stein H, Durkop H, Aversa F, Corneli P, Pizzolo G, et al.: Response of refractory Hodgkin's disease to monoclonal anti-CD30 immunotoxin. *Lancet* 339:1195-6, 1992
266. Grossbard ML, Freedman AS, Ritz J, Coral F, Goldmacher VS, Eliseo L, Spector N, Dear K, Lambert JM, Blattler WA, et al.: Serotherapy of B-cell neoplasms with anti-B4-blocked ricin: a phase I trial of daily bolus infusion. *Blood* 79:376-85, 1992
267. Roy DC, Ouellet S, Le Houillier C, Ariniello PD, Perreault C, Lambert JM: Elimination of neuroblastoma and small-cell lung cancer cells with an anti-neural cell adhesion molecule immunotoxin. *J Natl Cancer Inst* 88:1136-45, 1996
268. Byers VS, Henslee PJ, Kernan NA, Blazar BR, Gingrich R, Phillips GL, LeMaistre CF, Gilliland G, Antin JH, Martin P, et al.: Use of an anti-pan T-lymphocyte ricin A chain immunotoxin in steroid-resistant acute graft-versus-host disease. *Blood* 75:1426-32, 1990
269. Vitetta ES, Stone M, Amlot P, Fay J, May R, Till M, Newman J, Clark P, Collins R, Cunningham D, Ghetie V, Uhr JW, Thorpe PE: Phase I immunotoxin trial in patients with B-cell lymphoma. *Cancer Res* 51:4052-8, 1991
270. Amlot PL, Stone MJ, Cunningham D, Fay J, Newman J, Collins R, May R, McCarthy M, Richardson J, Ghetie V, Ramilo O, Thorpe PE, Uhr JW, Vitetta ES: A phase I study of an anti-CD22-deglycosylated ricin A chain immunotoxin in the treatment of B-cell lymphomas resistant to conventional therapy. *Blood* 82:2624-33, 1993
271. Sausville EA, Headlee D, Stetler-Stevenson M, Jaffe ES, Solomon D, Figg WD, Herdt J, Kopp WC, Rager H, Steinberg SM, et al.: Continuous infusion of the anti-CD22 immunotoxin IgG-RFB4-SMPT-dgA in patients with B-cell lymphoma: a phase I study. *Blood* 85:3457-65, 1995
272. Stone MJ, Sausville EA, Fay JW, Headlee D, Collins RH, Figg WD, Stetler-Stevenson M, Jain V, Jaffe ES, Solomon D, Lush RM, Senderowicz A, Ghetie V, Schindler J, Uhr JW, Vitetta ES: A phase I study of bolus versus continuous infusion of the anti-CD19 immunotoxin, IgG-HD37-dgA, in patients with B-cell lymphoma. *Blood* 88:1188-97, 1996
273. Multani PS, O'Day S, Nadler LM, Grossbard ML: Phase II clinical trial of bolus infusion anti-B4 blocked ricin immunoconjugate in patients with relapsed B-cell non-Hodgkin's lymphoma. *Clin Cancer Res* 4:2599-604, 1998

-
274. LeMaistre CF, Rosen S, Frankel A, Kornfeld S, Saria E, Meneghetti C, Drajek J, Fishwild D, Scanlon P, Byers V: Phase I trial of H65-RTA immunoconjugate in patients with cutaneous T- cell lymphoma. *Blood* 78:1173-82, 1991
275. LeMaistre CF, Meneghetti C, Rosenblum M, Reuben J, Parker K, Shaw J, Deisseroth A, Woodworth T, Parkinson DR: Phase I trial of an interleukin-2 (IL-2) fusion toxin (DAB486IL-2) in hematologic malignancies expressing the IL-2 receptor. *Blood* 79:2547-54, 1992
276. Vitetta ES, Thorpe PE, Uhr JW: Immunotoxins: magic bullets or misguided missiles? *Immunol Today* 14:252-9, 1993
277. Wawrzynczak EJ: Systemetic immunotoxin therapy of cancer: advances and prospects. *Br J Cancer* 64:624-30, 1991

IMMUNOTOXINS IN HEMATOLOGY



CONTENTS

- 2.1 Hematopoietic stem cell transplantation
- 2.2 Ex vivo purging of stem cell grafts
- 2.3 Treatment of GVHD and graft rejection
- 2.4 Treatment of leukemias and lymphomas
- 2.5 Starting point of thesis

2.1 HEMATOPOIETIC STEM CELL TRANSPLANTATION:

The transplantation of hematopoietic stem cells (SCT) forms a world-wide accepted method for the reconstitution of normal hematopoiesis in patients treated for a hematological malignancy or otherwise suffering from a defective hematopoietic or immunologic system. The stem cells can be derived from the patient himself (autologous transplantation) or, from a suitable donor (allogeneic transplantation). Major obstacles for successful outcome of SCT form recurrence of the malignancy and, in case of allogeneic SCT, graft-rejection and graft-versus-host disease (GVHD). ITs have been used in an attempt to improve on the outcome of transplantation by specifically removing undesirable cell types from either the stem cells graft or the patient. These can be malignant cells as well as over-reactive or misdirected immunocompetent cells (schematically depicted in Figure 1). Paragraph 2.2 and 2.3 review the current status of the SCT-related clinical applications of ITs described so far.




Figure 1 Schematic representation of the stem cell transplantation (STC) procedure

A patient suffering from a hematological malignancy is treated with extensive radio and/or chemotherapy for the elimination of malignant cells. In order to restore the affected hematopoiesis the patient is then transplanted with hematopoietic stem cells previously collected during clinical remission (autologous STC), or derived from a suitable donor (allogeneic STC). Adverse events negatively affecting the outcome of SCT are recurrence of the malignancy, and, in case of allogeneic STC, rejection of the stem cell graft and graft-versus-host disease (GVHD). With their specific toxicity, ITs can be used for the ex vivo or in vivo elimination of over-reactive or misdirected immunocompetent cells (either of donor or recipient origin: indicated as ID and IR, respectively), or residual malignant cells (indicated as M).

2.2 EX VIVO PURGING OF STEM CELL GRAFTS

Allogeneic stem cell grafts

Immunocompetent T cells in the graft play a key role in allogeneic SCT. Their (minimal) presence is required for a successful engraftment. Besides, they may contribute to the so-called graft-versus-leukemia effect (GVL), which involves the elimination of residual malignant cells. Unfortunately, they may also initiate graft-versus-host disease (GVHD), a major cause of morbidity and mortality after allogeneic STC. The partial depletion of T cells from allogeneic marrow grafts has been the first reported clinical use of ITs. The initial studies used ricin A-chain or ricin holotoxin conjugated to MoAbs against CD5,

CD3 or a mixture of CD3, CD5 and CD11a.¹⁻⁴ The *ex vivo* application allowed the addition of cytotoxicity enhancer NH₄Cl (ricin A-chain) or an excess of lactose to prevent aspecific binding (ricin holotoxin). The T-cell depletion ranged from 1-3 logs for the individual ITs, to up to 4 logs for the combination. All studies noted a significant decrease in the incidence of severe GVHD. Unfortunately, the removal of T cells appeared to be a double-edged sword, associated with poor engraftment and an increased incidence of relapse of the underlying disease. The same holds true for other immunological or physical methods used for (partial) T cell depletion or positive selection of CD34 positive cells.⁵⁻⁷ Even though some of these methods enable the exact control of the number of T cells in the final graft, SCT remains a complicated procedure, often balancing between graft-failure and leukemic relapse on one side and life-threatening GVHD at the other. Ideally, future purging techniques should deplete only those T cells destined to participate in the anti-host reaction, while conserving those with anti-leukemia reactivity. A step in that direction is the ongoing identification of dominant GVHD-associated CTL epitopes.⁸ In combination with the tetrameric HLA-peptide complexes technology ('tetramers'), this might bring back a role for ITs. Conjugating toxins to TCR-specific tetramers, may create highly selective ITs capable of killing CTL with undesired specificities.^{9,10} Alternatively, T-cell antigens might directly be fused to a toxin, as has been demonstrated for myelin basic protein.¹¹ The final success of these strategies is dependent on the degree to which GVHD and GVL can be separated.

Autologous stem cell grafts

Autologous SCT forms an alternative when no suitable donor is available. In this case, the stem cell graft is collected during remission and prior to the destruction of bone marrow by chemo- and/or radiotherapy. This scenario introduces the risk of persistent malignant cells within the graft contributing to a relapse upon re-infusion. ITs have been used for the purging of malignant cells from autologous grafts of patients with T-lineage acute lymphoblastic leukemia or T cell lymphoma. Though these studies demonstrated no adverse effect on engraftment, a clear clinical benefit regarding relapse free survival was hard to prove.^{4,12,13} A conclusive statistical evaluation will require large randomized clinical trials. Nevertheless, genetic marking studies clearly have demonstrated the contribution of 'graft-originating' malignant cells to the relapse following autologous SCT for neuroblastoma and CML.^{14,15} These results still argue for maximal tumor reduction of autologous grafts, e.g. by the IT-based purging of positively selected CD34-positive hematopoietic progenitor cells.^{16,17}

2.3 TREATMENT OF GVHD AND STEM CELL REJECTION

Acute GVHD (aGVHD) and accompanying immunosuppression continues to be the major cause of morbidity and mortality in patients receiving allogeneic SCT.¹⁸ With the

standard prophylaxis (typically cyclosporine, often combined with methotrexate) GVHD occurs in 30-70% of HLA-matched recipients and contributes to death in 20-40% of those affected. Even if a patient survives severe GVHD, this generally results in long-lasting disability and morbidity leading to repeated admission to hospital.

The etiology of aGVHD is considered a multi-step process, already initiated by the preparative treatment of the host with chemo- and/or radiotherapy (Figure 2).^{19,20} The




Figure 2 Schematic representation of the immunopathophysiology of GVHD

GVHD is considered a multi-step process. The pretransplant conditioning regimen induces damage of host tissues resulting in the secretion of inflammatory cytokines (TNF- α and IL-1) and the subsequent upregulation of MHC antigens. These conditions favor the interaction with allo-reactive donor T cells, which upon activation secrete IL-2, INF- γ . These cytokines induce further T-cell expansion and activate monocytes, macrophages and NK-cells to produce IL-1, TNF- α and IL-6. Altogether, this cascade may cause serious direct and indirect cytotoxicity to epithelial cells of skin, liver and gut.

Figure reprinted from Ref. 92 by courtesy of Marcel Dekker, Inc.

inevitable damage to the gastrointestinal tract, liver, and blood vessels induces an inflammatory reaction associated with the release of IL-1 and TNF- α and the over-expression of cytoadhesion molecules and MHC class II antigens. These events lead to an enhanced interaction of allo-reactive donor-T cells with major or minor HLA antigens on the affected tissues. Activation of these cells results in the production of

cytokines (IL-2 and INF- γ) which in turn activate additional effector cells, like monocytes, macrophages and NK-cells, to produce inflammatory proteins (IL-1, TNF- α and IL-6). Altogether, this may result in serious direct and indirect cytotoxicity to epithelial cells of skin, liver and gut.

When prophylaxis has failed, severe aGVHD is usually treated with low dose corticosteroids. If the reaction progresses, the dose may be increased or, alternatively, polyclonal antithymocyte/antilymphocyte globulin (ATG/ALG) or experimental immunosuppressive agents may be applied. Most of the experimental therapies are focussed on the removal of T cells or interference with their IL-2 signaling. Based on their capacity of eliminating specific cell populations, IT may also be suited for this type of therapy.

H65-RTA, a ricin A-based anti-CD5 IT, is the first IT clinically used for the *in vivo* treatment of aGVHD. In the initial studies, the use of H65-RTA resulted in extensive responses in about half of the patients with steroid-resistant aGVHD (~25% complete responses).²¹⁻²³ Compared to historical ATG-treated controls, the overall responses rates obtained with H65-RTA were higher (52 vs. 31%) and the median survival was longer (148 vs. 80 days).²³ Unfortunately, in later comparative trials H65-RTA was not more effective than high-dose corticosteroids or ATG.^{24,25} Additionally, the use of H65-RTA for the prophylaxis of aGVHD or stem cell rejection revealed no clear advantage over current cyclosporin-based regimens.²⁶⁻³⁰

Several animal studies now suggest that CD3-directed ITs may be more suitable for the control of GVHD or graft-rejection than their anti-CD5 counterparts. The group of Vallera and Blazer studied the therapeutic potential of a hamster derived anti-murine-CD3 MoAb in an aggressive murine model of established GVHD. Conjugated to ricin A, the CD3 MoAb appeared to be far more effective than a similar ITs directed against the murine CD5-equivalent (Lyt-1), despite both ITs having similar clearance and biodistribution.³¹ Its efficacy could partly be explained by the CD3 MoAb inducing modulation of the T-cell receptor and T cell depletion.³² Moreover, conjugation to ricin A dramatically improved the depletion efficacy of both the IgG and F(ab')₂-fragments.^{31,33} Vallera *et al.* are now trying to identify the most appropriate form of anti-CD3 ITs for clinical use. Their latest report addressed the evaluation of a single chain fusion toxin, comprising the Fv fragment of the anti-murine-CD3 MoAb genetically linked to truncated diphtheria toxin.³⁴⁻³⁶ Due to its relatively small size (68 kDA), this fusion toxins was readily filtrated into kidney, causing severe and irreversible renal damage. This problem appears now to be solved by the introduction of a cysteine residue at the c-terminus of the sFv part, resulting in the formation of disulfide bonded bivalent IT with an expanded therapeutic window.³⁶

Neville *et al.* studied the T-cell depletion efficacy of an anti-rhesus-CD3 murine IgG1 MoAb conjugated to a diphtheria toxin binding mutant (FN18-CRM9).³⁷ This construct was able of depleting >99% of T cells from both the blood and lymph node compart-

ments of healthy rhesus monkeys. Restoration of the T-cell compartment took 12 to 30 days and appeared to be thymus dependent. Depletion activity of native FN18 was limited to the blood compartment and lasted only for a day.³⁷ Additional primate studies focussed on the use of FN18-CRM9 in protocols aiming at tolerance induction. Encouraging results were obtained, including the induction of long-term tolerance in kidney and pancreatic islet transplant recipients.³⁸⁻⁴⁴ As an extension to these primate studies, an anti-porcine CD3-IT has been generated (pCD3-CRM9) for evaluation in a non-myeloblastic stem cell transplantation model established in miniature swine.⁴⁵ When pCD3-CRM9 was added to a mild preparative regimen, infusion of blood stem cells from leucocyte antigen-matched animals resulted in stable mixed chimerism without aGVHD occurring.⁴⁶ Moreover, this protocol provided donor-specific tolerance as indicated by long-term acceptance of donor skin and consistent rejection of third party skin.

2.4 TREATMENT OF LEUKEMIAS AND LYMPHOMAS

Their diffuse nature, and the relative abundance of tissue-specific antigens, make hematological malignancies an attractive target for IT therapy. Table 1 summarizes the outcome of the major studies reported so far. Most of them are directed at B cell derived malignancies, mainly due to their high incidence as compared to T-cell neoplasms (ratio of about 5:1).

Vitetta and colleagues performed a series of studies in which they evaluated RFB4-dgA (anti-CD22) and HD37-dgA (anti-CD19) for the treatment of relapsed Non-Hodgkin's Lymphoma (NHL). Initial studies with RFB4-dgA demonstrated no particular advantages of Fab'-ITs over whole IgG-ITs with regard to clinical responses, toxicities or immunogenicity. Subsequent studies were performed with the more cost-effective whole-IgG ITs. Both RFB4-dgA and HD37-dgA were administered as bolus and continuous infusions, revealing no obvious advantage of one schedule above the other, and resulting in a comparable maximal tolerable dose (MTD: ~16-20 mg/m²/8 days for both ITs). The most prominent toxicities were VLS, myalgia, rhabdomyolysis and aphasia. Overall response rates of RFB4-dgA were higher than those obtained with HD37-dgA (~25% and 10%, respectively). Based on the synergistic effect observed in preclinical studies⁴⁷, the combination of HD37-dgA and RFB4-dgA (Combotox) is now being clinically evaluated.⁴⁸ The first study appears to be hampered by unpredictable clinical courses, attributable to batch-dependent multimerization of HD37-dgA. Especially patients lacking circulating tumor cells were vulnerable for severe VLS and hemolytic uremic syndrome (HUS). Three deaths were noted, of which two were probably IT-related. Further studies are planned using non-aggregate-forming formulations of HD37-dgA.

Another comprehensive series of clinical studies is performed with B4-bR, an anti-CD19

Table 1 *Major clinical trials of IT for the treatment of hematological malignancies*

	Antigen	Disease ^a	Administration	#Patients
T101-RTA	CD5	B-CLL	b.i. twice wkly x 4 wk	9
H65-RTA	CD5	CTCL	b.i. daily x 10 d	14
Anti-CD7-dgA (DA7)	CD7	T-NHL, T-LGL	b.i. daily x 5 d	11
HD37-dgA	CD19	B-NHL	b.i. every 48h x 4 c.i. for 8 d b.i. every 4h x 4	23 10 8
B43-PAP	CD19	B-ALL	c.i. for 5 d, 1 or 3 cycles	17
RFB4-dgA (Fab')	CD22	B-NHL	b.i. every 48h x 2-6	15
RFB4-dgA (IgG)	CD22	B-NHL	b.i. every 48h x 2-12 c.i. for 8 d	26 18
RFB4-dgA (IgG) & HD37-dgA	CD19/CD22	B-NHL	c.i. for 8 d	22
B4-bR	CD22	B-cell neoplasms B-cell neoplasms B-NHL, postABMT, Ph I B-NHL, postABMT, Ph II B-NHL, postABMT, Ph III AIDS-NHL, relapsed AIDS-NHL, untreated	b.i. daily x 5 d c.i. for 7 d c.i. for 7 d c.i. for 7 d, every 14 d c.i. for 7 d, every 7 d c.i. for 28 d c.i. for 7 d	25 34 12 49 157 9 28
RFT5-dgA	CD25	HD	b.i. every 48h x 4, 1-4 cycles	25
Anti-Tac(Fv)-PE38 (LMB-2)	CD25	B & T lymph/leukemia, HD	b.i. every 48h x 3	35
BerH2-SAP	CD30	Hodgkin's	b.i. on day 1 \pm 7	12
DAB486IL-2	IL-2R	NHL, Hodgkin's, CLL NHL, Hodgkin's, CLL NHL CTCL & Sezary	b.i. on day 1, 3-5, 14-20 b.i. daily x 5 d b.i. daily x 5 d b.i. dly x 5 d, every 21 d	18 15 23 14
DAB389IL-2	IL-2R	Mycosis fungoides	b.i. dly x 5 d every 21 d	71

MTD	Toxicities ^b	Human anti-IT Ab	Responses ^c	Ref.
24-112 mg/m ² total	Fever, nausea, rash	1/9	Rapid transient fall in WBC	68,69
0.33 mg/kg/d	VLS, fever, chills, nausea, fatigue	10/14	4 PR	70
0.2 mg/kg/d	VLS, aphasia	1/11	0 CR, 2 PR	71
16 mg/m ² /8 d	VLS, rhabdomyolysis	5/15	1 CR, 1 PR	72
19.2 mg/m ² /8 d	VLS, acrocyanosis	2/8	1 PR	72
8 mg/m ²	VLS, hypotension, rhabdomyolysis	2/7	1 PR	73
1.25 mg/kg	VLS, myalgia	0	4 CR, 1 PR	74
75 mg/m ²	VLS, myalgias, aphasia, rhabdomyolysis	4/14	5 PR	75
26-40 mg/m ² total	VLS, myalgias, aphasia, rhabdomyolysis	9/24	1 CR, 5 PR	76
19.2 mg/m ² /8 d	VLS, aphasia, hypotension	6/14	4 PR	77
≤10 mg - 30 mg/m ²	VLS, HUS	6/22	2 PR	48
250 µg/kg total	AST, ALT, PLTs, VLS, fatigue	9/25	1 CR, 2 PR	78
350 µg/kg total	AST, ALT, PLTs, VLS, myalgia, edema	24/34	2 CR, 3 PR	79
280 µg/kg total	AST, ALT, PLTs, VLS, fever	7/12	7 relapse free, MFU 4 years	80
210 µg/kg total	AST, ALT, PLTs, VLS	23/49	27 relapse free, MFU 38 months	50
210 µg/kg total	PLTs, VLS, fatigue, dyspnea	NA	Comparable with control	81
560 µg/kg total	AST, ALT	3/9	1 CR, 1 PR	82
140 µg/kg total	AST, ALT, fever, myalgia, fatigue	8/28	14 CR, 12 PR	83
15 mg/m ² /cycle	VLS, myalgias, hypersensitivity	7/15	2 PR	84,85
120 µg/kg total	AST, ALT, cardiomyopathy, diarrhea, fever	11/35	1 CR, 7 PR	86
0.4 mg/kg/d	AST, ALT, PLTs, VLS, fever, fatigue	12/12	8 PR	87,88
0.1 mg/kg/d	AST, ALT, fever, dyspnea, rash	7/18	1CR, 2 PR	51
0.2 mg/kg/d	AST, ALT, hypoalbuminemia, hypersensitivity	6/14	1 CR	89
0.3 mg/kg/d	PLTs, renal insufficiency, hemolysis	15/20	2 PR	52
0.2 mg/kg/d	AST, ALT	8/14	1 PR	55
9 or 18 µg/kg/d	AST, ALT, VLS, hypersensitivity	70/71	3 CR, 18 PR	90,91

^aB-CLL, B-cell chronic lymphocytic leukemia; CTCL, cutaneous T-cell lymphoma; T-NHL, T-cell non-Hodgkin's lymphoma; T-LGL, T-cell large granular lymphocyte leukemia; B-NHL, B-cell non-Hodgkin's lymphoma; B-ALL, B-cell acute lymphoblastic leukemia; postABMT, post autologous bone marrow transplantation; AIDS-NHL, AIDS-related non-Hodgkin's lymphoma; HD, Hodgkin's disease; AML, acute myeloid leukemia.

^bVLS, vascular leak syndrome; HUS, hemolytic uremic syndrome; AST, aspartate amino transferase; ALT, amino alanine transferase; PLTs, thrombocytopenia

^cWBC, white blood cell counts; PR, partial response; CR, complete response; MFU, median follow up.

MoAb conjugated to blocked ricin (B4-bR). Following the evaluation of various dosing protocols and different clinical settings, B4-bR was the first IT to be tested in the minimal disease setting. B4-bR has been administered to patients with NHL brought in complete remission with high-dose chemotherapy and stem cell transplantation. Regrettably, no beneficial effects on disease free survival could be demonstrated in a randomized Phase III trial.^{49,50} The hepatotoxicity of bR was an important factor limiting therapeutic success by reducing the therapeutic window.

Of note is the good performance of IL-2 receptor (IL-2R) targeted recombinant ITs in certain hematological diseases. Truncated diphtheria toxin spliced to IL-2 (DAB486IL-2 and the improved version DAB389IL-2) showed impressive efficacy in treating CD25-positive refractory cutaneous T-cell lymphoma (CTCL), with overall response rates of approximately 30%.⁵¹⁻⁵⁸ DAB389IL-2 (Ontak) has been approved by the FDA (May 1999) for the treatment of patients with persistent or recurrent CD25-positive CTCL.

The group of Pastan created an alternative anti-IL-2R IT by genetically linking an anti-CD25 MoAb Fv-fragment to truncated *Pseudomonas* exotoxin (LMB-2).⁵⁹⁻⁶¹ This IT differs from DAB389IL-2 in that it binds CD25 (IL-2R α -chain) with high affinity also in the absence of the IL-2R β and γ chain. This may be an advantage for malignancies solely expressing the IL-2R α -chain, like the Reed Sternberg cells in Hodgkin's disease. LMB-2 has been administered to patients with various CD25-positive hematological malignancies. Remarkably, all four patients with hairy cell leukemia (HCL) responded to therapy, with one complete responder of more than 20 months of duration. Additionally, four partial responses were observed in CTCL ($n = 1$), CLL ($n = 8$), HD ($n = 11$), and adult T-cell leukemia ($n = 2$). Additional *in vitro* studies with freshly derived patient cells confirmed the particular vulnerability of HCL cells as compared to B-CLL cells expressing equal or higher numbers of CD25 per cell.⁶²

The above examples represent a proof of principal that IT-based therapy of hematological malignancies can be clinically useful. Several clinical responses could be noted in refractory patients with bulky disease. The ongoing and coming clinical trials will be crucial in determining the future role of ITs. Not only will they teach us which particular hematological diseases are vulnerable for what type of IT, they also should identify the optimal clinical setting for maximal therapeutic benefit.

2.5 STARTING POINT OF THESIS

The UMC St Radboud was among the first to introduce the IT research in the Netherlands, in 1983. This research was initially focused on identifying the factors influencing the activity of IT, like the nature and density of target-antigens, the internalization rate, the use of toxicity enhancers, and the activation state of the target cells.⁶³⁻⁶⁶ These stud-

ies resulted in the first institutional clinical application, the *ex vivo* purging of autologous marrow grafts of patients suffering from T cell leukemia or lymphoma.¹² Aim of this treatment was to reduce the incidence of a malignant relapse caused by residual malignant cells in the autologous graft. The more or less routinely *ex vivo* purging of bone marrow can be regarded as the starting point of this theses. As such, this application initiated additional research focussing at ways to improve the purging efficacy and to reliably monitor its efficacy in a heterogeneous population such as bone marrow grafts. At the mean time, the international excitement regarding the therapeutic possibilities of the ‘magic-bullit concept’ more or less drove the IT from the laboratory to the clinic. Several institutes, mostly US academic centers, initiated phase I/II studies based on the IT-targeting of diffuse and solid tumors and varying immune diseases. Encouraged by the outcome of these early studies, especially regarding the treatment of diffuse disorders, and motivated by the urgent clinical need, the Department of Hematology decided to develop an IT-based therapy for lifethreatening aGVHD. Additionally, another research line was started studying the anti-tumor activity of an anti-B cell IT in a mouse xenografted tumor model. The next part of this thesis forms a selection of the institutional IT-research during the last decade. It covers part of the early *in vitro* work (Chapter 3 to 5), and the set up and preliminary results of a clinical pilot study for the treatment of aGVHD (Chapters 6 and 7). Notably, the complementary work, addressing the IT-based treatment of B cell malignancies, is described in the thesis of Van Horssen.⁶⁷

REFERENCES

1. Fauser AA, Shustik C, Langleben A, Laurent G, Kanz L, Spurll GM, Price G, Ahlgren PD, Cooper BA: T-cell depletion with ricin A-chain T101 in allogeneic bone marrow transplantation to prevent severe graft-versus-host disease. *Clin Invest Med* 11:40-6, 1988
2. Martin PJ, Hansen JA, Torok-Storb B, Moretti L, Press O, Storb R, Thomas ED, Weiden PL, Vitetta ES: Effects of treating marrow with a CD3-specific immunotoxin for prevention of acute graft-versus-host disease [published erratum appears in *Bone Marrow Transplant* 1989 Mar;4(2):215]. *Bone Marrow Transplant* 3:437-44, 1988
3. Vallera DA: Immunotoxins for ex vivo bone marrow purging in human bone marrow transplantation. *Cancer Treat Res* 37:515-35, 1988
4. Filipovich AH, Vallera DA, Youle RJ, Haake R, Blazar BR, Arthur D, Neville DM, Jr., Ramsay NK, McClave P, Kersey JH: Graft-versus-host disease prevention in allogeneic bone marrow transplantation from histocompatible siblings. A pilot study using immunotoxins for T cell depletion of donor bone marrow. *Transplantation* 44:62-9, 1987
5. de Witte T, Hoogenhout J, de Pauw B, Holdrinet R, Janssen J, Wessels J, van Daal W, Hustinx T, Haanen C: Depletion of donor lymphocytes by counterflow centrifugation successfully prevents acute graft-versus-host disease in matched allogeneic marrow transplantation. *Blood* 67:1302-8, 1986
6. Hale G, Bright S, Chumbley G, Hoang T, Metcalf D, Munro AJ, Waldmann H: Removal of T cells from bone marrow for transplantation: a monoclonal antilymphocyte antibody that fixes human complement. *Blood* 62:873-82, 1983
7. de Wynter EA, Coutinho LH, Pei X, Marsh JC, Hows J, Luft T, Testa NG: Comparison of purity and enrichment of CD34+ cells from bone marrow, umbilical cord and peripheral blood (primed for apheresis) using five separation systems. *Stem Cells* 13:524-32, 1995
8. Mutis T, Gillespie G, Schrama E, Falkenburg JH, Moss P, Goulmy E: Tetrameric HLA class I-minor histocompatibility antigen peptide complexes demonstrate minor histocompatibility antigen-specific cytotoxic T lymphocytes in patients with graft-versus-host disease. *Nat Med* 5:839-42, 1999
9. Whelan JA, Dunbar PR, Price DA, Purbhoo MA, Lechner F, Ogg GS, Griffiths G, Phillips RE, Cerundolo V, Sewell AK: Specificity of CTL Interactions with Peptide-MHC Class I Tetrameric Complexes Is Temperature Dependent. *J Immunol* 163:4342-4348, 1999
10. Clark BR, Deshpande SV, Sharma SD, Nag B: Antigen-specific deletion of cloned T cells using peptide-toxin conjugate complexed with purified class II major histocompatibility complex antigen. *J Biol Chem* 269:94-9, 1994
11. Brenner T, Steinberger I, Soffer D, Beraud E, Ben-Nun A, Lorberboum-Galski H: A novel antigen-toxin chimeric protein: myelin basic protein- pseudomonas exotoxin (MBP-PE 40) for treatment of experimental autoimmune encephalomyelitis. *Immunol Lett* 68:403-10, 1999
12. Preijers FW, De Witte T, Wessels JM, De Gast GC, Van Leeuwen E, Capel PJ, Haanen C: Autologous transplantation of bone marrow purged in vitro with anti-CD7- (WT1-) ricin A immunotoxin in T-cell lymphoblastic leukemia and lymphoma. *Blood* 74:1152-8, 1989
13. Uckun FM, Kersey JH, Vallera DA, Ledbetter JA, Weisdorf D, Myers DE, Haake R, Ramsay NK: Autologous bone marrow transplantation in high-risk remission T-lineage acute lymphoblastic leukemia using immunotoxins plus 4- hydroperoxycyclophosphamide for marrow purging. *Blood* 76:1723-33, 1990
14. Brenner MK, Rill DR, Moen RC, Krance RA, Mirro J, Jr., Anderson WF, Ihle JN: Gene-marking to trace origin of relapse after autologous bone-marrow transplantation. *Lancet* 341:85-6, 1993

15. Deisseroth AB, Zu Z, Claxton D, Hanania EG, Fu S, Ellerson D, Goldberg L, Thomas M, Janicek K, Anderson WF, et al.: Genetic marking shows that Ph+ cells present in autologous transplants of chronic myelogenous leukemia (CML) contribute to relapse after autologous bone marrow in CML. *Blood* 83:3068-76, 1994
16. Spyridonidis A, Schmidt M, Bernhardt W, Papadimitriou A, Azemar M, Wels W, Groner B, Henschler R: Purging of mammary carcinoma cells during ex vivo culture of CD34+ hematopoietic progenitor cells with recombinant immunotoxins. *Blood* 91:1820-7, 1998
17. Straka C, Drexler E, Mitterer M, Langenmayer I, Pfefferkorn L, Stade B, Koll R, Emmerich B: Auto-transplantation of B-cell purged peripheral blood progenitor cells in B-cell lymphoma [letter]. *Lancet* 345:797-8, 1995
18. Ferrara JL, Deeg HJ: Graft-versus-host disease. *N Engl J Med* 324:667-74, 1991
19. Ferrara JL: Cytokines other than growth factors in bone marrow transplantation. *Curr Opin Oncol* 6:127-34, 1994
20. Antin JH, Ferrara JL: Cytokine dysregulation and acute graft-versus-host disease. *Blood* 80:2964-8, 1992
21. Kernan NA, Byers V, Scannon PJ, Mischak RP, Brochstein J, Flomenberg N, Dupont B, O'Reilly RJ: Treatment of steroid-resistant acute graft-vs-host disease by in vivo administration of an anti-T-cell ricin A chain immunotoxin. *Jama* 259:3154-7, 1988
22. Byers VS, Henslee PJ, Kernan NA, Blazar BR, Gingrich R, Phillips GL, LeMaistre CF, Gilliland G, Antin JH, Martin P, et al.: Use of an anti-pan T-lymphocyte ricin A chain immunotoxin in steroid-resistant acute graft-versus-host disease. *Blood* 75:1426-32, 1990
23. Lomen PL, Steward KK, Saks SR: Safety and efficacy of H65-RTA in the treatment of steroid-resistant acute graft versus host disease after allogeneic bone marrow transplantation. *Blood* 78(suppl 1):230, 1991
24. Martin PJ, Nelson BJ, Appelbaum FR, Anasetti C, Deeg HJ, Hansen JA, McDonald GB, Nash RA, Sullivan KM, Witherspoon RP, Scannon PJ, Friedmann N, Storb R: Evaluation of a CD5-specific immunotoxin for treatment of acute graft- versus-host disease after allogeneic marrow transplantation. *Blood* 88:824-30, 1996
25. Hings IM, Severson R, Filipovich AH, Blazar BR, Kersey JH, Ramsay NK, McGlave PB, Weisdorf DJ: Treatment of moderate and severe acute GVHD after allogeneic bone marrow transplantation. *Transplantation* 58:437-42, 1994
26. Koehler M, Hurwitz CA, Krance RA, Coustan-Smith E, Williams LL, Santana V, Ribeiro RC, Brenner MK, Heslop HE: XomaZyme-CD5 immunotoxin in conjunction with partial T cell depletion for prevention of graft rejection and graft-versus-host disease after bone marrow transplantation from matched unrelated donors. *Bone Marrow Transplant* 13:571-5, 1994
27. Phillips GL, Nevill TJ, Spinelli JJ, Nantel SH, Klingemann HG, Barnett MJ, Shepherd JD, Chan KW, Meharchand JM, Sutherland HJ, et al.: Prophylaxis for acute graft-versus-host disease following unrelated donor bone marrow transplantation. *Bone Marrow Transplant* 15:213-9, 1995
28. Przepiorka D, Chan KW, Champlin RE, Culbert SJ, Petropoulos D, Ippoliti C, Khouri I, Huh YO, Vreissendorp H, Deisseroth AB, et al.: Prevention of graft-versus-host disease with anti-CD5 ricin A chain immunotoxin after CD3-depleted HLA-nonidentical marrow transplantation in pediatric leukemia patients. *Bone Marrow Transplant* 16:737-41, 1995
29. Henslee-Downey PJ, Parrish RS, MacDonald JS, Romond EH, Marciniak E, Coffey C, Ciocchi G, Thompson JS: Combined in vitro and in vivo T lymphocyte depletion for the control of graft-versus-host disease following haploidentical marrow transplant. *Transplantation* 61:738-45, 1996
30. Pagliaro LC, Liu B, Munker R, Andreeff M, Freireich EJ, Scheinberg DA, Rosenblum MG: Humanized M195 monoclonal antibody conjugated to recombinant gelonin: an anti-CD33 immunotoxin with antileukemic activity. *Clin Cancer Res* 4:1971-6, 1998

31. Vallera DA, Taylor PA, Panoskaltsis-Mortari A, Blazar BR: Therapy for ongoing graft-versus-host disease induced across the major or minor histocompatibility barrier in mice with anti-CD3F(ab')₂-ricin toxin A chain immunotoxin. *Blood* 86:4367-75, 1995
32. Blazar BR, Taylor PA, Vallera DA: In vivo or in vitro anti-CD3 epsilon chain monoclonal antibody therapy for the prevention of lethal murine graft-versus-host disease across the major histocompatibility barrier in mice. *J Immunol* 152:3665-74, 1994
33. Blazar BR, Hirsch R, Gress RE, Carroll SF, Vallera DA: In vivo administration of anti-CD3 monoclonal antibodies or immunotoxins in murine recipients of allogeneic T cell-depleted marrow for the promotion of engraftment. *J Immunol* 147:1492-503, 1991
34. Vallera DA, Panoskaltsis-Mortari A, Jost C, Ramakrishnan S, Eide CR, Kreitman RJ, Nicholls PJ, Pennell C, Blazar BR: Anti-graft-versus-host disease effect of DT390-anti-CD3sFv, a single-chain Fv fusion immunotoxin specifically targeting the CD3 epsilon moiety of the T-cell receptor. *Blood* 88:2342-53, 1996
35. Vallera DA, Panoskaltsis-Mortari A, Blazar BR: Renal dysfunction accounts for the dose limiting toxicity of DT390anti-CD3sFv, a potential new recombinant anti-GVHD immunotoxin. *Protein Eng* 10:1071-6, 1997
36. Vallera DA, Kuroki DW, Panoskaltsis-Mortari A, Buchsbaum DJ, Rogers BE, Blazar BR: Molecular modification of a recombinant anti-CD3epsilon-directed immunotoxin by inducing terminal cysteine bridging enhances anti-GVHD efficacy and reduces organ toxicity in a lethal murine model. *Blood* 96:1157-65, 2000
37. Neville DM, Jr., Scharff J, Hu HZ, Rigaut K, Shiloach J, Slingerland W, Jonker M: A new reagent for the induction of T-cell depletion, anti-CD3-CRM9. *J Immunother Emphasis Tumor Immunol* 19:85-92, 1996
38. Knechtle SJ, Vargo D, Fechner J, Zhai Y, Wang J, Hanaway MJ, Scharff J, Hu H, Knapp L, Watkins D, Neville DM, Jr.: FN18-CRM9 immunotoxin promotes tolerance in primate renal allografts. *Transplantation* 63:1-6, 1997
39. Knechtle SJ, Fechner JH, Jr., Dong Y, Stavrou S, Neville DM, Jr., Oberley T, Buckley P, Armstrong N, Rusterholz K, Hong X, Tsuchida M, Hamawy MM: Primate renal transplants using immunotoxin. *Surgery* 124:438-46; discussion 446-7, 1998
40. Knechtle SJ, Kirk AD, Fechner JH, Jr., Hong X, Dong Y, Hamawy MM, Harlan DM: Inducing unresponsiveness by the use of anti-CD3 immunotoxin, CTLA4-Ig, and anti-CD40 ligand. *Transplant Proc* 31:27S-28S, 1999
41. Thomas JM, Neville DM, Contreras JL, Eckhoff DE, Meng G, Lobashevsky AL, Wang PX, Huang ZQ, Verbanac KM, Haisch CE, Thomas FT: Preclinical studies of allograft tolerance in rhesus monkeys: a novel anti-CD3-immunotoxin given peritransplant with donor bone marrow induces operational tolerance to kidney allografts. *Transplantation* 64:124-35, 1997
42. Thomas JM, Eckhoff DE, Contreras JL, Lobashevsky AL, Hubbard WJ, Moore JK, Cook WJ, Thomas FT, Neville DM, Jr.: Durable donor-specific T and B cell tolerance in rhesus macaques induced with peritransplantation anti-CD3 immunotoxin and deoxyspergualin: absence of chronic allograft nephropathy. *Transplantation* 69:2497-503, 2000
43. Thomas JM, Contreras JL, Jiang XL, Eckhoff DE, Wang PX, Hubbard WJ, Lobashevsky AL, Wang W, Asiedu C, Stavrou S, Cook WJ, Robbin ML, Thomas FT, Neville DM, Jr.: Peritransplant tolerance induction in macaques: early events reflecting the unique synergy between immunotoxin and deoxyspergualin. *Transplantation* 68:1660-73, 1999
44. Thomas F, Contreras J, Ricordi C, Ranuneolia A, Cartner S, Eckhoff D, Huang Z, Neville D, Thomas J: Reversal of naturally occurring insulinopenic diabetes and induction of isolated islet xenograft tolerance in a preclinical study. *Transplant Proc* 31:637-8, 1999
45. Huang CA, Yamada K, Murphy MC, Shimizu A, Colvin RB, Neville DM, Jr., Sachs DH: In vivo T cell depletion in miniature swine using the swine CD3 immunotoxin, pCD3-CRM9. *Transplantation* 68:855-60, 1999

46. Huang CA, Fuchimoto Y, Scheier-Dolberg R, Murphy MC, Neville DM, Jr., Sachs DH: Stable mixed chimerism and tolerance using a nonmyeloablative preparative regimen in a large-animal model. *J Clin Invest* 105:173-81, 2000
47. Ghetie MA, Tucker K, Richardson J, Uhr JW, Vitetta ES: The antitumor activity of an anti-CD22 immunotoxin in SCID mice with disseminated Daudi lymphoma is enhanced by either an anti-CD19 antibody or an anti-CD19 immunotoxin. *Blood* 80:2315-20, 1992
48. Messmann RA, Vitetta ES, Headlee D, Senderowicz AM, Figg WD, Schindler J, Michiel DF, Creekmore S, Steinberg SM, Kohler D, Jaffe ES, Stetler-Stevenson M, Chen H, Ghetie V, Sausville EA: A phase I study of combination therapy with immunotoxins IgG-HD37-dglycosylated ricin A chain (dgA) and IgG-RFB4-dgA (Combotox) in patients with refractory CD19(+), CD22(+) B cell lymphoma. *Clin Cancer Res* 6:1302-13, 2000
49. Grossbard ML, Multani PS, Freedman AS, O'Day S, Gribben JG, Rhuda C, Neuberg D, Nadler LM: A Phase II study of adjuvant therapy with anti-B4-blocked ricin after autologous bone marrow transplantation for patients with relapsed B-cell non-Hodgkin's lymphoma. *Clin Cancer Res* 5:2392-8, 1999
50. Multani PS, Grossbard ML: Immunotoxin therapy of lymphoma: studies with anti-B4-blocked ricin, in Grossbard ML (ed): *Monoclonal antibody-based therapy of cancer*, vol. 15, Basic and clinical oncology series. New York, Marcel Dekker, 1998, p 91-112
51. LeMaistre CF, Meneghetti C, Rosenblum M, Reuben J, Parker K, Shaw J, Deisseroth A, Woodworth T, Parkinson DR: Phase I trial of an interleukin-2 (IL-2) fusion toxin (DAB486IL-2) in hematologic malignancies expressing the IL-2 receptor. *Blood* 79:2547-54, 1992
52. LeMaistre CF, Craig FE, Meneghetti C, McMullin B, Parker K, Reuben J, Boldt DH, Rosenblum M, Woodworth T: Phase I trial of a 90-minute infusion of the fusion toxin DAB486IL-2 in hematological cancers. *Cancer Res* 53:3930-4, 1993
53. Kuzel TM, Rosen ST, Gordon LI, Winter J, Samuelson E, Kaul K, Roenigk HH, Nylen P, Woodworth T: Phase I trial of the diphtheria toxin/interleukin-2 fusion protein DAB486IL-2: efficacy in mycosis fungoides and other non-Hodgkin's lymphomas. *Leuk Lymphoma* 11:369-77, 1993
54. Hesketh P, Caguioa P, Koh H, Dewey H, Facada A, McCaffrey R, Parker K, Nylen P, Woodworth T: Clinical activity of a cytotoxic fusion protein in the treatment of cutaneous T-cell lymphoma. *J Clin Oncol* 11:1682-90, 1993
55. Foss FM, Borkowski TA, Gilliom M, Stetler-Stevenson M, Jaffe ES, Figg WD, Tompkins A, Bastian A, Nylen P, Woodworth T, et al.: Chimeric fusion protein toxin DAB486IL-2 in advanced mycosis fungoides and the Sezary syndrome: correlation of activity and interleukin-2 receptor expression in a phase II study. *Blood* 84:1765-74, 1994
56. Kiyokawa T, Williams DP, Snider CE, Strom TB, Murphy JR: Protein engineering of diphtheria-toxin-related interleukin-2 fusion toxins to increase cytotoxic potency for high-affinity IL-2-receptor-bearing target cells. *Protein Eng* 4:463-8, 1991
57. LeMaistre CF, Saleh MN, Kuzel TM, Foss F, Platanius LC, Schwartz G, Ratain M, Rook A, Freytes CO, Craig F, Reuben J, Nichols JC: Phase I trial of a ligand fusion-protein (DAB389IL-2) in lymphomas expressing the receptor for interleukin-2. *Blood* 91:399-405, 1998
58. Duvic M, Kuzel T, Olsen E, Martin A, Foss F, Kim Y, Heald P, Bacha P, Nichols J: Quality of life is significantly improved in ctcl patients who respond to DAB(389)IL-2 (ONTAK) fusion protein (abstr). *Blood* 92(suppl):2572, 1998
59. Chaudhary VK, Queen C, Junghans RP, Waldmann TA, FitzGerald DJ, Pastan I: A recombinant immunotoxin consisting of two antibody variable domains fused to *Pseudomonas* exotoxin. *Nature* 339:394-7, 1989
60. Kreitman RJ, Bailon P, Chaudhary VK, FitzGerald DJ, Pastan I: Recombinant immunotoxins containing anti-Tac(Fv) and derivatives of *Pseudomonas* exotoxin produce complete regression in mice of an interleukin-2 receptor-expressing human carcinoma. *Blood* 83:426-34, 1994

61. Kreitman RJ, Batra JK, Seetharam S, Chaudhary VK, FitzGerald DJ, Pastan I: Single-chain immunotoxin fusions between anti-Tac and *Pseudomonas* exotoxin: relative importance of the two toxin disulfide bonds. *Bioconjug Chem* 4:112-20, 1993
62. Robbins DH, Margulies I, Stetler-Stevenson M, Kreitman RJ: Hairy cell leukemia, a B-cell neoplasm that is particularly sensitive to the cytotoxic effect of anti-Tac(Fv)-PE38 (LMB-2). *Clin Cancer Res* 6:693-700, 2000
63. Preijers FW, Tax WJ, De Witte T, Janssen A, v.d. Heijden H, Vidal H, Wessels JM, Capel PJ: Relationship between internalization and cytotoxicity of ricin A-chain immunotoxins. *Br J Haematol* 70:289-94, 1988
64. Preijers FW, De Witte T, Rijke-Schilder GP, Tax WJ, Wessels JM, Haanen C, Capel PJ: Human T lymphocyte differentiation antigens as target for immunotoxins or complement-mediated cytotoxicity. *Scand J Immunol* 28:185-94, 1988
65. Preijers FW, Tax WJ, Wessels JM, Capel PJ, De Witte T, Haanen C: Different susceptibilities of normal T cells and T cell lines to immunotoxins. *Scand J Immunol* 27:533-40, 1988
66. Preijers FW, De Witte T, Wessels JM, Meyerink JP, Haanen C, Capel PJ: Cytotoxic potential of anti-CD7 immunotoxin (WT1-ricin A) to purge ex vivo malignant T cells in bone marrow. *Br J Haematol* 71:195-201, 1989
67. Van Horssen PJ: CD22-recombinant ricin A immunotoxin activity against malignant B-cells: Department of Hematology. Nijmegen, University Medical Center St Radboud, 1999, p 115
68. Hertler AA, Schlossman DM, Borowitz MJ, Laurent G, Jansen FK, Schmidt C, Frankel AE: A phase I study of T101-ricin A chain immunotoxin in refractory chronic lymphocytic leukemia. *J Biol Response Mod* 7:97-113, 1988
69. Hertler AA, Schlossman DM, Borowitz MJ, Blythman HE, Casellas P, Frankel AE: An anti-CD5 immunotoxin for chronic lymphocytic leukemia: enhancement of cytotoxicity with human serum albumin-monensin. *Int J Cancer* 43:215-9, 1989
70. LeMaistre CF, Rosen S, Frankel A, Kornfeld S, Saria E, Meneghetti C, Drajesk J, Fishwild D, Scanlon P, Byers V: Phase I trial of H65-RTA immunoconjugate in patients with cutaneous T-cell lymphoma. *Blood* 78:1173-82, 1991
71. Frankel AE, Laver JH, Willingham MC, Burns LJ, Kersey JH, Vallera DA: Therapy of patients with T-cell lymphomas and leukemias using an anti-CD7 monoclonal antibody-ricin A chain immunotoxin. *Leuk Lymphoma* 26:287-98, 1997
72. Stone MJ, Sausville EA, Fay JW, Headlee D, Collins RH, Figg WD, Stetler-Stevenson M, Jain V, Jaffe ES, Solomon D, Lush RM, Senderowicz A, Ghetie V, Schindler J, Uhr JW, Vitetta ES: A phase I study of bolus versus continuous infusion of the anti-CD19 immunotoxin, IgG-HD37-dgA, in patients with B-cell lymphoma. *Blood* 88:1188-97, 1996
73. Conry RM, Khazaeli MB, Saleh MN, Ghetie V, Vitetta ES, Liu T, LoBuglio AF: Phase I trial of an anti-CD19 deglycosylated ricin A chain immunotoxin in non-Hodgkin's lymphoma: effect of an intensive schedule of administration. *J Immunother Emphasis Tumor Immunol* 18:231-41, 1995
74. Uckun FM: Immunotoxins for the treatment of leukaemia. *Br J Haematol* 85:435-8, 1993
75. Vitetta ES, Stone M, Amlot P, Fay J, May R, Till M, Newman J, Clark P, Collins R, Cunningham D, Ghetie V, Uhr JW, Thorpe PE: Phase I immunotoxin trial in patients with B-cell lymphoma. *Cancer Res* 51:4052-8, 1991
76. Amlot PL, Stone MJ, Cunningham D, Fay J, Newman J, Collins R, May R, McCarthy M, Richardson J, Ghetie V, Ramilo O, Thorpe PE, Uhr JW, Vitetta ES: A phase I study of an anti-CD22-deglycosylated ricin A chain immunotoxin in the treatment of B-cell lymphomas resistant to conventional therapy. *Blood* 82:2624-33, 1993
77. Sausville EA, Headlee D, Stetler-Stevenson M, Jaffe ES, Solomon D, Figg WD, Herdt J, Kopp WC, Rager H, Steinberg SM, et al.: Continuous infusion of the anti-CD22 immunotoxin IgG-RFB4-SMPT-dgA in patients with B-cell lymphoma: a phase I study. *Blood* 85:3457-65, 1995

78. Grossbard ML, Freedman AS, Ritz J, Coral F, Goldmacher VS, Eliseo L, Spector N, Dear K, Lambert JM, Blattler WA, et al.: Serotherapy of B-cell neoplasms with anti-B4-blocked ricin: a phase I trial of daily bolus infusion. *Blood* 79:576-85, 1992
79. Grossbard ML, Lambert JM, Goldmacher VS, Spector NL, Kinsella J, Eliseo L, Coral F, Taylor JA, Blattler WA, Epstein CL, et al.: Anti-B4-blocked ricin: a phase I trial of 7-day continuous infusion in patients with B-cell neoplasms. *J Clin Oncol* 11:726-37, 1993
80. Grossbard ML, Gribben JG, Freedman AS, Lambert JM, Kinsella J, Rabinowe SN, Eliseo L, Taylor JA, Blattler WA, Epstein CL, et al.: Adjuvant immunotoxin therapy with anti-B4-blocked ricin after autologous bone marrow transplantation for patients with B-cell non-Hodgkin's lymphoma. *Blood* 81:2263-71, 1993
81. Grossbard ML, Niedzwiecki D, Nadler LM, et al: Anti-B4-blocked ricin (Anti-B4-bR) adjuvant therapy post-autologous bone marrow transplant (ABMT) (CALGB 9254): A phase III Intergroup study (abstr). *Proc Am Soc Clin Oncol* 17:A8, 1998
82. Tulpule A, Anderson LJJ, Levine AM, et al: Anti-B4 (CD19) monoclonal antibody, conjugated with ricin (B4-blocked ricin: B4-BR) in refractory AIDS lymphoma (abstr). *Proc Am Soc Clin Oncol* 13:A10, 1994
83. Scadden DT, Schenkein DP, Bernstein Z, Luskey B, Doweiko J, Tulpule A, Levine AM: Immunotoxin combined with chemotherapy for patients with AIDS-related non-Hodgkin's lymphoma. *Cancer* 83:2580-7, 1998
84. Engert A, Diehl V, Schnell R, Radszuhn A, Hatwig MT, Drillich S, Schon G, Bohlen H, Tesch H, Hansmann ML, Barth S, Schindler J, Ghetie V, Uhr J, Vitetta E: A phase-I study of an anti-CD25 ricin A-chain immunotoxin (RFT5-SMPT-dgA) in patients with refractory Hodgkin's lymphoma. *Blood* 89:403-10, 1997
85. Schnell R, Vitetta E, Schindler J, Borchmann P, Barth S, Ghetie V, Hell K, Drillich S, Diehl V, Engert A: Treatment of refractory Hodgkin's lymphoma patients with an anti-CD25 ricin A-chain immunotoxin. *Leukemia* 14:129-35, 2000
86. Kreitman RJ, Wilson WH, White JD, Stetler-Stevenson M, Jaffe ES, Giardina S, Waldmann TA, Pastan I: Phase I trial of recombinant immunotoxin anti-Tac(Fv)-PE38 (LMB-2) in patients with hematologic malignancies. *J Clin Oncol* 18:1622-36, 2000
87. Falini B, Bolognesi A, Flenghi L, Tazzari PL, Broe MK, Stein H, Durkop H, Aversa F, Corneli P, Pizzolo G, et al.: Response of refractory Hodgkin's disease to monoclonal anti-CD30 immunotoxin. *Lancet* 339:1195-6, 1992
88. Falini B, Pasqualucci L, Flenghi L, et al: Therapy of refractory Hodgkin's disease with anti-CD30/saporin immunotoxin. Fifth International Conference on Malignant Lymphoma. Lugano, Switzerland, 1993, June 9-12, p 66
89. Tepler I, Schwartz G, Parker K, Charette J, Kadin ME, Woodworth TG, Schnipper LE: Phase I trial of an interleukin-2 fusion toxin (DAB486IL-2) in hematologic malignancies: complete response in a patient with Hodgkin's disease refractory to chemotherapy. *Cancer* 73:1276-85, 1994
90. Bacha P, Fowlie A, Grimaldo D, et al: Antibody and pharmacokinetic analysis from a pivotal trial of ONTAK (DAB389IL-2) for the treatment of cutaneous T cell lymphoma (abstr). *Blood* 90:A1457, 1997
91. Kuzel T, Olsen E, Martin A, et al: Pivotal phase III trial of two dose levels of DAB389IL-2 (Ontak) for the treatment of mycosis fungoides (abstr). *Blood* 90:A2607, 1997
92. Bachier CR, LeMaistre CF: Immunotoxin therapy of Graft-versus-Host Disease, in Grossbard (ed): *Monoclonal antibody-based therapy of cancer*, vol. 15, Basic and clinical oncology series. New York, Marcel Dekker, 1998, p 213


**CYTOTOXICITY OF CD₃-RICIN A-CHAIN
IMMUNOTOXINS IN RELATION TO CELLULAR
UPTAKE AND DEGRADATION KINETICS**

Ypke V.J.M. van Oosterhout

Frank W.M.B. Preijers

Hans M.C. Wessels

Theo de Witte



Cancer Research 1992; 52: 5921-5925

ABSTRACT

The cytotoxicity of WT32 (CD3)-ricin A immunotoxin (IT) to the acute lymphoblastic leukemia T-cell line Jurkat was compared with the rate of internalization and intracellular degradation of WT32 and WT32-ricin A during continuous exposure. Moreover the influence of NH_4Cl and monensin on these processes was studied.

Based on protein synthesis inhibition ($[^3\text{H}]$ leucine incorporation), it appeared that cytotoxicity was not fully expressed directly after exposure to IT due to delay either in the internalization of membrane-bound IT or the action of intracellular ricin A. Varying the duration of incubation and postponing $[^3\text{H}]$ leucine addition up to 24 h after initiation showed that cytotoxicity occurred in two phases, rapid internalization of initially bound IT followed by a continuous but slower uptake possibly due to reexpression of the CD3 antigen.

No differences were found in the rate of internalization and degradation of ^{125}I -labeled WT32 and WT32-ricin A. Internalization started rapidly after binding at 37°C , was fastest during the first 12 h ($\sim 360,000$ molecules/cell) and continued for at least 24 h ($\sim 420,000$ molecules/cell). Exocytosis of intracellularly degraded molecules became measurable after 1 to 2 h of incubation at 37°C and increased up to approximately 400,000 molecules/cell in 24 h. After 4 h of incubation at 37°C the amount of internalized molecules exceeded the amount of WT32 that could maximally bind to the cell membrane ($\sim 150,000$ molecules/cell) confirming reexpression of antigen.

Addition of NH_4Cl and monensin enhanced the cytotoxicity of WT32-ricin A, probably due to an increased intracellular amount of IT. These agents appeared to reduce strongly the degradation of internalized WT32, resulting in an accumulation of intracellular molecules. NH_4Cl was most effective during the first 12 h of incubation, whereas monensin increased the amount of intracellular WT32 molecules after 2 to 24 h.

Our observations suggest that incubation conditions for the optimal cytotoxicity of IT treatment can be predicted by studying the internalization and degradation of the IT or respective monoclonal antibody.

INTRODUCTION

Monoclonal antibodies (MoAbs) coupled to toxins such as the A-chain of the phytolectin ricin appear to be highly specific ITs for the ex-vivo elimination of both malignant and normal T-cells which occur in autologous and allogeneic bone marrow (BM) grafts, respectively.¹⁻⁵ The efficacy of ricin A ITs depends on the number of molecules internalized by target cells and their intracellular routing and processing.⁶⁻¹¹ Ricin A-chain catalytically and irreversibly inhibits protein synthesis by cleaving the N-glycosidic bond of a single adenosine residue of 28S rRNA.¹² Theoretically, a cell can be killed by a single molecule that enters the cytosol¹³ but since the bulk of IT is usually transported into cellular compartments where ricin A is either proteolytically cleaved or inactivated by low pH^{7,14-16}, a sufficient number of molecules must be endocytosed to guarantee cytotoxicity.

Cells with a low density of antigen may survive treatment when the number of internalized IT molecules remains below the required threshold. This problem may be partly overcome by adding lysosomotropic amines (*e.g.*, ammonium chloride, chloroquine) or carboxylic ionophores (*e.g.*, monensin, nigericin) which enhance cytotoxicity by influencing different intracellular processes such as routing and metabolism.¹⁷⁻²¹ Alternatively, when the IT is continuously internalized as a result of reexpression of the target antigen, cytotoxic efficacy can be improved by prolonging incubation. However, this may be harmful to normal BM cells, and therefore it is important to determine the optimal period for individual ITs in purging BM grafts.

CD3-ricin A-chain ITs appear good candidates for the selective elimination of malignant and normal T lymphocytes.^{2,4,22,23} However, optimal incubation conditions remain to be determined. We therefore studied the kinetics of cytotoxicity of WT32-ricin A in the presence or absence of the enhancers ammonium chloride or monensin, and compared these with the intracellular uptake and processing of 125I-WT32 and 125I-WT32-ricin A.

MATERIALS AND METHODS

Cell line

The human acute lymphoblastic leukemia T-cell line Jurkat was cultured in RPMI 1640 (Flow Labs, Irvine, Scotland) supplemented with 10% heat-inactivated fetal calf serum (FCS) (Gibco, Paisley, Scotland), glutamine (2mM), penicillin (100 U/ml) and streptomycin (100 µg/ml) in a humidified incubator with 5% CO₂ in air at 37°C. Cells were maintained in the log phase.

Monoclonal antibody and immunotoxin preparation

Murine anti-human T-cell MoAb WT32 (IgG2a, CD3)²⁴ was purified from ascitic fluid by ammonium sulfate precipitation and affinity chromatography with staphylococcal protein A-sepharose (Pharmacia, Uppsala, Sweden). WT32 was conjugated on a 1 to 1.1 ratio to ricin A using *N*-succinimidy-3-(2-pyridyldithio)propionate (SPDP) (Pharmacia) and assayed as described elsewhere.⁹ Antibody-binding activity was confirmed by flow cytometry (FCM) and enzyme-linked immunosorbent assay.⁹ Nonspecific cytotoxicity was tested on cell lines which did not express the CD3-antigen.⁹ Ricin A was kindly provided by Dr. F.K. Jansen (Centre de Recherches Clin Midy, Montpellier, France).

Radioiodination

WT32 and WT32-ricin A were labeled with ¹²⁵I (Amersham International, UK) using the Chloramine-T method (Hunter, 1973), and then purified by chromatography on a Sephadex G25 column (PD-10; Pharmacia). Eluted fractions containing ¹²⁵I-WT32 and ¹²⁵I-WT32-ricin A were pooled, and molecules smaller than *M_r* 30,000 were removed by means of a centrprep-30 concentrator (Grace B.V., Amicon Division, Rotterdam, Holland). Thereafter radioactivity due to free ¹²⁵I and fragments smaller than *M_r* 30,000 was less than 0.3% of the total radioactivity. The concentration of the labeled antibody and its ricin A conjugate was determined by radioimmunoassay, while immunoreactivity was assayed by means of binding limiting quantities of labeled WT32 and WT32-ricin A. Conjugation of ricin A to the antibody was confirmed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) of ¹²⁵I-labeled IT, followed by autoradiography and scanning densitometry (data not shown).

Cytotoxicity of WT32-ricin A

Culture medium containing 10⁶ cells/ml was incubated at 37°C with WT32 ricin A (10⁻⁸ M) with and without either 6 mM NH₄Cl or freshly prepared 100 nM monensin (Sigma, St. Louis, MO). After 1, 2, 4, 8, 16, and 24 h of incubation, cells were washed with RPMI 1640 containing 1% FCS and then resuspended in fresh medium. They were distributed in triplicate into 96-well microtiter plates (Costar, Cambridge, MA) at a concentration of 10⁵ cells/well. Subsequently, cells were immediately incubated with 1 µCi [³H]leucine (Amersham) at 37°C for 1 h, following which they were harvested. The radioactivity was determined, and protein synthesis was expressed as the percentage [³H]leucine incorporation of untreated cells after correcting for non-specific incorporation in the presence of 1 mM cycloheximide.

Cytotoxicity caused by continuous internalization of WT32-ricin A

The increase of cytotoxicity after prolonging incubation might be explained by delayed internalization of IT already bound soon after exposure or slow onset of cytotoxicity. To eliminate differences in cytotoxicity caused by these phenomena rather than by continuous internalization of IT, the assay was modified by introducing an IT-free incubation after cells were washed up to 24 h after the initiation of the experiment (23, 22, 20, 16, 8, and 0 h, respectively), and then proceeding with [³H]leucine incorporation.

Intracellular and surface bound ^{125}I -WT32 and ^{125}I -WT32-ricin A

Following continuous incubation of Jurkat cells with ^{125}I -labeled antibody or its ricin A conjugate, the number of intracellular molecules was determined using a modification of the procedure previously described.⁹ Briefly, a suspension of 5×10^5 target cells was incubated with 10^{-8} M ^{125}I -WT32 or ^{125}I -WT32-ricin A for various times at either 37°C or 4°C at a final volume of 100 μl . Thereafter cells were washed and counted to determine the total amount of associated radioactivity. Subsequently, cells were incubated with isotonic glycine buffer (pH 2) at 4°C for 30 min, to remove surface-bound molecules and then washed, after which radioactivity was again determined. The absolute number of intracellular molecules was calculated from the glycine buffer-treated cells initially incubated at 37°C after correcting for the cells treated with the same buffer at 4°C.

The number of molecules bound to the cell surface was determined by subtracting the amount of intracellular radioactivity from total radioactivity after incubation at 37°C. The maximal number of binding sites was determined by incubation with increasing amounts of ^{125}I -WT32 at 4°C for 1 h.

Degradation of ^{125}I -WT32 and ^{125}I -WT32-ricin A

The degraded and subsequently exocytosed amount of internalized molecules was determined by measuring radioactivity of the trichloroacetic acid (TCA)-soluble fraction of culture supernatant. After incubation with ^{125}I -WT32 or ^{125}I -WT32-ricin A at 37°C or 4°C, cells were pelleted by centrifugation. Seventyfive μl of supernatant was treated with 1 ml of 25% TCA and 75 μl 1% bovine serum albumin (BSA) at 4°C for 15 min and then centrifuged at 1500 G for 15 min. The radioactivity of the soluble fraction from cells incubated at 4°C was subtracted from that obtained at 37°C to remove any background radioactivity.

Extracellular degradation was excluded since there was negligible radioactivity found in the TCA-soluble fraction of B-cell line Daudi, which does not express CD3, and of purified surface membranes from disrupted Jurkat cells after incubation with ^{125}I -WT32 or ^{125}I -WT32-ricin A as described by Press *et al.*^{19,25} Similar results were obtained after incubating Jurkat with a ^{125}I -labeled control antibody (CD19, IgG2a) or its ricin A conjugate (data not shown), suggesting that no extracellular degradation occurred during incubation.

Internalization ^{125}I -WT32 and ^{125}I -WT32-ricin A

Internalization was defined by the total number of molecules transported across the cell membrane during incubation. The degree of internalization was calculated by adding the fraction which was intracellularly degraded and subsequently exocytosed to the amount of intracellular molecules.

RESULTS

Kinetics of cytotoxicity of WT32-ricin A directly after IT treatment

Prolonging IT treatment led to a reduction in [^3H]leucine incorporation from 85% after 1 h to 9% after 24 h (Figure 1). This might be explained by continuous binding and internalization of IT. Alternatively it might be the result of slow internalization of IT bound to cells in the first phase of incubation or of a delay in the toxicity of intracellular ricin A molecules.




Figure 1

Kinetics of cytotoxicity of WT32-ricin A

Jurkat cells were incubated at 37°C with 10^{-8} M WT32-ricin A for various periods. After IT treatment cells were washed, and directly after (■) or 24 h after (▲) initiation of the experiment they were incubated with [^3H]leucine for 1 h. Protein synthesis is expressed as a percentage of [^3H]leucine incorporation of untreated cells. Untreated cells incorporated ~20,000 cpm; background incorporation was ~100 cpm. Assays were performed in triplicate; SDs were less than 10%.

Cytotoxicity caused by continuous internalization

Introduction of an IT-free incubation up to 24 h after the initiation of the experiment resulted in an additional decrease of protein synthesis which became less pronounced after prolonged treatment (Fig. 1). The kinetics of protein synthesis measured after the post-IT-treatment period showed a biphasic pattern: a considerable decrease of protein synthesis during the first 2 h of IT treatment (60%), followed by a slower decrease for up to 24 h (89% after 24 h). In this setting the reduction of protein synthesis could only be the result of continuous internalization of IT.

Influence of enhancers of cytotoxicity

The presence of NH_4Cl as well as of monensin resulted in significantly greater decrease of protein synthesis after 1 h (Figure 2), although monensin appeared to be slightly more effective than NH_4Cl (83%, 74% and 47% reduction with monensin, NH_4Cl and without enhancers, respectively). Similar concentrations of NH_4Cl or monensin in the absence of IT were non toxic since [^3H]leucine incorporation was unaffected (data not shown). Enhancement of cytotoxicity became less pronounced after prolonging incubation in the presence of IT.




Figure 2
Influence of cytotoxicity enhancers

Jurkat cells were incubated at 37°C with 10^{-8} M WT32-ricin A for various periods in the absence (▲) or presence (●) of 6 mM NH_4Cl or 100 nM monensin (□). After IT treatment cells were washed, followed by an incubation in the absence of IT and enhancers up to 24 h after initiation of the experiment, and then by a 1-h uptake of [^3H]leucine. Protein synthesis is expressed as percentage [^3H]leucine incorporation of untreated cells. Untreated cells incorporated ~20,000 cpm; background incorporation was ~100 cpm. Assays were performed in triplicate; SDs were less than 10%.

Kinetics of internalization and intracellular degradation of ^{125}I -WT32 and ^{125}I -WT32-ricin A

To confirm the possibility of continuous internalization due to reexpression of antigen as suggested by the pattern of protein synthesis inhibition, the kinetics of internalization and intracellular degradation were studied by means of ^{125}I -labeled WT32 and its ricin A conjugate. The maximal number of binding sites was determined with increasing concentrations of ^{125}I -WT32 at 4°C and was achieved with 4 $\mu\text{g}/\text{ml}$ ^{125}I -WT32 (~150,000 molecules) (Figure 3).




Figure 3
Binding of ^{125}I -WT32 to Jurkat cells at 4°C

Jurkat cells were incubated with 1, 2, 4, 6 and 8 $\mu\text{g}/\text{ml}$ ^{125}I -WT32 at 4°C for 1 h. After incubation unbound ^{125}I -WT32 was washed out, and the number of cell-associated ^{125}I -WT32 molecules was determined (●). Assays were performed in triplicate; SDs were less than 10%.

Both ^{125}I -WT32 and ^{125}I -WT32-ricin A were internalized at the same rate (Figure 4). At 37°C ^{125}I -labeled antibody and its ricin A conjugate were rapidly internalized during the first 12 h (~360,000 molecules); thereafter internalization continued at a slower rate for up to at least 24 h (between 420,000 and 440,000 molecules). After 4 h the number of internalized molecules surpassed the amount of ^{125}I -WT32 that could maximally bind




Figure 4 Internalization and intracellular degradation of WT32 and WT32-ricin A

Jurkat was incubated at 37°C or 4°C with ^{125}I -labeled WT32 (A) or WT32-ricin A (B) for various periods. Results of cells incubated at 37°C corrected for cells treated at 4°C, as described in "Material and Methods", are depicted. The number of internalized and subsequently degraded and exocytosed molecules was measured by counting TCA-soluble radioactivity of the supernatant of incubation medium (Δ). The number of surface bound molecules was measured by counting cell bound radioactivity that was removable by glycine buffer pH2 (\square). The intracellular number of molecules was measured by counting cell associated radioactivity not-removable by the glycine buffer (\bullet). Absolute amount of internalized molecules was determined as the sum of intracellular molecules and intracellular broken down and subsequently exocytosed molecules (\blacktriangle). One of three (A) and one of five (B) concordant experiments are shown.

to the cell membrane ($\sim 150,000$), confirming the continuous reexpression of antigen. The amount of membrane-bound molecules at 37°C ($\sim 60,000$ after 2 h) decreased slightly during the 24-h period. Binding of ^{125}I -WT32 and ^{125}I -WT32-ricin A at the concentration used was optimal after 2 h at 4°C and remained unchanged up to 24 h (data not shown).

The number of intracellular molecules reached a maximum after a 2-h incubation at 37°C ($\sim 80,000$ molecules) and declined to about 18,000 molecules after 24 h.

Breakdown and exocytosis of internalized molecules were measurable after 1 h and increased to about 330,000 molecules in 12 h. Thereafter the exocytosis rate was reduced but still continued up to about 400,000 molecules after 24 h.

Influence of cytotoxicity enhancers on the internalization and intracellular degradation

To determine the influence of the cytotoxicity enhancers on the uptake and intracellular degradation of ^{125}I -labeled WT32 and its ricin A conjugate at 37°C, the above-described experiments were repeated with ^{125}I -WT32 in the presence of NH_4Cl (Figure 5A) or monensin (Figure 5B).

The addition of NH_4Cl resulted in a considerable increase in the amount of intracellular ^{125}I -WT32, with a maximum of at least 140,000 molecules between 4 and 8 h. There-




Figure 5 Internalization and intracellular degradation of WT32

Jurkat was incubated at 37°C or 4°C with ^{125}I -WT32 for various periods in the presence of 6mM NH_4Cl (A) or 100 nM monensin (B). Results from cells incubated at 37°C corrected for cells treated at 4°C, as described in "Material and Methods", are depicted. Exocytosis of degraded WT32 (△), surface-bound WT32 (□), intracellular WT32 (●) and internalized WT32 (▲) were determined. For additional legends see Fig. 4. One of four (A) and one of three (B) concordant experiments are shown.

after this increase diminished, almost disappearing after 24 h of incubation (Figures 4 and 5A).

Monensin also resulted in an increase of intracellular ^{125}I -WT32, but this was less pronounced during the first 8 h. After 4 h the amount of intracellular ^{125}I -WT32 reached a peak of approximately 100,000 molecules, which gradually declined to 80,000 molecules after 24 h of incubation. However this was larger than the yield either with NH_4Cl or without enhancer after 24 h.

The absolute number of internalized molecules in the presence of NH_4Cl was slightly decreased in comparison to the conditions without enhancers, suggesting that NH_4Cl did not influence the internalization of ^{125}I -WT32 (~380,000 and 410,000 molecules after 24 h, with and without NH_4Cl , respectively). The addition of monensin resulted in a larger reduction in the number of molecules entering the cell (~330,000 molecules after 24 h).

DISCUSSION

The cytotoxicity of WT32-ricin A was not fully expressed directly due to a delay in either the internalization of initially bound IT or the action of intracellular ricin A. This has also been reported by Ravel *et al.*²⁰ and Sung *et al.*²⁶, using T101-ricin A (CD5) and 454-ricin A (CD71), respectively. They observed that cytotoxicity increased when measuring protein synthesis was postponed. This phenomenon can strongly influence the outcome of kinetics studies for the determination of the optimal incubation time. To determine the efficacy of prolonging IT-treatment, it is therefore important to standardize the period between initiation of IT incubation and the moment of [³H]leucine exposure. This was assessed by introducing an IT-free incubation up to 24 h after the initiation of the experiment. The reduction of protein synthesis in the current setting showed a biphasic pattern of strong inhibition during the first period followed by a slower inhibition during the second period. This suggests a rapid internalization of initially bound WT32-ricin A during the first period of IT treatment, followed by a more slowly continuous uptake of WT32-ricin A, possibly due to reexpression of CD3 antigen.

In order to confirm the continuous uptake of IT due to reexpression of CD3 antigen, binding, internalization and degradation of WT32 as well as WT32-ricin A were examined after continuous exposure of Jurkat cells to ¹²⁵I-labeled ligands. Free MoAb and the respective IT were both studied because conjugation of ricin A to the MoAb may affect the internalization or intracellular routing of the complex, depending on the target antigen. It has been reported that conjugation of RFT11 (CD2) to ricin A decreases the intracellular degradation of ligand.¹⁰ In this study we show that there is neither difference in internalization nor in intracellular degradation between WT32 and WT32-ricin A. This has also been described for other MoAbs and ricin A ITs.^{6,9,11,27}

Press *et al.*¹⁹ and Geissler *et al.*¹⁸ studied the internalization and degradation of CD3-MoAb ¹²⁵I-64.1 in T-cells and the T-cell line HPB-ALL after initial pulse labeling of the cell surface with ¹²⁵I-64.1 at 4°C for 1 h. Washing out unbound ligand prior to incubation at 37°C, however, would reduce the amount that can be maximally internalized due to: (a) disturbing the equilibrium between free and bound ligand, resulting in the dissociation of membrane-bound ligand; and (b) preventing the binding and internalization of ligand induced by reexpression of the antigen. We therefore performed internalization studies in the continuous presence of an excess of free MoAb or IT in the medium. We were able to demonstrate the reexpression of CD3 antigen or the expression of newly synthesized CD3 antigen by the observation that the amount of internalized ligands exceeded 2.5 times the maximal number of antigen binding sites under saturating conditions in 24 h. This is in accordance with observations of Krangel²⁸, who demonstrated a recycling of the CD3/TCR complex to the cell membrane following

endocytosis in the presence of CD3 MoAb. The capacity of the CD3 antigen to induce the internalization of potential cytotoxic ITs for up to 24 h resembles the observation that prolongation of treatment with WT32-ricin A up to 24 h increases cytotoxicity.

Cytotoxicity of WT32-ricin A to T-cells and T-cell lines is enhanced by NH_4Cl ^{9,22,23} and monensin. Although the mechanism of potentiating is not fully understood, different intracellular compartments and transport mechanisms might be influenced (reviewed in Ref. 29). Previous studies have shown that NH_4Cl and monensin considerably decreased the degradation of 64.1(CD3) by T cells and the T-cell line HPB-ALL.^{18,19,24} We demonstrated that NH_4Cl and monensin reduced the degradation of intracellular ¹²⁵I-WT32 by Jurkat but found differences in their action. Intracellular degradation was decreased during different periods of the continuous incubation, resulting in a temporary accumulation of MoAb. NH_4Cl was most effective during the first 12 h of incubation, whereas monensin decreased the degradation of intracellular ¹²⁵I-WT32 after 2 h to at least 24 h of incubation. Moreover, NH_4Cl resulted in a higher intracellular accumulation of ligand. As was published before, NH_4Cl did not influence the total amount of internalization.^{9,27} Monensin, however, reduced the number of internalized molecules in contrast to observations of Ravel *et al.*²⁷, who found that monensin had no influence on internalization. Though these results seem to be in favor of NH_4Cl as an enhancing agent, monensin may act more effectively on intracellular handling of ricin A than NH_4Cl , which was confirmed in the cytotoxicity kinetics, since monensin was at least as effective as NH_4Cl .

The efficacy of WT32-ricin A might be extended to cells with low antigen density when treatment is prolonged and either NH_4Cl or monensin is added due to continuous uptake and intracellular accumulation of ITs. Since exposure to ricin A-ITs for 24 h is not harmful to either bone marrow (BM) progenitor cells or stem cells²³, it may now be possible to improve considerably the purging of BM intended for autologous transplantation and therefore the success of this form of therapy.

ACKNOWLEDGEMENT

We would like to thank Peter Donnelly for helpful comments and suggestions during the preparation of this manuscript.

REFERENCES

1. Fauser AA, Shustik C, Langleben A, Laurent G, Kanz L, Spurl GM, Price G, Ahlgren PD, Cooper BA: T-cell depletion with ricin A-chain T101 in allogeneic bone marrow transplantation to prevent severe graft-versus-host disease. *Clin Invest Med* 11:40-6, 1988
2. Filipovich AH, Valleria DA, Youle RJ, Haake R, Blazar BR, Arthur D, Neville DM, Jr., Ramsay NK, McGlave P, Kersey JH: Graft-versus-host disease prevention in allogeneic bone marrow transplantation from histocompatible siblings. A pilot study using immunotoxins for T cell depletion of donor bone marrow. *Transplantation (Baltimore)* 44:62-9, 1987
3. Laurent G, Maraninchi D, Gluckman E, Vernant JP, Derocq JM, Gaspard MH, Rio B, Michalet M, Reiffers J, Dreyfus F, et al.: Donor bone marrow treatment with T101 Fab fragment-ricin A-chain immunotoxin prevents graft-versus-host disease. *Bone Marrow Transplant* 4:367-71, 1989
4. Martin PJ, Hansen JA, Torok-Storb B, Moretti L, Press O, Storb R, Thomas ED, Weiden PL, Vitetta ES: Effects of treating marrow with a CD3-specific immunotoxin for prevention of acute graft-versus-host disease. *Bone Marrow Transplant* 3:437-44, 1988
5. Preijers FW, De Witte T, Wessels JM, De Gast GC, Van Leeuwen E, Capel PJ, Haanen C: Autologous transplantation of bone marrow purged in vitro with anti-CD7-(WT1)-ricin A immunotoxin in T-cell lymphoblastic leukemia and lymphoma. *Blood* 74:1152-8, 1989
6. Byers VS, Pawluczyk IZ, Hooi DS, Price MR, Carroll S, Embleton MJ, Garnett MC, Berry N, Robins RA, Baldwin RW: Endocytosis of immunotoxin-791T/36-RTA by tumor cells in relation to its cytotoxic action. *Cancer Res* 51:1990-5, 1991
7. Chignola R, Colombatti M, Dell'Arciprete L, Candiani C, Tridente G: Distribution of endocytosed molecules to intracellular acidic environments correlates with immunotoxin activity. *Int J Cancer* 46:1117-23, 1990
8. May RD, Wheeler HT, Finkelman FD, Uhr JW, Vitetta ES: Intracellular routing rather than cross-linking or rate of internalization determines the potency of immunotoxins directed against different epitopes of sIgD on murine B cells. *Cell Immunol* 135:490-500, 1991
9. Preijers FW, Tax WJ, De Witte T, Janssen A, v.d. Heijden H, Vidal H, Wessels JM, Capel PJ: Relationship between internalization and cytotoxicity of ricin A-chain immunotoxins. *Br J Haematol* 70:289-94, 1988
10. Press OW, Martin PJ, Thorpe PE, Vitetta ES: Ricin A-chain containing immunotoxins directed against different epitopes on the CD2 molecule differ in their ability to kill normal and malignant T cells. *J Immunol* 141:4410-7, 1988
11. Wargalla UC, Reisfeld RA: Rate of internalization of an immunotoxin correlates with cytotoxic activity against human tumor cells. *Proc Natl Acad Sci U S A* 86:5146-50, 1989
12. Endo Y, Tsurugi K: The RNA N-glycosidase activity of ricin A-chain. The characteristics of the enzymatic activity of ricin A-chain with ribosomes and with rRNA. *J Biol Chem* 263:8735-9, 1988
13. Olsness S, Phil A: Chimeric toxins. *Pharmacol Ther* 15:355, 1982
14. Carriere D, Casellas P, Richer G: Immuntoxins: ultrastructural cytotoxic effect of ricin and immunotoxin on lymphoma WEHI-7 cells. Antibodies as carriers of anticancer drugs or toxins. Satellite Symposium of the IV International Congress of Immunology. Paris: Quo Vadis, 1980
15. Press OW, Vitetta ES, Farr AG, Hansen JA, Martin PJ: Evaluation of ricin A-chain immunotoxins directed against human T cells. *Cell Immunol* 102:10-20, 1986
16. Carriere D, Casellas P, Richer G, Gros P, Jansen FK: Endocytosis of an antibody ricin A-chain conjugate (immuno-A-toxin) adsorbed on colloidal gold. Effects of ammonium chloride and monensin. *Exp Cell Res* 156:327-40, 1985
17. Casellas P, Bourrie BJ, Gros P, Jansen FK: Kinetics of cytotoxicity induced by immunotoxins. Enhancement by lysosomotropic amines and carboxylic ionophores. *J Biol Chem* 259:9359-64, 1984

18. Geissler F, Anderson SK, Press O: Intracellular catabolism of radiolabeled anti-CD3 antibodies by leukemic T cells. *Cell Immunol* 137:96-110, 1991
19. Press OW, DeSantes K, Anderson SK, Geissler F: Inhibition of catabolism of radiolabeled antibodies by tumor cells using lysosomotropic amines and carboxylic ionophores. *Cancer Res* 50:1243-50, 1990
20. Ravel S, Casellas P: Internalization of the cytotoxic molecules of T101 F(ab')₂-(ricin-A-chain) immunotoxin into human T-leukemic cells. *Eur J Biochem* 192:469-73, 1990
21. Sandvig K, Olsnes S: Entry of the toxic proteins abrin, modeccin, ricin and diphtheria toxin into cells. II. Effect of pH, metabolic inhibitors, and ionophores and evidence for toxin penetration from endocytotic vesicles. *J Biol Chem* 257:7504-13, 1982
22. Preijers FW, Tax WJ, Wessels JM, Capel PJ, De Witte T, Haanen C: Different susceptibilities of normal T cells and T cell lines to immunotoxins. *Scand J Immunol* 27:533-40, 1988
23. Preijers FW, De Witte T, Rijke-Schilder GP, Tax WJ, Wessels JM, Haanen C, Capel PJ: Human T lymphocyte differentiation antigens as target for immunotoxins or complement-mediated cytotoxicity. *Scand J Immunol* 28:185-94, 1988
24. Tax WJM, Willems HW, Reekers PPM, Capel PJA, Koene RAP: Polymorphism in mitogenic effect of IgG1 monoclonal antibodies against T₃ antigen on human T cells. *Nature (Lond.)* 304:445-7, 1983
25. Press OW, Hansen JA, Farr A, Martin PJ: Endocytosis and degradation of murine anti-human CD3 monoclonal antibodies by normal and malignant T-lymphocytes. *Cancer Res* 48:2249-57, 1988
26. Sung C, Wilson D, Youle RJ: Comparison of protein synthesis inhibition kinetics and cell killing induced by immunotoxins. *J Biol Chem* 266:14159-62, 1991
27. Ravel S, Colombatti M, Casellas P: Internalization and intracellular fate of anti-CD5 monoclonal antibody and anti-CD5 ricin A-chain immunotoxin in human leukemic T cells. *Blood* 79:1511-7, 1992
28. Krangel MS: Endocytosis and recycling of the T₃-T cell receptor complex. *J Exp Med* 165:1141-59, 1987
29. Casellas P, Jansen FK: Immunotoxin enhancers. In: A. E. Frankel (ed.), *Immunotoxins*, pp. 351-369. Boston, Kluwer, 1988

**EFFECT OF ISOTYPE ON
INTERNALIZATION AND CYTOTOXICITY
OF CD19-RICIN A-IMMUNOTOXINS**

Ypke V.J.M. van Oosterhout

Ingrid E. van den Herik-Oudijk


Hans M.C. Wessels

Theo de Witte

Jan G. J. van de Winkel

Frank W.M.B. Preijers

Cancer Research 1994; 54: 3527-3532



ABSTRACT

We analyzed the effect of isotype variation on effectiveness of B-cell CD19 immunotoxins (ITs) by using class switch variants (CLB-B4-IgG1 and CLB-B4-IgG2a) conjugated to ricin A. Notably, IgG1-IT appeared to be ~100-fold more potent than IgG2a-IT toward B-cell lines Daudi and KM3. Binding and internalization studies with ^{125}I -labelled monoclonal antibodies (MoAbs) revealed a higher cellular uptake of IgG1 compared to IgG2a, despite similar binding affinities. Following removal of the Fc-part, both MoAb internalized at the same rate as IgG2a, indicating that the Fc part of IgG1 is involved in enhanced cellular uptake. The involvement of Fc γ RII (CD32) in this process was demonstrated by a decreased cytotoxicity of IgG1-IT (and not IgG2a-IT) in the presence of Fc γ RII-blocking MoAbs AT10 or IV.3. To identify the isoform responsible for this phenomenon, internalization of IgG1 and IgG2a in 11 B-cell lines, and malignant B-cells of 8 patients was compared with expression of Fc γ RII subclasses. In addition to four cell lines (Daudi, KM3, Nalm6, and Raji), the malignant B-cells of two patients showed enhanced uptake of IgG1 relative to IgG2a. Only the Fc γ RIIa transcript was found in all B-cells. Furthermore, enhanced uptake of IgG1 correlated with rosetting of erythrocytes sensitized with anti-glycophorin A MoAb of IgG1 isotype rather than with Fc γ RIIa membrane expression levels. These data support the idea that functional Fc γ RIIa is involved in the enhanced IgG1 uptake observed in a subset of B-cells. Our study, therefore, points to an important role for the Fc region of IT in delivery of cytotoxic effects.

INTRODUCTION

During the past decade, a large number of MoAbs have been conjugated to the highly toxic A chain of the phytolectin ricin in an effort to create tumor-toxic agents. These may be of use for treatment of bone marrow in vitro or may be useful in vivo for patients suffering from hematological malignancies (reviewed in Refs. 1-3). The killing potencies

of such ricin A-chain ITs appeared to vary widely. Firstly, the cytotoxic efficacy of IT depends upon the number of IT molecules internalized by target cells, as well as the intracellular route followed upon internalization.⁴⁻⁶ Both factors are influenced by the nature of the target cell, type of antigen^{7,8}, target antigen density^{6,9}, MoAb binding affinity^{9,10}, and the IT epitope¹¹⁻¹³. Furthermore, the biological effects of ITs have been observed to vary depending upon the use of F(ab')₂, F(ab'), or intact IgG.¹⁴⁻¹⁶ The effect of variation in isotype on the cytotoxic effectiveness of ITs remained unaddressed.

We constructed two anti-CD19 ITs, which differ in isotype only, by conjugating the murine anti-CD19 MoAb CLB-B4-IgG1 and its isotype-switch variant CLB-B4-IgG2a¹⁷ to ricin A. These ITs differed only in their Fc-parts and expressed identical binding affinities. Interestingly, however, a profound difference in their cytotoxic efficacy was observed using B-cell lines such as Daudi and KM3 as targets. We analyzed the mechanisms underlying the difference in effectiveness between these two IT. Our data point to a role of significant role for the IIa subclass of FcγRII (CD32) molecules expressed on B cells.

MATERIALS AND METHODS

Malignant B-cells and cell lines

Malignant B-cells were obtained from bone marrow or blood of B-cell acute lymphocytic leukemia patients after informed consent. B-cell acute lymphocytic leukemia cells were enriched by Ficoll (1.077 g/ml) density gradient centrifugation. The isolated cell populations consisted of more than 90% of CD19-positive cells as determined by flow cytometric analysis (FCM) (Coulter Epics Elite, Hialeah, FL). The cells were suspended in culture medium and were either used directly or cryopreserved (with 10% DMSO) without loss of functional potential.¹⁸

The human B-cell lines CRL, Daudi, DOHH2, Fravel, KM3, Nalm6, Raji, Ramos, Ros-1, RPMI, and the T-cell line Jurkat were cultured in a culture medium consisting of RPMI 1640 medium (Flow, Irvine, Scotland) supplemented with 10% heat-inactivated fetal calf serum (FCS) (Gibco, Paisley, Scotland), glutamine (2mM), penicillin (100 units/ml) and streptomycin (100 μg/ml) in a humidified incubator with 5% CO₂ in air at 37°C. 3T6 fibroblasts expressing human FcγRIIa, FcγRIIb₁, or FcγRIIb₂ molecules were grown in culture medium supplemented with geneticin (0.8 mg/ml; GIBCO BRL, Grand Island, NY).

Monoclonal antibodies

Murine anti-CD19 MoAb CLB-B4-IgG1, and isotype switch-variant CLB-B4-IgG2a¹⁷ were kindly provided by Dr I. Slaper (CLB, Amsterdam, The Netherlands). Hybridoma cells secreting TS2.18 (CD2 and IgG1)¹⁹ and 35.1 (CD2 and IgG2a)²⁰ were purchased from

American Type Culture Collection (Rockville, MD). The CD2-MoAbs were used as isotype-matched controls. MoAbs were purified from ascites fluid by ammonium sulfate precipitation and affinity chromatography with staphylococcal protein A-sepharose (Pharmacia, Uppsala, Sweden).

MoAb 197 (IgG2a)²¹ was used to block FcγRI (CD64), and was obtained from Medarex (Clinton, NJ). IgG and F(ab')₂ fragments of MoAb AT10 (IgG1)²², generously provided by Dr M. Glennie (Tenovus Research Laboratory, Southampton, United Kingdom), and IV.3 (IgG2b)²³ from Medarex served to inhibit binding to FcγRII (CD32).

Preparation of F(ab')₂ fragments

F(ab')₂ fragments of CLB-B4-IgG1 and CLB-B4-IgG2a were prepared using a F(ab')₂ preparation kit (Pierce, Rockford, IL) according to the manufacturer's protocol. Briefly, CLB-B4-IgG1 and CLB-B4-IgG2a were incubated with immobilized pepsin at pH 4.2 (20 mM sodium acetate buffer) for 4 and 16 h, respectively. Undigested IgG molecules and Fc-fragments were removed by affinity chromatography with protein A-Sepharose. Remaining fragments smaller than *M_r* 30,000 were removed by means of a Centrprep 30 concentrator (Amicon, Beverly, MA). The purity of F(ab')₂ fragments was determined by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), which revealed less than 1% contamination with either intact IgG or Fc fragments (data not shown).

IT preparation

CLB-B4-IgG1, CLB-B4-IgG2a, TS2.18, and 35.1 were conjugated to ricin A (kindly provided by Dr. F. K. Jansen, Centre de Recherches Clin Midy, Montpellier, France) using *N*-succinimide 3-(2-pyridyldithio)propionate (SPDP) (Pharmacia), as described.⁶ The conjugation ratios of ricin A to MoAb were estimated by measurement of absorbance at 280 nm and radioimmunoassay (RIA) and were determined to be 0.9, 1.1, 0.9, and 0.8 for CLB-B4-IgG1, CLB-B4-IgG2a, TS2.18, and 35.1, respectively. Preservation of antibody-binding activity following conjugation was assessed by FCM.

Cytotoxic potential of IT

Cells in culture medium were distributed into 96-well microtiter plates (Costar, Cambridge, MA) at a concentration of 10⁵ cell/well and incubated in triplicate at 37°C with varying concentrations of IT (10⁻⁸-10⁻¹³ M) in the presence or absence of 6 mM NH₄Cl. After 16 h of incubation, 0.5 μCi [³H]leucine (Amersham, Buckinghamshire, United Kingdom) was added, and cells were further incubated at 37°C for another 24 h and harvested. The radioactivity was quantitated, and protein synthesis was expressed as the percentage [³H]leucine incorporation of cells cultured at 37°C without IT.⁶

In some of the cytotoxicity assays, FcγRs were blocked by pre-incubation of target cells with MoAb 197 (purified IgG, 2 μg/ml), AT10 (ascites fluid, diluted 1:500) or IV.3 (purified IgG, 2 μg/ml) for 45 min at 4°C. Cells were washed, and cytotoxicity was assayed

as above. Effectiveness and specificity of blocking with these MoAbs was verified by inhibition of EA-rosetting (Refs. 24 and 25; data not shown).

Radioiodination

MoAbs CLB-B4-IgG1 and CLB-B-4-IgG2a (intact IgG and F(ab')₂-fragments), TS2.18 and 35.1 were labelled with Na¹²⁵I (Amersham) using the Chloramine-T method²⁶, and purified by chromatography on Sephadex G25 (PD-10; Pharmacia). Eluted fractions containing ¹²⁵I-IgG and ¹²⁵I-F(ab')₂-fragments, respectively, were pooled, and molecules smaller than *M_r* 30,000 were removed by Centriprep 30 concentration. Radioactivity due to free ¹²⁵I and fragments smaller than *M_r* 30,000 were always less than 0.3% of total radioactivity. The concentration of labelled IgG and F(ab')₂ fragments was determined by Mancini assays²⁷, and specific binding activity was assessed by performing binding studies with limiting amounts of labelled IgG and F(ab')₂ fragments.²⁸

Scatchard analysis

Suspension of 5×10^5 Daudi cells were incubated with various concentrations of ¹²⁵I-labelled IgG or F(ab')₂ fragments for 2 h at 4°C (final volume, 100 µl). Thereafter, cells were washed, and radioactivity was counted to determine the number of cell-associated molecules. Binding to Daudi cells not mediated by antigen was assessed in the presence of a 100-fold excess of unlabelled antibody and was less than 5% of total cell-associated radiolabel for both CD19 MoAbs. Nonspecific binding to incubation tubes was assessed by performing the experiment in the absence of targets. The affinity constant (*K_a*) and the number of bound molecules/cell under saturating conditions were calculated according to Scatchard.²⁹

Internalization of ¹²⁵I-IgG and ¹²⁵I-F(ab')₂

The cellular uptake and processing of ¹²⁵I-labelled IgG, or F(ab')₂ fragments, were determined as described.²⁸ Briefly, a suspension of 5×10^5 cells was incubated with 10^{-8} M ¹²⁵I-IgG or ¹²⁵I-F(ab')₂ in culture medium at 37°C (final volume, 100 µl). Following different times, cells were pelleted, and the amount of (internalized, and subsequently) degraded and exocytosed MoAb was estimated by measurement of TCA (25%)-soluble radioactivity of the supernatant. Cells were then washed twice and assessed for total cell-associated (surface-bound, and intracellular) radioactivity. Surface radioactivity was subsequently released by incubation with isotonic glycine-HCl buffer (pH2; 4°C for 30 min); then cells were washed, and intracellular radioactivity was determined. Nonantigen-mediated binding and internalization of ¹²⁵I-ligand was assessed in the presence of a 100-fold excess of unlabelled antibody and was typically less than 5% of total cell-associated radiolabel. Experiments performed at 4°C were used to correct for surface radioactivity not removable by glycine-HCl buffer (usually <5%) and background radioactivity of the TCA-soluble fraction. Extracellular degradation at 37°C could be excluded since negligible radioactivity was found in the TCA-soluble fraction of T-cell line Jurkat, which does not express CD19, and in purified surface membranes from dis-

rupted Daudi or KM3 cells following incubation with ^{125}I -IgG or ^{125}I -F(ab')₂.

Internalization was defined as the total number of molecules transported across the cell membrane, and calculated by summation of TCA-soluble radioactivity to intracellular radioactivity.

Detection of FcγRIIa, FcγRIIb1 and FcγRIIb2 transcripts in the panel of B-cell lines

The presence of FcγRIIa, FcγRIIb1 and FcγRIIb2 transcripts in the panel of B-cell lines was assessed by RT-PCR as described before.³⁰ Briefly, total cellular RNA was extracted from 5×10^6 cells using the RNazol B isolation method³¹, concentrated by ethanol precipitation, resuspended in diethylpyrocarbonated water, and stored at -70°C. Total cellular RNA (5 μg) was reverse transcribed and the presence of FcγRIIa and FcγRIIb cDNA were amplified by PCR, using FcγRIIa- and FcγRIIb-specific primer pairs. PCR products were analyzed by electrophoresis on 1.8% (w/v) agarose gels stained with ethidium bromide, followed by Southern blotting to Zeta-probe nylon membranes (Bio-rad). To detect FcγRIIa and FcγRIIb PCR products, blots were hybridized overnight with FcγRIIa- and FcγRIIb-specific ^{32}P -labelled oligonucleotide probes, which were then washed, and analyzed using a 400A Phosphor-imager System (Molecular Dynamics, Sunnyvale, CA). cDNA of FcγRIIa, FcγRIIb1, and FcγRIIb2 transfectants served as controls.

Membrane expression of FcγRII

Surface expression levels of FcγRII (CD32) were determined via indirect immunofluorescence measured by FCM. Cells in phosphate-buffered saline (PBS) supplemented with 0.1% (w/v) bovine serum albumine and 0.1% sodium azide were incubated with MoAb IV.3 (2 μg/ml) for 45 min at 4°C and then washed and incubated with fluorescein isothiocyanate-labeled F(ab')₂ fragments of goat-anti-mouse IgG (Nutacon, American Qualex Division, Amsterdam, the Netherlands) for another 45 min at 4°C. Subsequently, cells were washed and the mean fluorescence intensity (MFI) of 10,000 cells was evaluated by FCM and expressed in arbitrary units.

EA-IgG rosetting

Rosette formation between B-cells and E sensitized with anti-glycophorin A switch-variant MoAb of IgG1-isotype³² was performed as described before.³⁰ Briefly, a 0.5% suspension of human E in PBS was incubated for 30 min at 37°C with the anti-glycophorin A MoAb and then washed and suspended in RPMI 1640-5% FCS. Fifty μl were then added to 50 μl B-cells (4×10^5), centrifugated (4 min at 40 x g), and incubated for 1 h at 4°C. Cells with at least three bound E were microscopically scored as EA-rosettes.

Statistical analysis

Statistical evaluation of results was performed using the Spearman correlation.

RESULTS

Cytotoxicity of CLB-B4-ricin A isotype-switch variants

The efficacy of the class switch variant ITs was determined by protein synthesis inhibition assayed with B-cell lines Daudi and KM3, which show high and intermediate membrane-expression levels of CD19, respectively. Both ITs demonstrated poor cytotoxicity (ID_{50} s of $\geq 10^{-8}$ M) in the absence of NH_4Cl . In the presence of NH_4Cl (6 mM) which enhances IT cytotoxicity³³, the IgG1-IT became about 100 times more potent than IgG2a-IT (ID_{50} s of $\sim 10^{-10}$ M, and 10^{-8} M, respectively). Neither IT, even in the presence of NH_4Cl , showed any cytotoxicity against T-cell line Jurkat, which lacks expression of CD19 (Figure 1).

In order to confirm that the cytotoxic effect of the CLB-B4-IT is mediated through the CD19 antigen, the blocking effect of free CD19 MoAb was examined. The addition of a 100-fold excess of CLB-B4-IgG2a completely blocked the cytotoxic action of the CLB-B4-IT against Daudi cells ($ID_{50} > 10^{-8}$ M in the presence of NH_4Cl for both the IgG1-IT and IgG2a-IT; data not shown). Moreover, non-binding control IT TS2.18-ricin A (CD2 and IgG1) and 35.1-ricin A (CD2 and IgG2a) demonstrated no cytotoxicity against Daudi cells (data not shown).

Scatchard analysis of the CLB-B4 IgGs and F(ab')₂ fragments

In order to determine whether the observed differences in cytotoxicity were attributable to differences in affinity, we performed binding studies with both MoAb (and their F(ab')₂-fragments) on Daudi cells. Both affinity ($\sim 8 \times 10^9 M^{-1}$), and number of cell-bound molecules ($\sim 67,000$) were found comparable for both MoAbs (and their F(ab')₂ fragments; Table 1).

Table 1 *Scatchard analysis of CLB-B4-IgG1 and CLB-B4-IgG2a binding to Daudi cells^a*

Antibody form	CLB-B4-IgG1		CLB-B4-IgG2a	
	K_a ($\times 10^{-9}$ M)	Molecules Bound/cell ($\times 10^{-4}$)	K_a ($\times 10^{-9}$ M)	Molecules Bound/cell ($\times 10^{-4}$)
IgG	7.1 ± 1.0	6.7 ± 1.2	8.3 ± 2.1	6.4 ± 0.8
F(ab') ₂	8.6 ± 1.4	6.9 ± 0.9	7.5 ± 1.8	6.9 ± 1.2

^aCells were incubated with various concentrations of ¹²⁵I-MoAb for 2 h at 4°C and then washed and assessed for membrane-bound radioactivity to calculate the affinity constant (K_a); the number of bound molecules/cell were counted under saturating conditions according to Scatchard (29). The mean \pm SD of three independent experiments is shown.




Figure 1 Cytotoxicity mediated by CLB-B4-ricin A-IgG1 and CLB-B4-ricin A-IgG2a

Daudi (A and B), KM3 (C and D), or Jurkat (E and F) cells were incubated with various concentrations of CLB-B4-ricin A-IgG1 (A, C, and E) or CLB-B4-ricin A-IgG2a (B, D, and F) in the absence (○, □) or presence (●, ■) of 6 mM NH₄Cl at 37°C. After 16 h, cells were washed and cultured with [³H]leucine for another 24 h. Protein synthesis is expressed as the percentage [³H]leucine incorporation relative to cells cultured in medium alone (control cells). Control Daudi and KM3 cells incorporated ~15,000 cpm, and Jurkat cells incorporated ~20,000 cpm. Background incorporation approximated 100 cpm. Assays were performed in triplicate (*n* = 3), and SD was <10%.

Binding and internalization kinetics of CLB-B4-IgG1 and CLB-B4-IgG2a

Because the efficacy of IT strongly depends on numbers of internalized molecules⁶, we studied binding and internalization kinetics of the switch variants. Following incubation




Figure 2 Cellular uptake and processing of CLB-B4-IgG1 and CLB-B4-IgG2a

Daudi (A and B) and KM3 (C and D) were incubated with ^{125}I -labeled CLB-B4-IgG1 (A and C) or CLB-B4-IgG2a (B and D) for various times at 37°C . Intracellularly degraded and subsequently exocytosed molecules (TCA-soluble radioactivity of supernatants; ▲) and intracellular molecules (cell associated radioactivity not-removable by glycine buffer; ●) were assessed as described in "Material and Methods". Internalized molecules were determined as the sum of intracellular and intracellularly degraded and subsequently exocytosed molecules (□). Assays were performed in triplicate ($n = 3$), and SD was $<10\%$. One of three concordant experiments is shown.

at 4°C with 10^{-8} M, equal amounts of IgG1 and IgG2a MoAbs were bound to the cell membranes (Daudi and KM3 cells, $\sim 40,000$ and $\sim 15,000$ molecules, respectively; data not shown). Incubation at 37°C , however, resulted in consistent internalization differences (Figure 2). Daudi cells internalized 3.5 times and KM3 cells 2.5 times more IgG1 MoAb than IgG2a MoAb (Daudi, 35,000 IgG1 molecules and 10,000 IgG2a molecules; KM3, 10,000 IgG1 molecules and 4,000 IgG2a molecules) during a 12-h incubation. The highest number of intracellularly present molecules was reached after 4 and 8 h for Daudi ($\sim 25,000$ IgG1 molecules and 7,500 IgG2a-molecules), and KM3 cells ($\sim 4,000$ IgG1 molecules and 3000 IgG2a molecules), respectively. Thereafter the number of intracellularly present molecules declined as a result of exocytosis (of intracellularly degraded MoAb).

Because the switch-variants differ only in subclass, we next tested the effect of removing the CD19-Fc fragment on internalization. Daudi cells were incubated with 10^{-8} M of




Figure 3 Effect of isotype on internalization

Daudi cells were incubated for 1 h at 37°C/4°C with ¹²⁵I-labeled IgG or F(ab')₂-fragments of CLB-B4-IgG1 and CLB-B4-IgG2a, or with isotype-matched MoAb TS2.18 (CD2 and IgG1) or 35.1 (CD2 and IgG2a). Data are presented as molecules/cell incubated at 37°C (corrected for cells treated at 4°C) as described in "Material and Methods". Experiments were performed in triplicate, and SD was <10%. One of three experiments yielding similar data is shown.

intact IgG or F(ab')₂ fragments of CLB-B4 for 1 h at 37°C; then the amount internalized molecules was determined (Figure 3). Both the IgG1- and the IgG2a-F(ab')₂-fragments were internalized to a comparable extent as intact IgG2a MoAb, indicating that the Fc part of the IgG1 MoAb was responsible for the observed increase in internalization. The binding of CD19 antigen appeared to be a prerequisite because no cellular uptake could




Figure 4 Effect of FcγR-blocking MoAb on IT-cytotoxicity

Daudi cells were incubated with various concentrations of the IgG1-IT (A) or IgG2a-IT (B) with NH₄Cl, in the absence (○) or presence (●) of FcγRI-blocking MoAb 197, or FcγRII-blocking MoAb AT10 (▲), or IV.3 (□) at 37°C. After 16 h, cells were washed and cultured with [³H]leucine for another 24 h. Protein synthesis is expressed as the percentage [³H]leucine incorporation relative to cells cultured in medium lacking IT. Untreated Daudi cells incorporated ~15,000 cpm. Background incorporation was 100 cpm. Assays were performed in triplicate, and SD was <10%. The experiment was repeated three times, yielding almost identical results.

be observed of isotype-matched irrelevant MoAb TS2.18 (CD2 and IgG1) or of 35.1 (CD2 and IgG2a).

Involvement of FcγRII in CLB-B4-ricin A cytotoxicity

Because the involvement of the IgG1-Fc fragment suggested an interaction with receptors for the Fc domain of IgG, we performed cytotoxicity assays with Daudi cells in the presence of FcγR-blocking MoAb. Addition of FcγRI-blocking MoAb 197 did not affect cytotoxicity of the IgG1-IT or IgG2a-IT. Coincubation of cells with FcγRII-blocking MoAb AT10 or IV.3 reduced the effectiveness of IgG1-IT without affecting IgG2a-IT cytotoxicity (Figure 4). Addition of AT10 F(ab')₂ fragments (2 μg/ml) resulted in a similar blocking to AT10 IgG (data not shown; *n* = 3), excluding the reduced cytotoxicity to be caused by the Fc part of this FcγRII-blocking MoAb. These results indicated that FcγRII is involved in the observed differences between the class switch ITs.

Table 2 *Relation between CLB-B4 internalization, FcγRIIa expression levels, and EA-rosette formation by B-cell lines and patient cells*

Cells	Internalization ^a (Molecules/cell)			FcγRIIa expression ^b (MFI)	EA-rosetting ^c (%)
	IgG1	IgG2a	IgG1/IgG2a ^d		
CRL	2,879	2,624	1.1	14.9	0
Daudi	9,540	3,085	3.1	18.8	96
DOHH2	11,245	11,063	1.0	1.1	0
Fravel	2,328	2,154	1.1	18.7	0
KM3	2,008	800	2.5	1.9	0
Nalm6	10,381	2,002	3.5	10.4	95
Raji	8,550	5,700	1.5	11.8	10
Ramos	11,251	11,941	1.0	0.4	0
REH	1,009	900	1.1	1.6	0
Ros1	4,532	4,498	1.0	22.0	0
RPM11788	1,259	1,290	1.0	5.1	0
Patient 1	1,019	917	1.1	6.3	0
Patient 2	3,192	3,002	1.1	2.1	2
Patient 3	2,201	1,928	1.1	1.0	0
Patient 4	2,675	1,574	1.7	17.8	20
Patient 5	3,172	1,233	2.6	3.5	60
Patient 6	4,841	4,586	1.1	15.0	0
Patient 7	1,775	1,714	1.0	1.0	0
Patient 8	515	498	1.0	8.5	0

^aCells were incubated for 1 h with 10⁻⁸ M ¹²⁵I-labeled MoAb at 37°C; then the number of internalized molecules/cell was determined as described in "Materials and Methods". The mean of three independent experiments is shown; SD < 10%. ^bFcγRII-expression levels were determined by indirect immunofluorescence. Cells were incubated with MoAb IV.3 (predominantly interacting with FcγRIIa) for 45 min at 4°C, washed and cultured for another 45 min with fluorescein isothiocyanate-labeled F(ab')₂ fraction of anti-mouse immunoglobulin antiserum. Data shown represent the MFI of 10,000 cells. ^cEA-rosette formation was used to estimate functional binding capacity of FcγRIIa. Cells were incubated with EA sensitized with IgG1 for 1 h at 4°C and washed and assessed for EA-rosettes. Data represent the percentage of cells with three or more bound EA-IgG. The mean of three experiments is shown; SD < 10%. ^dRatio of internalized IgG1 and IgG2a molecules upon incubation for 1h at 37°C.

Relation between internalization, FcγRII-isoform expression and EA-IgG rosette formation

To determine the FcγRII-isoform involved in the internalization of the IgG1 MoAb, we studied the relation between enhancement of IgG1 uptake and expression of the different isoforms. The enhancement of IgG1 uptake in 11 B-cell lines and malignant B-cells derived from 8 patients (determined after 1 h at 37°C) was expressed as the ratio of internalized IgG1 and IgG2a molecules (Table 2). As in some cases, the number of internalized molecules was rather low; nonspecific endocytosis due to a fluid phase uptake was examined with MoAbs TS2.18 (CD2 and IgG1) and 35.1 (CD2 and IgG2a). No cellular uptake of these control antibodies could be observed (data not shown). Besides Daudi, KM3, Nalm6 and Raji cells, malignant B-cells from two of the patients (patients nos. 4 and 5) showed an enhanced internalization of CD19-IgG1 molecules.




Figure 5 Expression of FcγRIIa, FcγRIIb1 and FcγRIIb2 transcripts in a panel of B-cell lines assessed by reverse transcription-PCR

RNA was isolated and reverse transcribed, and FcγRIIa and FcγRIIb transcripts were selectively amplified (described in "Material and Methods"). Products were run on a 1.8% agarose gel, blotted, and hybridized with isoform-specific oligonucleotide probes. FcγRIIa, FcγRIIb1, and FcγRIIb2 transfectants served as controls (Lanes 1-3). Blots were probed with either ³²P-labelled FcγRIIa (A) or FcγRIIb (B) specific oligonucleotides.

The expression of various FcγRII transcripts in the panel of B-cell lines was analyzed by performing an FcγRIIa- and FcγRIIb-specific PCR. Total cellular RNA was isolated and reverse transcribed to cDNA; specific primer pairs were used to amplify FcγRIIa, FcγRIIb1 and FcγRIIb2 transcripts (Figure 5). FcγRIIa was found to be present in all B-cell lines tested. The FcγRIIb1 and FcγRIIb2 expression levels varied between the different B-cell lines. Notably, FcγRIIa represented the sole FcγRII transcript present in Nalm6, KM3, DOHH2 and REH cells, suggesting that: (a) the FcγRIIa isoform is involved in enhancement of IgG1 uptake (Nalm6 and KM3 showing IgG1:IgG2a ratios of 3.5 and 2.5, respectively); and (b) FcγRIIa RNA expression as such is not indicative for enhanced IgG1 uptake (DOHH2 and REH, IgG1:IgG2a ratios of ~1).

Because the presence of RNA transcripts does not necessarily imply membrane expression of the receptor, we determined the membrane-expression levels of FcγRIIa using MoAb IV.3, which predominantly interacts with FcγRIIa (34; Table 2). Cell lines DOHH2, KM3, Ramos, REH, and B cells derived from patients 1,2,3,5, and 7 showed no or poor expression. All other cells showed intermediate to high expression. The relation between membrane expression levels and enhanced uptake of CD19-IgG1 is plotted in Figure 6. No correlation could be found ($r = 0.32$; $P < 0.185$; Fig. 6A).




Figure 6 Relation between enhancement of IgG1 uptake, FcγRII expression levels, and EA-rosette formation

To assay FcγRII expression levels (A), cells (as indicated in Table 2) were incubated with MoAb IV.3 for 45 min at 4°C, washed and cultured for another 45 min with fluorescein isothiocyanate-labeled anti-mouse serum. Data represent the mean fluorescence intensities (MFI). EA-rosette formation (B) was determined after incubation of cells with EA-IgG1 for 1 h at 37°C. Data represent the percentage of cells (indicated in Table 2) with three or more bound EA-IgG indicator cells. Assays were performed in duplicate. The experiment was repeated three times, yielding almost identical results.

The relation between expression levels and enhanced CD19-IgG1 uptake may be complicated by membrane-expression of “nonfunctional” FcγRIIa molecules. Therefore, we assessed functional IgG binding capacity of the FcγRIIa molecules by determining EA-rosette formation using IgG1-sensitized E (EA-IgG1; Table 2). A highly significant correlation between enhanced IgG1 uptake and EA-rosette formation was found ($r = 0.76$; $P < 0.001$; Fig. 6B). Remarkably, KM3 cells showed an enhanced uptake of IgG1 (ratio 2.5) without forming rosettes. This may have been due to the exceptionally low FcγRII-expression level on KM3 (MFI, 2.2). These results suggest that FcγRIIa molecules capable of IgG1 are involved in the enhancement of internalization observed upon interaction of cells with CD19-IgG1-IT.

DISCUSSION

Several reports addressed the use of CD19 MoAb-toxin conjugates as potent and specific reagents for neoplastic B-cell elimination. These include MoAb HD37 conjugated to ricin A¹⁰, to saporin³⁵, as well as MoAb B4 conjugated to blocked ricin³⁶ and MoAb B43 conjugated to pokeweed antiviral protein (PAP)³⁷, all of the mouse IgG1 isotype. In the present study, we evaluated the cytotoxic potency of CD19 class switch variants CLB-B4-IgG1 and CLB-B4-IgG2a upon conjugation to ricin A. The IgG1-IT required the presence of the cytotoxicity enhancer NH₄Cl to show similar cytotoxicity as reported for HD37-ricin A against Daudi cells (ID₅₀s of $\sim 10^{-10}$ M; Fig. 1)³⁸. This difference in potency is probably due to a variation in epitope specificity and/or binding affinity. More surprisingly, in view of both MoAbs sharing identical epitopes and binding characteristics, the CLB-B4-IgG2a-IT appeared to be ~ 100 -fold less cytotoxic than the CLB-B4-IgG1-IT when assayed on Daudi and KM3 cells (ID₅₀ of $\sim 10^{-8}$ M).

Internalization studies with ¹²⁵I-labelled MoAb and F(ab')₂ fragments revealed that the higher efficacy of the IgG1-IT is attributable to the IgG1 Fc fragment, enhancing the cellular uptake following membrane binding (Figs. 2 and 3). The enhancement of IgG1 uptake was observed in 4 of 11 B-cell lines and in B-cells of 2 of 8 patients examined, suggesting the involvement of a heterogeneously expressed receptor on B-cells. A likely candidate to be involved in the enhanced uptake of IgG1 may be FcγRII (CD32), which is both present on B-cells and capable of interacting with IgG1.^{30,39} By interaction with the Fc part of membrane-bound CD19-IgG1, FcγRII may be capable of supporting internalization of antigen-MoAb complexes, analogous to the mechanisms described by Kurlander.⁴⁰ This hypothesis was supported by the observation that incubation with FcγRII-blocking MoAb strongly reduced the cytotoxicity of the IgG1-IT (and not IgG2a-IT; Fig. 4). The increase in potency resulting in the interaction of the IgG1-Fc part with FcγRII (~ 100 -fold; Fig. 1) is tremendous, considering the relatively modest increase in internalization (~ 2.5 -fold; Fig. 2). Further studies are needed to address this disproportional increase in cytotoxicity, but it is tempting to speculate that the interaction with FcγRII alters the intracellular routing and/or handling of the antigen-IT complex in a way more favorable for ricin A to exert its cytotoxic action.

Expression of the FcγRII-isoform involved in the internalization of IgG1-IT in B cells was assayed by isoform-specific PCR experiments. Notably, FcγRIIa was the only isoform expressed by cell lines KM3 and Nalm6, showing enhanced IgG1-uptake. The strong correlation of enhancement of IgG1-uptake with EA-IgG1 rosetting, rather than with expression levels determined using MoAb IV.3 (predominantly interacting with FcγRIIa)³⁴, suggested that part of the expressed FcγRIIa molecules are incapable of IgG1-binding (and, thereby, incompetent to promote cellular uptake). This is in accordance with observations of Koenderman *et al*, who demonstrated that not all FcγRIIa molecules expressed by eosinophils are able to bind EA-IgG.²⁵ Remarkably, the cell line KM3 did

not form rosettes with EA-IgG1 but does show enhanced IgG1-uptake. In our view, this may be attributable to the exceptionally low FcγRIIa expression levels on KM3, which might be sufficient for enhancing IgG1 uptake in a catalytic manner (Table 2), but too low to support stable EA-IgG1 binding in EA-rosette experiments (Fig. 6B).

Several factors have been described to affect internalization and intracellular handling and, thereby, the cytotoxicity of ricin A ITs. Among these are the nature of the target cell and antigen^{7,8}, antigen density^{6,9}, binding affinity of the MoAb^{9,10}, and the epitope to which the IT is directed¹¹⁻¹³. Our observation of class switch variant ITs displaying different levels of cytotoxicities suggests also that the MoAb isotype is important for the efficacy of anti-B-cell ITs. Additional studies, however, are needed to address whether or not this phenomenon is restricted to CD19-ITs.

From our study, we conclude that the Fc part of ITs may be a parameter of importance for the efficacy of anti-B-cell ITs. Selection of the most effective isotype may improve the killing potency of an IT and, therefore, the success of these reagents in human therapy.

REFERENCES

1. Vitetta ES: Immunotoxins: new therapeutic reagents for autoimmunity, cancer, and AIDS. *J Clin Immunol* 10:155-185, 1990
2. Hertler AA, Frankel AE: Immunotoxins in the therapy of leukemias and lymphomas. *Cancer Invest* 9:211-9, 1991
3. Wawrzynczak EJ, Derbyshire EJ: Immunotoxins: the power and the glory. *Immunol Today* 13:381-3, 1992
4. Chignola R, Colombatti M, Dell'Arciprete L, Candiani C, Tridente G: Distribution of endocytosed molecules to intracellular acidic environments correlates with immunotoxin activity. *Int J Cancer* 46:1117-23, 1990
5. May RD, Wheeler HT, Finkelman FD, Uhr JW, Vitetta ES: Intracellular routing rather than cross-linking or rate of internalization determines the potency of immunotoxins directed against different epitopes of sIgD on murine B cells. *Cell Immunol* 135:490-500, 1991
6. Preijers FW, Tax WJ, De Witte T, Janssen A, v.d. Heijden H, Vidal H, Wessels JM, Capel PJ: Relationship between internalization and cytotoxicity of ricin A-chain immunotoxins. *Br J Haematol* 70:289-94, 1988
7. Press OW, Vitetta ES, Farr AG, Hansen JA, Martin PJ: Evaluation of ricin A-chain immunotoxins directed against human T cells. *Cell Immunol* 102:10-20, 1986
8. Embleton MJ, Byers VS, Lee HM, Scannon P, Blackhall NW, Baldwin RW: Sensitivity and selectivity of ricin toxin A chain-monoclonal antibody 791T/36 conjugates against human tumor cell lines. *Cancer Res* 46:5524-8, 1986
9. Youle RJ, Neville DM, Jr.: Kinetics of protein synthesis inactivation by ricin-anti-Thy 1.1 monoclonal antibody hybrids. Role of the ricin B subunit demonstrated by reconstitution. *J Biol Chem* 257:1598-601, 1982
10. May RD, Vitetta ES, Moldenhauer G, Dorken B: Selective killing of normal and neoplastic human B cells with anti-CD19- and anti-CD22-ricin A chain immunotoxins. *Cancer Drug Deliv* 3:261-72, 1986
11. Press OW, Martin PJ, Thorpe PE, Vitetta ES: Ricin A-chain containing immunotoxins directed against different epitopes on the CD2 molecule differ in their ability to kill normal and malignant T cells. *J Immunol* 141:4410-7, 1988
12. Luo Y, Seon BK: Marked difference in the in vivo antitumor efficacy between two immunotoxins targeted to different epitopes of common acute lymphoblastic leukemia antigen (CD10). Mechanisms involved in the differential activities of immunotoxins. *J Immunol* 145:1974-82, 1990
13. May RD, Finkelman FD, Wheeler HT, Uhr JW, Vitetta ES: Evaluation of ricin A chain-containing immunotoxins directed against different epitopes on the delta-chain of cell surface-associated IgD on murine B cells. *J Immunol* 144:3637-42, 1990
14. Derocq JM, Casellas P, Laurent G, Ravel S, Vidal H, Jansen F: Comparison of the cytotoxic potency of T101 Fab, F(ab')₂ and whole IgG immunotoxins. *J Immunol* 141:2837-43, 1988
15. Ghetie MA, Richardson J, Tucker T, Jones D, Uhr JW, Vitetta ES: Antitumor activity of Fab' and IgG-anti-CD22 immunotoxins in disseminated human B lymphoma grown in mice with severe combined immunodeficiency disease: effect on tumor cells in extranodal sites. *Cancer Res* 51:5876-80, 1991
16. Better M, Bernhard SL, Lei SP, Fishwild DM, Lane JA, Carroll SF, Horwitz AH: Potent anti-CD5 ricin A chain immunoconjugates from bacterially produced Fab' and F(ab')₂. *Proc Natl Acad Sci U S A* 90:457-61, 1993

17. de Rie MA, Zeijlemaker WP, von dem Borne AE: Inhibition, by vinca alkaloids and colchicine, of antigenic modulation induced by anti-CD19 monoclonal antibodies. *Leuk Res* 12:135-41, 1988
18. Van de Ouweland F, De Witte T, Geerdink P, Haanen C: Enrichment and cryopreservation of bone marrow progenitor cells for autologous reinfusion. *Cryobiology* 19:292-8, 1982
19. Sanchez-Madrid F, Krensky AM, Ware CF, Robbins E, Strominger JL, Burakoff SJ, Springer TA: Three distinct antigens associated with human T-lymphocyte-mediated cytotoxicity: LFA-1, LFA-2, and LFA-3. *Proc Natl Acad Sci U S A* 79:7489-93, 1982
20. Martin PJ, Longton G, Ledbetter JA, Newman W, Braun MP, Beatty PG, Hansen JA: Identification and functional characterization of two distinct epitopes on the human T cell surface protein Tp50. *J Immunol* 131:180-5, 1983
21. Guyre PM, Graziano RF, Vance BA, Morganelli PM, Fanger MW: Monoclonal antibodies that bind to distinct epitopes on Fc gamma RI are able to trigger receptor function. *J Immunol* 143:1650-5, 1989
22. Greenman J, Tutt AL, George AJ, Pulford KA, Stevenson GT, Glennie MJ: Characterization of a new monoclonal anti-Fc gamma RII antibody, AT10, and its incorporation into a bispecific F(ab')₂ derivative for recruitment of cytotoxic effectors. *Mol Immunol* 28:1243-54, 1991
23. Looney RJ, Abraham GN, Anderson CL: Human monocytes and U937 cells bear two distinct Fc receptors for IgG. *J Immunol* 136:1641-7, 1986
24. Parren PW, Warmerdam PA, Boeijs LC, Arts J, Westerdaal NA, Vlug A, Capel PJ, Aarden LA, van de Winkel JG: On the interaction of IgG subclasses with the low affinity Fc gamma RIIa (CD32) on human monocytes, neutrophils, and platelets. Analysis of a functional polymorphism to human IgG2. *J Clin Invest* 90:1337-46, 1992
25. Koenderman L, Hermans SW, Capel PJ, van de Winkel JG: Granulocyte-macrophage colony-stimulating factor induces sequential activation and deactivation of binding via a low-affinity IgG Fc receptor, hFc gamma RII, on human eosinophils. *Blood* 81:2413-9, 1993
26. Hunter WH: Radioimmunoassay. Oxford, Blackwell Scientific Publications, 1973
27. Ouchterlony O, Nilsson LA: Immunodiffusion and immunoelectrophoresis, vol. Section 32. Oxford, Blackwell Scientific Publications, 1986
28. van Oosterhout YV, Preijers FW, Wessels HM, de Witte T: Cytotoxicity of CD3-ricin A chain immunotoxins in relation to cellular uptake and degradation kinetics. *Cancer Res* 52:5921-5, 1992
29. Scatchard G: The attraction of proteins for small molecules and ions. *Ann NY Acad Sci* 51:660-672, 1949
30. Van Den Herik-Oudijk IE, Westerdaal NA, Henriquez NV, Capel PJ, Van De Winkel JG: Functional analysis of human Fc gamma RII (CD32) isoforms expressed in B lymphocytes. *J Immunol* 152:574-85, 1994
31. Kindt GC, van de Winkel JG, Moore SA, Anderson CL: Identification and structural characterization of Fc gamma-receptors on pulmonary alveolar macrophages. *Am J Physiol* 260:L403-11, 1991
32. Boot JH, Geerts ME, Aarden LA: Functional polymorphisms of Fc receptors in human monocyte-mediated cytotoxicity towards erythrocytes induced by murine isotype switch variants. *J Immunol* 142:1217-23, 1989
33. Casellas P, Bourrie BJ, Gros P, Jansen FK: Kinetics of cytotoxicity induced by immunotoxins. Enhancement by lysosomotropic amines and carboxylic ionophores. *J Biol Chem* 259:9359-64, 1984
34. Van de Winkel JGJ, Anderson CL: Cluster report: CD32. Oxford, Oxford University Press, 1994
35. Bregni M, Siena S, Formosa A, Lippi DA, Martineau D, Malavasi F, Dorken B, Bonadonna G, Gianni AM: B-cell restricted saporin immunotoxins: activity against B-cell lines and chronic lymphocytic leukemia cells. *Blood* 73:753-62, 1989

36. Lambert JM, Goldmacher VS, Collinson AR, Nadler LM, Blattler WA: An immunotoxin prepared with blocked ricin: a natural plant toxin adapted for therapeutic use. *Cancer Res* 51:6236-42, 1991
37. Uckun FM, Gajl-Peczalska KJ, Kersey JH, Houston LL, Vallera DA: Use of a novel colony assay to evaluate the cytotoxicity of an immunotoxin containing pokeweed antiviral protein against blast progenitor cells freshly obtained from patients with common B-lineage acute lymphoblastic leukemia. *J Exp Med* 163:347-68, 1986
38. Ghetie MA, May RD, Till M, Uhr JW, Ghetie V, Knowles PP, Relf M, Brown A, Wallace PM, Janossy G, et al.: Evaluation of ricin A chain-containing immunotoxins directed against CD19 and CD22 antigens on normal and malignant human B-cells as potential reagents for in vivo therapy. *Cancer Res* 48:2610-7, 1988
39. van de Winkel JG, Anderson CL: Biology of human immunoglobulin G Fc receptors. *J Leukoc Biol* 49:511-24, 1991
40. Kurlander RJ: Blockade of Fc receptor-mediated binding to U-937 cells by murine monoclonal antibodies directed against a variety of surface antigens. *J Immunol* 131:140-7, 1983

FLOW CYTOMETRIC EVALUATION OF IMMUNOTOXIN INDUCED CELL KILL

Ypke V.J.M. van Oosterhout


Peter J. van Horssen

Arie H.M. Pennings

J. Liesbeth van Ernst

Theo de Witte

Frank W.M.B. Preijers



Partly published in:

Prog Clin Biol Res 1994; 389: 89-95

and

Leukemia 1999; 13: 241-249

ABSTRACT

Immunotoxins (ITs) represent an attractive drug targeting concept with therapeutic potential for the treatment of a variety of malignancies and immunological disorders. Prior to their clinical evaluation, the reactivity of IT towards target and non-target cells has to be extensively evaluated. In this report, a method is described for the preclinical evaluation of IT-induced target cell killing. This method is based on the flow cytometric (FCM) identification of viable cells escaping IT treatment. The simultaneous use of two viability markers, propidium iodine (PI) and calceine (Calc), minimizes the occurrence of falsely positive/negative cells due to autofluorescence, or extensive DNA-disintegration preventing PI-incorporation. The admixture of a fixed concentration of inert beads prior to analysis enables the precise quantification of surviving cells. Using this method, it appeared that IT-induced cell kill is not immediately accompanied by the physical disruption of the cell membrane. An IT-free lag period of four days proves essential for full exposure of efficacy. With a 1:9 mixture of leukemic Ramos cells and normal bone marrow cells, it was demonstrated that the method can also be used for the evaluation of IT-based killing of non-clonogenic target cells in a heterogeneous cell sample. With its capacity of monitoring target cell kill in freshly derived heterogeneous patient samples, the flow cytometric assay forms a valuable addition to the conventional clonogenic assays and protein synthesis inhibition assays.

INTRODUCTION

Immunotoxins (ITs) are hybrid molecules consisting of a cell-specific compound, mostly monoclonal antibodies (MoAbs), conjugated to a potent bacterial or plant derived toxin, *e.g.* modified diphtheria toxin or the A chain of plant toxin ricin. Several studies demonstrated the potential of ITs to selectively kill target cells in a highly efficient manner.¹⁻³ Moreover, the past two decades, ITs have been tested both in animals and

human studies for the treatment of life-threatening diseases, including various cancers and immunological disorders.⁴⁻¹¹

Prior to its clinical use, ITs have to be extensively screened for reactivity against target and non-target cells. Their potency is regularly expressed as the dose resulting in 50% inhibition of protein synthesis (ID_{50}). In general, ITs having an ID_{50} of 10^{-10} M or less are considered suitable for clinical use, supposed they do not cross-react with life-sustaining tissues.¹² It should be realized, though, that the presence of a small fraction of IT-resistant target cells might give rise to recurrence of the disease without having much impact on the ID_{50} . Therefore, in designing the optimal therapy, it is also important to determine the number of cells escaping various treatment regimens. So far, clonogenic assays have proven to be an useful tool for an accurate quantification of cells resistant to IT-treatment, especially when using cell lines as target.¹³⁻¹⁵ Unfortunately, the in vitro clonogenicity of patient derived target-cells is often too low to perform a reliable clonogenic assay. Because of this limitation, a quantitative flowcytometric (FCM) method was developed capable of determining IT-induced cell kill in freshly derived patient-samples as well. This study determines the prerequisites for a reliable quantification of surviving target-cells in a heterogeneous population such as human bone marrow.

MATERIALS AND METHODS

Malignant B-cell lines and bone marrow cells

Human B-cell lines Daudi, KM3 and Ramos were obtained from the American Type Culture Collection (ATTC, Rockville, MD). Normal bone marrow cells were obtained from patients undergoing cardiac surgery, after informed consent. Erythrocytes and mature granulocytes were removed by Ficoll (1.077 g/ml) density gradient centrifugation. Mononuclear cells were suspended in culture medium and were either used directly, or cryopreserved (10% dimethyl sulfoxide) without loss of functional potential.¹⁶

Cells were cultured in medium consisting of RPMI 1640 (Flow, Irvine, Scotland) supplemented with 10% heat-inactivated FCS (Gibco, Paisley, Scotland), glutamine (2mM), penicillin (100 units/ml) and streptomycin (100 µg/ml) in a humidified incubator with 5% CO₂ in air at 37°C.

Immunotoxins (IT)

MoAb CLB-B4 (anti-CD19, IgG1) and CLB-B-ly/1 (anti-CD22, IgG1) were obtained from the Central Laboratory of the Red Cross Blood Bank (Amsterdam, the Netherlands). Recombinant ricin A was generously provided by Zeneca Pharmaceuticals (Macclesfield, UK). ITs were constructed by biochemical linkage of MoAb and ricin A using *N*-succinimidyld-3-(2-pyridyldithio)propionate (SPDP) (Pierce, Rockford, IL), as previously described.¹³ Following conjugation, preservation of MoAb-binding activity was assessed

by FCM (Coulter Epics Elite, Hialeah, FL). Retention of ricin A activity was determined in a rabbit reticulocyte lysate assay, as described by Press *et al.*¹⁷

IT-treatment

Cells in culture medium at a concentration of 10^6 cells/ml were incubated with IT in triplicate at 37°C. IT concentrations were expressed in molar, ranging from 10^{-13} - 10^{-8} M. In some experiments, cytotoxicity enhancers NH_4Cl (6 mM) or nigericin (10 nM) were added to the culture medium to enhance the cytotoxicity of the ricin A-IT. After 16 h of IT-treatment, cells were washed and assayed for the number of surviving cells by either flow cytometry or a clonogenic assays.

Flow cytometric evaluation of surviving cells

Following IT-treatment, cells were cultured for varying days in IT-free culture medium at 37°C. After this lag period, cells were incubated with 2 $\mu\text{g/ml}$ propidium iodine (PI) (Molecular Probes, Junction City, OR) for 30 minutes at room temperature to stain death cells. In later experiments, also 2 $\mu\text{g/ml}$ calcein AM (Calc) (Molecular Probes) was added as an additional viability marker. Samples were then analyzed in triplicate on a flow cytometer (Coulter Epics Elite, Hialeah, FL) using a minimum of 10,000 cells. Viable cells were identified as being PI-negative and, if applicable, Calc-positive. Overlap of emission spectra of PI and Calc was adjusted by electronic compensation using single-labeled samples. Prior to FCM analysis, a fixed amount of inert beads (DNA-check or Flow-Count fluorospheres, Coulter) was added to each sample (*e.g.* 5×10^4 beads/ml) to enable the quantification of surviving cells. To prevent sedimentation of the beads, the FCM analysis was performed under continuous gently shaking of the sample.

Clonogenic assays

The clonogenic capacity of IT-treated cells was determined as described earlier.^{13,14} Briefly, IT-treated cells were resuspended in cloning medium consisting of RPMI 1640 supplemented with 1 mM α -ketoglutarate (Sigma, St. Louis, MO), 1 mM sodium oxaloacetate (Sigma), 5% fetal calf serum, and 10% heat inactivated horse serum (Hyclone, Logan, UT). Varying numbers of cells (10^2 - 10^6) were seeded in double layer agarose (3%) and cultured for 15 days at 37°C. Absolute numbers of surviving cells were calculated from the number of colonies, the number of plated cells and the cloning frequency of untreated cells (12% for Daudi and 28% for KM3).

Detection of IT-efficacy in a heterogeneous sample

Ramos cells were stained with the green-fluorescent membrane intercalating dye D-275 (Molecular probes, Eugene, OR), according to the manufacturer's protocol. Labeled Ramos cells were mixed in a 1:9 ratio with normal bone marrow and subsequently treated with IT and analyzed by FCM as described above. Overlap of emission spectra of PI and D-275 could be adjusted by electronic compensation using single-labeled samples.

RESULTS

Kinetics of IT-Induced Cell Kill as Determined by FCM

PI incorporates in the DNA of dead cells after passing through their compromised membranes. For a reliable assessment of IT-induced cell kill it is essential to determine the ‘lag-period’ between the irreversible inhibition of protein synthesis and the resulting physical disruption of the cell membranes.




Figure 1
PI Uptake of Daudi treated cells with anti-CD19 IT (CLB-B4-ricin A)

Cells were treated with anti-CD19 IT for 24 h and washed. After an IT-free incubation of various days, PI was added and cells were evaluated by FCM for viability. Cells treated accordingly without IT served as control. The percentage PI-positive cells is indicated. One of three concordant experiments in shown.

Figure 1 illustrates the fate of Daudi cells treated with CLB-B4-ricin A (anti-CD19) in the presence of NH_4Cl as enhancer of cytotoxicity. Prolongation of the post-treatment (IT-free) incubation period up to four days resulted in a progressive increase in the number of dead or dying cells, characterized by a strongly reduced forward scatter. This

change in light-scatter signals was accompanied by a disturbance of the physical integrity of the cell membranes, as reflected by the resulting PI⁺ phenotype. The maximal number of dead cells was observed four days after treatment with CLB-B4-ricin A (95% PI⁺), and did not increase upon further prolongation of the post-treatment period. The control sample treated with unconjugated MoAb CLB-B4 and ricin A, contained about 10% dead cells following the same incubation period. Similar patterns were observed with different combinations of ITs and cell lines (data not shown, n = 5).




Figure 2
Quantification
of IT-induced cell kill

Ramos cells were incubated with a suboptimal concentration anti-CD22 IT (CLB-B-Ly/1-ricin A, 5×10^{-11} M), in the presence (C) or absence (B) of nigericin (10 nM). Cells incubated with culture medium served as control (A). After 24 h, cells were washed and incubated without IT for an additional 4 days. Subsequently, cells were stained with PI and Calc and analyzed by FCM for viability (PI/Calc⁺) in the presence of a fixed concentration of inert beads (5×10^4 /ml). Analysis was stopped when 1,000 beads had passed the flow cell. One of three concordant experiments is shown.

Quantification of surviving cells

Treatment with IT may result in the elimination of several logs of antigen-bearing target cells, especially in the presence of cytotoxicity enhancers as NH_4Cl , monensin or nigericin.^{13,14,18} For an accurate FCM-based quantification of surviving cells, an internal standard in the form of inert beads was added to the IT-treated samples prior to FCM-

analysis. The amount of beads that passes the flow cell during analysis, enables the precise calculation of the number of viable cells. Moreover, to decrease the number of false positive/negative cells (*e.g.* due to auto-fluorescence or the extensive disintegration of DNA preventing PI-incorporation) Calc was added as a second viability marker. Only cells being PI/Calc⁺ were considered viable cells.

Figure 2 shows a typical FCM-analysis of Ramos cells 4 days after treatment with a sub-optimal concentration CLB-B-ly/1-ricin A (5×10^{-11} M) in the presence or absence of cytotoxicity enhancer nigericin. The inert beads ($10^5/\text{ml}$) could easily be distinguished from viable as well as non-viable cells based on their relatively small forward scatter and large right angle scatter. As an alternative, beads could be identified by their strong broad-band fluorescence (data not shown). FCM-analyses were stopped when 1,000 beads had passed the flow cell. Comparison of the viable fractions of the control and IT-treated samples revealed a 90-fold increase of efficacy upon addition of nigericin (~ 0.24 and 2.2 logs of kill with and without nigericin, respectively). Nigericin alone did only marginally effect the viability ($<7\%$ reduction of viable cells).

Comparison of FCM-evaluation and clonogenic assays

Conventional clonogenic assays have proven to be a valuable tool for estimation of cell kill. Table 1 presents the reduction in viable cells after incubation different combinations of target-cells and ITs, as determined with FCM-evaluation and clonogenic assays. Both assays revealed comparable results, demonstrating the superiority of the anti-CD22 IT as compared to its anti-CD19 counterpart. Unconjugated MoAbs and ricin A did not influence the viability of any of the target-cells in neither assay (data not shown, $n = 3$).

Table 1 *Comparison of FCM-evaluation and clonogenic assays.*

	Depletion of target cells (logs)			
	CD19-ricin A		CD22-ricin A	
	clonogenic assay	flow cytometry	clonogenic assay	flow cytometry
Daudi	1.3	1.5	3.0	3.3
KM3	1.3	1.1	2.0	2.7

SDs were less than 10% ($n = 3$)

Monitoring of cell kill in a heterogeneous sample

Fluorescent labeled Ramos cells were mixed in a 1 to 9 ratio with BM in order to mimic residual leukemic blasts. This mixture was treated with CLB-B-ly/1-ricin A in the presence of NH_4Cl , and assayed for dead and viable cells by FCM-evaluation Figure 3. Their green fluorescence made Ramos cells easily distinguishable from BM cells, including CD19-positive lymphocytes derived from blood admixture. Comparison of the viable fractions of the IT-treated sample and control, demonstrated an approximately 3 logs

of depletion of fluorescent Ramos cells. Omission of cytotoxicity enhancer NH_4Cl reduced the depletion-efficacy to approximately 1 logs ($n = 3$, data not shown).




Figure 3 Monitoring of IT-induced cell kill in bone marrow

Green fluorescent (D-275) labeled Ramos cells were mixed in a 1:9 ratio with BM cells. The mixture was treated with or without 10^{-8} M anti-CD22 IT (CLB-B-Ly/1-ricin A) in the presence of NH_4Cl (6 mM) at 37°C . Following 24 h, cells were washed and incubated in culture medium for four additional days. Subsequently, cells were stained with PI and evaluated for viability by FCM. Before analysis, a fixed concentration of inert beads ($10^5/\text{ml}$) was added to the samples. Analysis was stopped after 1,000 beads had passed the flow cell. One of three concordant experiments is shown.

DISCUSSION

Viability staining in combination with flow cytometric analysis has been used for diverse applications like monitoring of lymphocyte mediated target-cell lysis and toxicity testing of chemicals.¹⁹⁻²⁴ The use of this method for evaluation of IT-induced cell kill appeared to be somewhat more complicated. Reason for this is that the toxin-based inhibition of protein synthesis is not accompanied immediately by physical disruption of the cell

membrane. An IT-free incubation period of four days appeared to be necessary for full exposure of IT-toxicity. Moreover, the high potency of ITs (killing up to several logs of target cells) requires a sensitive and precise method for the quantification of residual cells. Simply counting the number of flow cytometric events seems not very adequate, as this parameter is influenced by dead cell particles and fluctuating sampling rates. Therefore, inspired by others, we added a fixed amount of inert beads to the IT-treated sample prior to FCM-analysis.²⁵ This enabled an accurate quantification of viable cells, which closely correlated with the outcome of conventional clonogenic assays. This is somewhat in contrast with the results obtained by Laurant *et al.*²⁵ Already in 1986, they described the use of viability staining to evaluate the sensitivity of fresh leukemic cells to ITs. Their assay appeared to be indicative rather than quantitative, though, only roughly qualifying leukemic cells as modestly, intermediate or highly susceptible to IT-treatment. This may be explained by the moment of analysis, directly following IT-treatment, which is before maximal exposure of efficacy.

When evaluating target cell kill in a heterogeneous population as blood or BM, the death/alive analysis needs to be combined with the identification of target-cells (Fig. 3). The use of fluorescent cell specific MoAbs is most convenient, enabling both the viability staining and target-cell identification in one analysis. However, this requires the presence of one or more 'identification-antigens' besides the 'target-antigen' already used for therapy. The latter is inappropriate as it might be modulated or covered by the therapeutic mAb. Moreover, residual malignant cells may be enriched for cells with low expression of target-antigen. As an alternative, target cells may be identified by the PCR-amplification of a cell specific DNA-sequence.²⁶⁻²⁸ In that case, viable cells should first be presorted to prevent DNA from IT-eliminated cells contributing to the PCR signal. Based on the specifications of current flow cytometers, a triple-sorted fraction should contain less than one in a million erroneously sorted non-viable cells, equalizing the detection limit of sensitive PCR-assay.

Protein synthesis inhibition is widely used to express the potency of an IT. Estimation of the ID_{50} is relatively easy, fast and independent of *in vitro* clonogenic capacity. Its major disadvantage, though, is that the ID_{50} is hardly influenced by the presence of a minor population of therapy-resistant (potentially tumorigenic) target-cells. The potential relevance of this is demonstrated by observations of Flavel *et al.*, who demonstrated that the combined use of an anti-CD19 and anti-CD38 IT did not reduce the ID_{50} towards Burkitt's lymphoma cell line Ramos, as compared to either IT alone.²⁹ The survival of Ramos-xenografted mice, in contrast, dramatically increased upon simultaneous administration of both ITs. Apparently, the IT-combination eliminated target-cells with low antigen density which otherwise would escape single IT-treatment. The flow cytometric method is more likely to identify such relatively small cell population, which appears to be of no significance in terms of overall protein synthesis.

From our study, we conclude that the FCM-evaluation of IT-induced cell kill forms a useful addition to the conventional clonogenic assays and protein synthesis inhibition assay. Several studies confirmed that this assay can be used for the monitoring of IT-induced target-cell elimination in freshly derived heterogeneous patient samples.^{10,18} As such, this assay may prove a valuable tool to help improve this form of therapy.

REFERENCES

1. Vitetta ES, Thorpe PE, Uhr JW: Immunotoxins: magic bullets or misguided missiles? *Immunol Today* 14:252-9, 1993
2. Vallera DA: Immunotoxins: will their clinical promise be fulfilled? *Blood* 83:309-17, 1994
3. Kreitman RJ: Immunotoxins in cancer therapy. *Curr Opin Immunol* 11:570-8, 1999
4. Preijers FW, De Witte T, Wessels JM, De Gast GC, Van Leeuwen E, Capel PJ, Haanen C: Autologous transplantation of bone marrow purged in vitro with anti-CD7- (WT1-) ricin A immunotoxin in T-cell lymphoblastic leukemia and lymphoma. *Blood* 74:1152-8, 1989
5. Byers VS, Henslee PJ, Kernan NA, Blazar BR, Gingrich R, Phillips GL, LeMaistre CF, Gilliland G, Antin JH, Martin P, *et al.*: Use of an anti-pan T-lymphocyte ricin A chain immunotoxin in steroid-resistant acute graft-versus-host disease. *Blood* 75:1426-32, 1990
6. Vitetta ES, Stone M, Amlot P, Fay J, May R, Till M, Newman J, Clark P, Collins R, Cunningham D, Ghetie V, Uhr JW, Thorpe PE: Phase I immunotoxin trial in patients with B-cell lymphoma. *Cancer Res* 51:4052-8, 1991
7. Engert A, Diehl V, Schnell R, Radszuhn A, Hatwig MT, Drillich S, Schon G, Bohlen H, Tesch H, Hansmann ML, Barth S, Schindler J, Ghetie V, Uhr J, Vitetta E: A phase-I study of an anti-CD25 ricin A-chain immunotoxin (RFT5-SMPT-dgA) in patients with refractory Hodgkin's lymphoma. *Blood* 89:403-10, 1997
8. Senderowicz AM, Vitetta E, Headlee D, Ghetie V, Uhr JW, Figg WD, Lush RM, Stetler-Stevenson M, Kershaw G, Kingma DW, Jaffe ES, Sausville EA: Complete sustained response of a refractory, post-transplantation, large B-cell lymphoma to an anti-CD22 immunotoxin. *Ann Intern Med* 126:882-5, 1997
9. Lynch TJ, Jr., Lambert JM, Coral F, Shefner J, Wen P, Blattler WA, Collinson AR, Ariniello PD, Brame G, Cook S, Esseltine D, Elias A, Skarin A, Ritz J: Immunotoxin therapy of small-cell lung cancer: a phase I study of N901- blocked ricin. *J Clin Oncol* 15:723-34, 1997
10. van Oosterhout YV, van Emst L, Schattenberg AV, Tax WJ, Ruiter DJ, Spits H, Nagengast FM, Masereeuw R, Evers S, de Witte T, Preijers FW: A combination of anti-CD3 and anti-CD7 ricin A-immunotoxins for the in vivo treatment of acute graft-versus-host disease. *Blood* 95:3693-3701, 2000
11. Kreitman RJ, Wilson WH, White JD, Stetler-Stevenson M, Jaffe ES, Giardina S, Waldmann TA, Pastan I: Phase I trial of recombinant immunotoxin anti-Tac(Fv)-PE38 (LMB-2) in patients with hematologic malignancies. *J Clin Oncol* 18:1622-36, 2000
12. Ghetie V, Vitetta E: Immunotoxins in the therapy of cancer: from bench to clinic. *Pharmacol Ther* 63:209-34, 1994
13. Preijers FW, De Witte T, Wessels JM, Meyerink JP, Haanen C, Capel PJ: Cytotoxic potential of anti-CD7 immunotoxin (WT1-ricin A) to purge ex vivo malignant T cells in bone marrow. *Br J Haematol* 71:195-201, 1989
14. Casellas P, Canat X, Fauser AA, Gros O, Laurent G, Poncelet P, Jansen FK: Optimal elimination of leukemic T cells from human bone marrow with T101-ricin A-chain immunotoxin. *Blood* 65:289-97, 1985
15. Myers DE, Uckun FM, Ball ED, Vallera DA: Immunotoxins for ex vivo marrow purging in autologous bone marrow transplantation for acute nonlymphocytic leukemia. *Transplantation* 46:240-5, 1988
16. Van de Ouweland F, De Witte T, Geerdink P, Haanen C: Enrichment and cryopreservation of bone marrow progenitor cells for autologous reinfusion. *Cryobiology* 19:292-8, 1982
17. Press OW, Vitetta ES, Martin PJ: A simplified microassay for inhibition of protein synthesis in reticulocyte lysates by immunotoxins. *Immunol Lett* 14:37-41, 1986

18. van Horssen PJ, van Oosterhout YV, Evers S, Backus HH, van Oijen MG, Bongaerts R, de Witte T, Preijers FW: Influence of cytotoxicity enhancers in combination with human serum on the activity of CD22-recombinant ricin A against B cell lines, chronic and acute lymphocytic leukemia cells. *Leukemia* 13:241-9, 1999
19. Mastbergen SC, Duivenvoorden I, Versteegh RT, Geldof AA: Cell cycle arrest and clonogenic tumor cell kill by divergent chemotherapeutic drugs. *Anticancer Res* 20:1833-8, 2000
20. Ackermann MF, Lamm KR, Wiegand GW, Luster MI: Antitumor activity of murine neutrophils demonstrated by cytometric analysis. *Cancer Res* 49:528-32, 1989
21. Shenker BJ, Datar S, Mansfield K, Shapiro IM: Induction of apoptosis in human T-cells by organomercuric compounds: a flow cytometric analysis. *Toxicol Appl Pharmacol* 143:397-406, 1997
22. Sumitran S, Anderson P, Widner H, Holgersson J: Porcine embryonic brain cell cytotoxicity mediated by human natural killer cells. *Cell Transplant* 8:601-10, 1999
23. Papadopoulos NG, Dedoussis GV, Spanakos G, Gritzapis AD, Baxevas CN, Papamichail M: An improved fluorescence assay for the determination of lymphocyte-mediated cytotoxicity using flow cytometry. *J Immunol Methods* 177:101-11, 1994
24. Hatam L, Schuval S, Bonagura VR: Flow cytometric analysis of natural killer cell function as a clinical assay. *Cytometry* 16:59-68, 1994
25. Laurent G, Kuhlein E, Casellas P, Canat X, Carayon P, Poncelet P, Correll S, Rigal F, Jansen FK: Determination of sensitivity of fresh leukemia cells to immunotoxins. *Cancer Res* 46:2289-94, 1986
26. Gribben JG, Freedman AS, Neuberg D, Roy DC, Blake KW, Woo SD, Grossbard ML, Rabinowe SN, Coral F, Freeman GJ, *et al.*: Immunologic purging of marrow assessed by PCR before autologous bone marrow transplantation for B-cell lymphoma. *N Engl J Med* 325:1525-33, 1991
27. Meijerink JP, Smetsers TF, Raemaekers JM, Bogman MJ, De Witte T, Mensink EJ: Quantitation of follicular non-Hodgkin's lymphoma cells carrying t(14;18) by competitive polymerase chain reaction. *Br J Haematol* 84:250-6, 1993
28. Willems P, Verhagen O, Segeren C, Veenhuizen P, Guikema J, Wiemer E, Groothuis L, Jong TB, Kok H, Bloem A, Bos N, Vellenga E, Mensink E, Sonneveld P, van Der Schoot HL, Group RR: Consensus strategy to quantitate malignant cells in myeloma patients is validated in a multicenter study. Belgium-Dutch Hematology-Oncology Group. *Blood* 96:637-0, 2000
29. Flavell DJ, Boehm DA, Emery L, Noss A, Ramsay A, Flavell SU: Therapy of human B-cell lymphoma bearing SCID mice is more effective with anti-CD19- and anti-CD38-saporin immunotoxins used in combination than with either immunotoxin used alone. *Int J Cancer* 62:337-44, 1995

**LABORATORY PRODUCTION OF ANTI-CD₃ AND
ANTI-CD₇ RICIN A-IMMUNOTOXINS
FOR A CLINICAL PILOT STUDY**

Ypke V.J.M. van Oosterhout

J. Liesbeth van Emst

Hans H. Bakker

Frank W.M.B. Preijers


Anton V.M.B. Schattenberg

Dirk J. Ruiter

Sabine Evers

Joop P. Koopman

Theo de Witte



Int J Pharm, in press

ABSTRACT

This report describes the preparation of an immunotoxin-combination, consisting of an anti-CD3 and anti-CD7 monoclonal antibody (MoAb) both conjugated to the A-chain of plant toxin ricin, for the experimental treatment of graft-versus-host disease. MoAbs and toxin were conjugated by conventional biochemical and chromatographic techniques. Raw materials, intermediate and final products were evaluated conform the relevant 'points to consider' of the FDA. Yields, purity and sterility of the two final products were all satisfactory. Preservation of MoAb-affinity and toxin-activity were confirmed in biological assays. The LD₅₀, 25-45 mg immunotoxin-combination/kg mouse, equaled that of similar immunotoxins already in clinical use. Because *in vitro* cross-reactivity screening revealed an unexpected binding of the CD3-MoAb to the esophagus epithelium, human doses immunotoxin-combination were administered to two cynomolgus monkeys. Clinically relevant serum concentrations were obtained without irreversible toxicities occurring. The T_{1/2} varied between ~6-9 hours, the C_{max} ranged from 1.8-3.9 µg/ml. The main side effect was a transient rise of serum creatine kinase. Importantly, neither damage nor binding of the CD3-immunotoxin to the monkey esophagus epithelium could be demonstrated. It was concluded that sufficient material of proper quality and with an acceptable toxicity profile was produced, warranting the evaluation in a clinical pilot-study.

INTRODUCTION

Monoclonal antibodies (MoAbs) conjugated to bacterial or plant-derived toxins (*e.g.* pseudomonas exotoxin or the deglycosylated A-chain of plant toxin ricin), so called immunotoxins, form attractive agents for the selective elimination of disease causing cells (reviewed in 1-3). Within the last two decades, various immunotoxins have been tested in clinical studies for the treatment of life-threatening disorders, including solid

and diffuse tumors and immunological disorders.⁴⁻⁹ Despite differences in specificity and efficacy, their mechanism of action is basically comparable. The MoAb moiety binds specifically to an antigen expressed on the surface of target cells, whereafter the entire antigen-immunotoxin complex is internalized. Once inside the cell, the toxin-subunit is translocated to the cytosol where it inhibits protein synthesis irreversibly by means of a catalytic reaction culminating in the cell's death. The potency of a given immunotoxin depends in essence on the number of immunotoxin-molecules internalized, and on the intracellular handling following internalization.¹⁰⁻¹⁴ Recently, we demonstrated in laboratory experiments that a combination of two deglycosylated ricin A-chain (dgA) based immunotoxins, directed against lymphocyte differentiation antigens CD3 and CD7, acts synergistically in killing activated T cells.⁹ This makes this particular immunotoxin-combination potentially valuable to control certain immunological disorders, like transplantation related rejection and several auto-immune diseases. To explore its therapeutic potential, a clinical pilot study is set up for the treatment of patients with severe graft-versus-host disease, a serious complication of hematological stem cell transplantation initiated by donor-derived T cells.⁹ This document describes the semi-large scale laboratory production and quality control of the clinical grade immunotoxin-combination used for the pilot-study.

MATERIALS AND METHODS

Processes, facilities and documentation

The manufacturing processes and the production facilities were designed to minimize the risk of contamination of raw materials, intermediates and the two final products. Prior to the production, the formulae, manufacturing method and specifications of all (intermediate) products were established and documented. The documentation included instructions for operation of equipment, the manufacturing, packaging and storage.

MoAb and dgA

Hybridoma cells producing murine MoAb SPV-T3a (α CD3, IgG2b κ)¹⁵ and WT1 (α CD7, IgG2a λ)¹⁶ were kindly provided by Drs. H. Spits (NKI, Amsterdam, The Netherlands) and W. Tax (UMC St Radboud, Nijmegen, The Netherlands), respectively. From both hybridomas, a master cell bank (MCB) and manufacturing working cell bank (MWCB) were prepared. The production of MoAb WT1 was performed at the institutional Central Hematology Laboratory. The production of MoAb SPV-T3a, as well as the protein A purification of both MoAb, was performed by IQ Corporation (Groningen, The Netherlands).

For antibody production, hybridoma WT1 was grown in 162 cm² culture flasks in RPMI1640 medium (Flow, Irvine, Scotland) supplemented with 8% fetal calf serum

(FCS) (Integro, Zaandam, The Netherlands), glutamine (2mM), penicillin (100 U/ml) and streptomycin (100 µg/ml) in a humidified incubator with 5% CO₂ in air at 37°C. Hybridoma SPV-T3a was cultured under the same conditions in RPMI1640 supplemented with 8% FCS, D(+)-glucose (2.5 g/l), sodium hydrogencarbonate (2.0 g/l), L-glutamine (0.4 g/l), gentamycine (0.04 g/l), and HybridCult supplement (IQ Corporation). Cells were first expanded in log-phase and then cultured for another two weeks without medium changes, resulting in a final viability of about 10%. Subsequently, cell cultures were centrifuged, and the supernatants were filter sterilized (crude harvest) and stored at -20°C (SPV-T3a) or 4°C (WT1). For MoAb purification, the crude harvest was 1:1 diluted with 'bindings-buffer', containing 1.5 M glycine and 3.0 M NaCl (pH 8.9), and applied to a protein A column. Affinity bound MoAbs were eluted with citrate buffer (0.1 M citrate, pH 2.8-3.0) and directly neutralized with 0.1 volume 1M Tris-HCl. Subsequently, purified MoAbs were dialyzed against 0.01 M phosphate buffered saline (pH 7.4) (PBS), filter sterilized and stored at a concentration of approximately 4 mg/ml at 4°C.

Clinical grade deglycosylated ricin A (dgA) was purchased from Inland Laboratories (Austin, TX). The dgA contained the two naturally occurring A-chain forms, of 30 and 32 kDa, found in the seeds of *Ricinus communis*.^{17,18} The dgA was stored at a concentration of 3.06 mg/ml in phosphate buffered saline (pH 7.2) containing 50% glycerol at -20°C.

MoAb-dgA conjugation

Preparation of MoAb-dgA conjugates was based on the method described by Ghetie *et al.*¹⁹ Sterile and endotoxin-free solutions were prepared by the Department of Clinical Pharmacy. The preparative chromatographic columns were set up as a closed system except for the inlet and outlet which were positioned in a class 100 laminar flow cabinet. Chromatographic media and columns (all Amersham Pharmacia Biotech, Uppsala, Sweden) were sanitized for every conjugation according to the supplier's 'product support files'. The conjugation procedure started with the derivatization of MoAb with an 8 to 12-fold molar excess of crosslinker 4-succinimidylloxycarbonyl- α -methyl- α -(2-pyridyldithio)toluene (SMPT)¹⁸ (Pierce, Rockford, IL), and the reduction of dgA with 5 mM DTT. Both reactions were carried out at 37°C for 1 hour. Excess SMPT and DTT were subsequently removed by gel filtration on Sephadex G-25 (XK100/50 column) equilibrated with PBS containing 0.05% Tween-20 (PBST). Average MoAb-MPT loading was determined spectrophotometrically, as described elsewhere.²⁰ For the actual conjugation reaction, MoAb-MPT was then mixed with a 2.5-fold molar excess of freshly reduced dgA-SH. Subsequently, 0.2 volume 85% glycerol was added and the conjugation mixture was gently stirred and the reaction was allowed to proceed at 20-25°C for 48 hours. The resulting crude conjugate was purified by affinity chromatography on Blue-Sepharose (XK50/60 column) for removal of unconjugated MoAb (The Sepharose-conjugated Cybacron Blue has a high affinity for dgA). In case of WT1-dgA,

the Blue-Sepharose was equilibrated with PBST. For SPV-T3a-dgA, the Na⁺ concentration of the reaction mixture and loading buffer had to be adjusted to 300 mM in order to prevent affinity-binding of unconjugated SPV-T3a. Affinity-bound dgA and MoAb-dgA conjugates were eluted with PBST containing 1M NaCl. High molecular weight material and free dgA were subsequently removed by size exclusion chromatography on Sephacryl S-300 HR (K100/100 column) equilibrated with PBST (formulation buffer). The resulting immunotoxins were concentrated to approximately 1 mg/ml and stored as infusion concentrates in sterile 5 ml glass vials at -20°C at the Department of Clinical Pharmacy. The immunotoxin-combination is prepared directly for use by diluting equal amounts (w/w) of SPV-T3a-dgA and WT1-dgA in formulation buffer.

Immunotoxin composition and biological activity

The composition of purified MoAb-dgA conjugates was analyzed by nonreducing SDS-PAGE using 4-15% gradient gels (PhastSystem, Amersham Pharmacia Biotech). The ratio of different reaction products was determined by densitometric scanning of the separated product bands of Coomassie blue stained gels. Samples containing a 1:1 mixture of unconjugated dgA and MoAb were used for compensation of variance in dye-uptake between both proteins.

Binding capacity of the MoAb before and after conjugation to dgA was examined by a standard flowcytometric titration assay using an antigen positive cell line and a fluorescent labeled goat-anti-mouse antibody to detect cell bound MoAb. Preservation of Ricin A toxicity was determined by measurement of protein synthesis inhibition in a cell free reticulocyte assay.²¹

These analyses were periodically repeated and compared to baseline values to monitor the chemical stability and biological activity of the infusion concentrates during storage.

Sterility, endotoxins and adventitious agents

The FCS added to the culture medium was derived from healthy calves housed in a BSE-free environment, and tested negative for Bovine Diarrhoea virus, Infectious Bovine Rhinotracheitis, Parainfluenza and Bovine Polyoma virus. The MWCB of SPV-T3a and WT1 were tested for bacterial and fungal contamination and for mycoplasma. An 'extended S⁺L⁻ focus assay' and 'extended XC-plaque assay' was performed by Inveresk Research (Tranent, Scotland) for detection of xenotropic and ecotropic retroviruses, respectively. Species specific viruses were tested by the institutional Department of Virology using a mouse antibody production test (MAP-assay), detecting 16 different exogenous mouse viruses including sendai virus, reovirus type 3, Mouse rotavirus (EDIM), hantaan virus, and lymphocytic choriomeningitis virus. The presence of adventitious viruses was tested by the Department of Virology using 8 different indicator cell lines (including 3 human) and by inoculation of embryonated chicken eggs. Purified MoAb were tested for endotoxins, sterility and mycoplasma. The final immunotoxins were tested for the presence of hybridoma DNA by Inveresk Research. The leak-

age of affinity chromatography media was assayed using commercial enzyme immunoassays according to the manufacturer's instructions: Protein A using a kit from Cygnus Technologies (Wrentham, MA, USA), Cybacron Blue with a kit from Affinity Chromatography Ltd (Girton, Cambridge, UK). The stored infusion concentrates were tested for endotoxins, sterility and mycoplasma.

Cross reactivity with human tissues

Potential cross-reactivity of MoAb SPV-T3a and WT1 was tested by immunostaining of the following panel of quickly frozen human tissue sections derived from the tissue bank at the Department of Pathology (# donors): heart (3), lymph node (3), spleen (3), lung (3), esophagus (3), stomach (3), ileum (3), colon (3), liver (3), pancreas (3), testis (3), prostate (3), bladder (3), pyelum (3), kidney (3), ovary (3), Fallopian tube (2), uterus (3), breast (3), cerebrum (3), cerebellum (3), spinal cord (1), skin (3), striated muscle (3), thyroid (3), parathyroid (2), adrenal gland (3), pituitary (2), eyeball (1). Reactivity against hematopoietic cell lines and stem cells has been tested extensively during the initial characterization of the MoAbs. Potential cross-reactivity against embryonic/fetal tissues has not been determined yet since pregnant women are no candidates for allogeneic stem cell transplantation.

In vitro dissociation in human plasma

Dissociation of MoAb and dgA during incubation in human plasma at 37°C was monitored using enzyme immunoassays which detect intact MoAb-dgA conjugates.⁹

LD₅₀ mice

All animal experiments were performed by personnel of the Central University Animal Laboratory in accordance with the 'Principles of Laboratory Animal Care' (NIH publication #85-23, revised 1985). The LD₅₀ experiment was performed with male BALB/C mice of eight weeks weighing ~17 g, as described by Weil.²² Four doses were administered: 96, 193, 385 and 770 µg immunotoxin-combination per mouse, in a final volume of 200 µl. Administration was intravenously through one of the tail veins. Upon administration, mice were weighed daily and followed for survival for two weeks. The control group was injected with formulation buffer only.

Administration to cynomolgus monkeys

The monkeys received two doses of immunotoxin-combination, administered 48 hours apart as 100 ml infusions over a period of 4 hours. Prior to infusion, the monkeys were sedated (ketamine 10 mg/kg, atropine 0,5 mg) and subsequently anesthetized (ratio oxygen and nitrous oxide 1:2, enflurane 1,5%). The immunotoxin-combination was administered via a subclavian (first infusion) or femoral (second infusion) central venous catheter. For detection of any esophagus toxicity, endoscopy was performed on day -7, 4 and 9 (day 1 being the first infusion day). During endoscopy, biopsies were taken for microscopic analysis of potential tissue damage. Moreover, biopsy derived tissue was stained with a peroxidase-labeled anti-mouse-IgG2b antibody for detection of

SPV-T3a-dgA. Tissue pre-incubated *in vitro* with saturating amounts of SPV-T3a-dgA served as positive control. In addition, both monkeys underwent regular physical examinations and blood chemistries, complete blood counts and leukocyte differentials were obtained for general safety assessment. One of the monkeys (receiving 0.1 and 0.2 mg/kg immunotoxin-combination) suffered from diabetes mellitus, but showed no physical disorders which were expected to influence the outcome of the experiment.




Figure 1 Immunotoxin preparation

MoAb and dgA are derivatized with SMPT (m -fold, with $m = 0, 1, 2, \dots$) and reduced with DTT, respectively. The purified reaction mixtures (Sephadex G25) are mixed and incubated at room temperature for 48 hours. The resulting crude conjugate, containing protein aggregates, MoAb conjugated to dgA (n -fold, with $n \leq m$), and non-conjugated molecules, are then separated by BlueSepharose-affinity chromatography and S300-size exclusion chromatography.

RESULTS

Immunotoxin production

The crude harvest of SPV-T3a and WT1 contained approximately 50 $\mu\text{g}/\text{ml}$ MoAb. The recovery following Protein A purification was $\sim 85\%$, about 1 g of purified SPV-T3a and WT1 were produced. The subsequent conjugation process is schematically depicted in figure 1. To reduce the risk of losing all material by a technical failure, MoAbs were divided in two fractions which were conjugated separately to dgA and subsequently partly purified by BlueSepharose. The MoAb-dgA containing BlueSepharose eluates of each MoAb were then pooled and applied as one fraction to the Sephacryl S-300 column. The final MoAb-recovery following conjugation was 34% and 26% for SPV-T3a and WT1, respectively. Figure 2 displays a representative non-reducing SDS-PAGE analysis of an SPV-T3a-dgA conjugation.




Figure 2 SDS-PAGE analysis of SPV-T3a-dgA conjugation

A: high molecular weight marker; B: SPV-T3a (IgG2a); C: reduced dgA; D: crude conjugate; E: run-through fraction BlueSepharose; F: high-affinity eluate BlueSepharose; G: pooled fractions of Sephacryl S300HR eluate containing SPV-T3a-dgA.

Table 1 summarizes the main characteristics of the final products, the left column displaying the proposed release criteria. Ideally, the final product consists entirely of MoAb conjugated to one or two dgA molecules. In practice, the final product contains some unconjugated MoAb, and MoAb bound to three or more dgA molecules, as well. The composition of WT1-dgA did not strictly meet the (arbitrary) release criteria as for its relatively high amount of free MoAb (23% instead of <10%). However, the acceptance of WT1-dgA in its present form was justified by the absence of 'contamination' with MoAb conjugated to three or more dgA-molecules, as well as by the high preservation of antigen-binding activity (90%) and effective *in vitro* killing capacity.⁹ Another point of

Table 1 *Product specifications*

	Proposed Release Criteria	SPV-T3a-dgA (CD3)	WT1-dgA (CD7)
MoAb-Isoform	IgG2bκ/IgG2aλ	IgG2bκ	IgG2aλ
Composition:			
MoAb(dgA) _{≥3}	<10%	6%	0%
MoAb(dgA) ₁₋₂	>80%	88%	78%
MoAb	<10%	6%	23%
dgA	<3%	<1% (detection limit)	<1%
Sterility	Sterile	Sterile	Sterile
Endotoxins (EU/mg protein)	< 20	< 0.8	< 0.7
DNA content (pg/mg protein)	< 5	< 11	< 9
Protein A (ng/mg protein)	< 10	1.1 ± 0.3	2.7 ± 0.8
Cybacron Blue (nM)	< 10 (detection limit)	< 10	< 10
A-chain activity (% native dgA)	> 75	84	91
Binding activity (% native MoAb)	> 50	70	90
Dissociation in vitro (% in 24h)	~15	10	10
<i>In vitro</i> cross reactivity	Absent. If not, relevant animal studies should follow	Esophagus epithelium (++) Smooth muscle cells (+/-)	Kupfer cells (+/-)
LD ₅₀ in mice (μg/g)	~14 μg/g mouse	25-45 μg/g mouse for the IT-combination	
Toxicity for cynomolgus monkey	No severe irreversible toxicities	Reversible toxicity to skeletal muscles after 0.1-0.25 mg/kg IT-combination	

attention was the presence of genomic DNA. The IT-combination contains <10 pg DNA/mg protein, being the detection limit of the assay. The relevant FDA's 'Points to Consider in the Manufacture and Testing of Monoclonal Antibody Products for Human Use' suggests that, whenever possible, the final product contains no more than 100 pg cellular DNA per dose. This implies that total human doses of 10 mg and higher (equalizing ~5-7 mg/m²) acquire additional testing (e.g. with concentrated immunotoxin to lower the detection limit).

So far, the repeated testing for composition, MoAb-binding and dgA-activity demonstrated stability of both SPV-T3a-dgA and WT1-dgA at -20°C for a minimum period of three years.

Cross reactivity with human tissues

MoAb SPV-T3a showed two unexpected cross-reactivities: against (1) smooth muscle cells, and (2) basal epithelial cells of the esophagus. The staining of smooth muscle cells was weak (+/-) but consistently found throughout various tissues. The diffuse staining pattern suggested a cytoplasmatic binding of physically disrupted cells. This is supported by the observation that SPV-T3a-dgA (10⁻⁸M for 24 hours) appeared to be non-toxic to cultured intact human leptomeningial smooth muscle cells (data not shown). The staining intensity of SPV-T3a with the esophagus-basal epithelium was as strong (++) as observed with positive lymphocytes, and appeared to be characteristic for SPV-T3a since it was not observed with any of the other CD3 MoAbs tested (OKT3, UCHT and WT32). The *in vivo* consequence of this cross reactivity will depend on the local concentration SPV-T3a-dgA, and whether SPV-T3a-dgA will be internalized (a prerequisite for toxicity). To gain more insight in these parameters, it was decided to administer the immunotoxin-combination to cynomolgus monkeys, demonstrating *in vitro* cross-reactivity of comparable intensity and localization.

The screening with MoAb WT1 revealed a weak staining (+/-) of some Kupffer cells located in the liver sinus wall. This staining pattern was not observed during previous preclinical tissue screenings performed with WT1.²³ No further unexpected cross-reactivities were observed.

LD₅₀ mice

The LD₅₀ in mice was determined for comparison with reference immunotoxin RFB4-dgA, a comparable immunotoxin already tested in FDA-approved phase I/II trials (having a LD₅₀ of 14 mg/kg).²⁴ Three of the four mice treated with the highest dose immunotoxin-combination (770 µg/mouse) died within the follow-up period of two weeks. One mouse already died during injection, probably due to an administration-related toxicity (dgA-related deaths are normally first seen after 1-2 days). All mice treated with 385 µg immunotoxin-combination or less survived. Based on the average initial weight, this implies an LD₅₀ of 25 - 45 mg/kg mouse.

Apart from the survival score, mice were daily weighed and observed for behavior. Administration of IT combination caused weight loss in all groups, which was most pronounced after 9 days (Figure 3). The extent and duration correlated well with the dosage. Besides weight-loss, mice treated with 770 or 385 μg IT-combination demonstrated lethargy, lack of appetite and a neglected fur. The mice treated with the two lower doses could not be distinguished visually from the control group.




Figure 3 Average weight-changes of immunotoxin treated mice

Mice were treated with formulation buffer (○) or with 96 μg (●), 193 μg (■), 385 μg (▲) or 770 μg immunotoxin-combination/mouse (◆). Upon administration, mice were weighed daily for two weeks.

Administration of the immunotoxin-combination to cynomolgus monkeys

The immunotoxin-combination was administered to two male cynomolgus monkeys. As the administration was performed under general anesthesia, the animals had to be deprived of nutrition before each infusion. Since this may negatively impact their physical condition, the administration was restricted to two doses at a 48-hour interval (the patients in the pilot-study were supposed to receive four doses at the same interval). This decision was justified by observations of Amlot *et al.* who demonstrated that the toxicity of a dgA-based immunotoxin mainly depends on the height of the individual doses and not on the number of doses administered.²⁴ Both monkeys received two of the higher individual doses of the proposed human dose escalation scheme (ranging from 2 to 10 mg/m^2), translated to mg/kg . One monkey (about 12 years and 7.6 kg) received 0.1 and 0.2 mg/kg , the other monkey (4 years and 4.1 kg), received 0.2 and 0.25 mg/kg (equalizing human doses of 4, 8 and 10 mg/m^2). In human, the immunotoxin-combination will be given as a 100 ml infusion over 4 hours. For the monkeys, this was changed to 20 ml administered in 1 hour because of their smaller blood volume.




Figure 4 Immunotoxin-combination plasma concentrations in cynomolgus monkeys

Two male cynomolgus monkeys were each given two doses immunotoxin-combination with an interval of 48 hours (arrows beneath the x-axis). One monkey received 0.1 and 0.2 mg/kg (□), the other 0.2 and 0.25 mg/kg (●).

Figure 4 shows the plasma concentrations of the immunotoxin-combination. The clearance curves best fitted a two-compartment model for both SPV-T3a-dgA and WT1-dgA individually and given in combination. The pharmacokinetic parameters are listed in

Table 2 Pharmacokinetic parameters (\pm SD)^a

Monkey (dose, mg/kg)	Immunotoxin	$T_{1/2\beta}$ (h)	AUC ($\mu\text{g}\cdot\text{h}/\text{ml}$)	Cl (ml/h)	V_d (ml)	C_{\max} (ng/ml)
A (0.1 and 0.2)	SPV-T3a-dgA	9.1 ± 0.6	9.4 ± 1.6	26.5 ± 4.5	347.3 ± 29.1	1,830
	WT1-dgA	9.4 ± 1.6	12.8 ± 1.7	19.6 ± 2.6	265.8 ± 17.5	2,090
	Combination	9.3 ± 1.7	22.3 ± 3.1	4.5 ± 0.6	60.3 ± 4.2	3,920
B (0.2 and 0.25)	SPV-T3a-dgA	9.3 ± 2.0	23.5 ± 3.8	8.5 ± 1.4	113.6 ± 9.2	2,280
	WT1-dgA	2.2 ± 0.2	4.7 ± 0.3	42.9 ± 3.1	133.7 ± 7.4	1,450
	Combination	5.7 ± 1.0	24.8 ± 3.3	8.1 ± 1.1	65.8 ± 4.6	3,840

^aAbbreviations: $T_{1/2\beta}$, plasma half-life; AUC, area under the concentration versus time curve; Cl, clearance; V_d , volume of distribution; and C_{\max} , maximum plasma concentration.

Table 2. The $T_{1/2}$ of the immunotoxin-combination was 9.3 and 5.7 hours in monkeys A and B, respectively. Notably the $T_{1/2}$ of SPV-T3a and WT1 in 'monkey B' differed considerably, being approximately 9 and 2 hours, respectively. This may be explained by the capture of WT1-dgA by the CD7-antigen expressed on T and NK cells (virtually all these cells being CD7⁺ in 'monkey B'). 'Monkey A' with only ~30% of its T/NK cells being

CD7⁺, demonstrated equal half lives for SPV-T3a-dgA and WT1-dgA (~9 hours). It should be noted that SPV-T3a-dgA is not influenced by expression of the CD3-antigen as SPV-T3a does not bind 'monkey-CD3'. Peak plasma levels were attained directly following each infusion and decreased (nearly) to baseline level in about 48 hours. The C_{max} of the immunotoxin-combination ranged from 1.8 to 3.9 $\mu\text{g/ml}$ and strongly correlated with the dose administered. In general, plasma levels of ~1.8 $\mu\text{g/ml}$ (10^{-8} M) are considered therapeutic concentrations.²⁵ In both monkeys, these plasma concentrations were maintained for 8-10 hours following the second infusion.




Figure 5 Serum CK-levels in cynomolgus monkeys

Following administration of the immunotoxin-combination, both cynomolgus monkeys demonstrated a transient rise of serum CK-levels. One monkey received 0.1 and 0.2 mg/kg (□), the other 0.2 and 0.25 mg/kg (●) immunotoxin-combination, separated by a 48-hour interval.

Regarding the cross-reactivity of SPV-T3a, neither macroscopic nor microscopic examinations revealed any evidence of toxicity towards the esophagus. Moreover, immunostaining of biopsy derived tissue could not demonstrate any *in vivo* binding of SPV-T3a-dgA to the esophagus epithelium. Blood chemistry analyses revealed a transient drug-related toxicity. Each administration was followed by a sharp increase in serum creatine kinase (CK), reaching a maximum at 7 hours after infusion and normalizing to baseline within 2-7 days (Figure 5). Following the second peak, both monkeys demonstrated a third, more modest, rise of CK-levels that might be explained by a secondary release of dgA. The increase in CK levels is indicative for destruction of striated muscles and/or heart muscle. Damage of the heart muscle could be excluded by measurement of heart-muscle isomer CKMB. In accordance with the destruction of striated

muscle cells, a concomitant (modest) increase in LD, ASAT and ALAT serum levels could be noted. Four weeks after completion of the study all biochemical parameters had returned to baseline levels. Complete blood counts and leukocyte differentials displayed a normal distribution. Immediately after the infusions, both monkeys seemed a bit weakened. Aside from the general anesthesia, this may have resulted from the destruction of striated muscle tissue. One day after the second dose, both monkeys ate almost as normal again. Within a week, both monkeys functioned as before treatment.

DISCUSSION

This report describes the laboratory preparation and control of two dgA-based immunotoxins that are currently under evaluation in a clinical pilot-study. The design of this study was strongly inspired by earlier reports on RFB4-dgA, a comparable immunotoxin for the treatment of Non-Hodgkin's lymphoma.^{24,26} The murine LD₅₀ of the immunotoxin-combination compared favorably with that of RFB4-dgA (25-45 mg and 14 mg per kg mouse, respectively).¹⁹ Moreover, the monkey experiment demonstrated pharmacokinetics resembling those previously observed with RFB4-dgA and comparable immunotoxins in man. Importantly, clinically significant plasma concentrations were reached for several hours without being accompanied by acute irreversible toxicities. With respect to the potential cross-reactivity of SPV-T3a-dgA, it was particularly of interest that no binding or toxicity to the esophagus was observed. The main toxicity observed was a clear but transient rise in CK-levels. This is in accordance with myalgias, sometimes associated with rhabdomyolysis, being the second dose limiting toxicity associated with dgA-immunotoxins in man (after vascular leak syndrome, VLS).²⁵ Consistent with the human studies, this toxicity resolved upon ending the administration of immunotoxin. It should be noted, though, that the animal studies do not eliminate all uncertainties with regard to clinical administration of the immunotoxin-combination. The esophagus of patients with severe graft-versus-host disease may be more accessible for SPV-T3a-dgA (*e.g.* due to disease-associated lesions) than those of healthy cynomolgus monkeys. Another concern is that monkeys are hardly vulnerable for dgA-induced VLS, the main toxicity reported in man. However, as these issues can not be clarified by animal studies, the current data were considered maximally supportive for performing the clinical pilot-study.

So far, the preliminary results of the clinical pilot-study are very encouraging and do warrant the set up of multi-center Phase I/II studies. Consequently, in addition to a further scaling-up of production, product-specifications will probably have to meet stricter criteria. A point for improvement forms the reduction of the relatively high amount of unconjugated WT₁. Moreover, stricter guidelines might ask for future MoAb productions to be performed in serum-free medium. This inevitably will result in repeated

hybridoma cell banking and subsequent virus testing, as changes in growing conditions might provoke the expression of different latent viruses. As an additional requirement, the actual viral removing capacity of the production process will need to be demonstrated by virus spiking using a scale-down model. Though the FDA foresees in the human testing of (modified) MoAb combinations, it is anticipated that the required animal toxicity studies (*e.g.* acute/subchronic toxicity and reproduction toxicity studies) will have to be performed with both immunotoxins separately.

Notwithstanding the foreseen adaptations, the current batch is very well suited for its intended use: the evaluation in a clinical pilot study for the treatment of a life-threatening disease. As such, the described production forms a stepping stone towards the future productions as required for the actual registration process. It provided sufficient quantities of proper quality to enable the clinical evaluation in an academic setting. Especially for niche indications like severe graft-versus-host disease, some clinical proof of concept appears a prerequisite for further development by the pharmaceutical industry. The in house production and clinical evaluation of biological therapeutics by academic centers might therefore contribute increasingly to the development of new drugs within this area.

ACKNOWLEDGEMENTS

We would like to thank Mr. Albert Peters and colleagues for taking care of the test animals, Dr. Fokke Nagengast for performing endoscopy and Dr. Roos Masereeuw for pharmacokinetic analysis. The RIVM (Bilthoven, The Netherlands) is acknowledged for generously providing the chromatography columns. This work was supported by a grant from the Technology Foundation STW (Utrecht, The Netherlands).

REFERENCES

1. Vitetta ES, Thorpe PE, Uhr JW: Immunotoxins: magic bullets or misguided missiles? *Immunol Today* 14:252-9, 1993
2. Valleria DA: Immunotoxins: will their clinical promise be fulfilled? *Blood* 83:309-17, 1994
3. Kreitman RJ: Immunotoxins in cancer therapy. *Curr Opin Immunol* 11:570-8, 1999
4. Byers VS, Henslee PJ, Kernan NA, Blazar BR, Gingrich R, Phillips GL, LeMaistre CF, Gilliland G, Antin JH, Martin P, et al.: Use of an anti-pan T-lymphocyte ricin A chain immunotoxin in steroid-resistant acute graft-versus-host disease. *Blood* 75:1426-32, 1990
5. Vitetta ES, Stone M, Amlot P, Fay J, May R, Till M, Newman J, Clark P, Collins R, Cunningham D, Ghetie V, Uhr JW, Thorpe PE: Phase I immunotoxin trial in patients with B-cell lymphoma. *Cancer Res* 51:4052-8, 1991
6. Engert A, Diehl V, Schnell R, Radszuhn A, Hatwig MT, Drillich S, Schon G, Bohlen H, Tesch H, Hansmann ML, Barth S, Schindler J, Ghetie V, Uhr J, Vitetta E: A phase-I study of an anti-CD25 ricin A-chain immunotoxin (RFT5-SMPT-dgA) in patients with refractory Hodgkin's lymphoma. *Blood* 89:403-10, 1997
7. Lynch TJ, Jr., Lambert JM, Coral F, Shefner J, Wen P, Blattler WA, Collinson AR, Ariniello PD, Braman G, Cook S, Esseltine D, Elias A, Skarin A, Ritz J: Immunotoxin therapy of small-cell lung cancer: a phase I study of Ng01- blocked ricin. *J Clin Oncol* 15:723-34, 1997
8. Kreitman RJ, Wilson WH, White JD, Stetler-Stevenson M, Jaffe ES, Giardina S, Waldmann TA, Pastan I: Phase I trial of recombinant immunotoxin anti-Tac(Fv)-PE38 (LMB-2) in patients with hematologic malignancies. *J Clin Oncol* 18:1622-36, 2000
9. van Oosterhout YV, van Emst L, Schattenberg AV, Tax WJ, Ruiter DJ, Spits H, Nagengast FM, Masereeuw R, Evers S, de Witte T, Preijers FW: A combination of anti-CD3 and anti-CD7 ricin A-immunotoxins for the in vivo treatment of acute graft-versus-host disease. *Blood* 95:3693-3701, 2000
10. Preijers FW, Tax WJ, De Witte T, Janssen A, v.d. Heijden H, Vidal H, Wessels JM, Capel PJ: Relationship between internalization and cytotoxicity of ricin A-chain immunotoxins. *Br J Haematol* 70:289-94, 1988
11. Chignola R, Colombatti M, Dell'Arciprete L, Candiani C, Tridente G: Distribution of endocytosed molecules to intracellular acidic environments correlates with immunotoxin activity. *Int J Cancer* 46:1117-23, 1990
12. May RD, Wheeler HT, Finkelman FD, Uhr JW, Vitetta ES: Intracellular routing rather than cross-linking or rate of internalization determines the potency of immunotoxins directed against different epitopes of sIgD on murine B cells. *Cell Immunol* 135:490-500, 1991
13. van Oosterhout YV, van den Herik-Oudijk IE, Wessels HM, de Witte T, van de Winkel JG, Preijers FW: Effect of isotype on internalization and cytotoxicity of CD19-ricin A immunotoxins. *Cancer Res* 54:3527-32, 1994
14. van Horssen PJ, van Oosterhout YV, de Witte T, Preijers FW: Cytotoxic potency of CD22-ricin A depends on intracellular routing rather than on the number of internalized molecules. *Scand J Immunol* 41:563-9, 1995
15. Spits H, Keizer G, Borst J, Terhorst C, Hekman A, de Vries JE: Characterization of monoclonal antibodies against cell surface molecules associated with cytotoxic activity of natural and activated killer cells and cloned CTL lines. *Hybridoma* 2:423-37, 1983
16. Tax WJ, Greaves MF, Willems HM, Leeuwenberg HF, Capel PJ, Koene RA: WT1: a monoclonal antibody reactive with T-ALL but not with other leukemias. *Haematol Bluttransfus* 28:139-41, 1983
17. Fulton RJ, Blakey DC, Knowles PP, Uhr JW, Thorpe PE, Vitetta ES: Purification of ricin A1, A2, and B chains and characterization of their toxicity. *J Biol Chem* 261:5314-9, 1986

18. Thorpe PE, Wallace PM, Knowles PP, Relf MG, Brown AN, Watson GJ, Blakey DC, Newell DR: Improved antitumor effects of immunotoxins prepared with deglycosylated ricin A-chain and hindered disulfide linkages. *Cancer Res* 48:6396-403, 1988
19. Ghetie V, Thorpe P, Ghetie MA, Knowles P, Uhr JW, Vitetta ES: The GLP large scale preparation of immunotoxins containing deglycosylated ricin A chain and a hindered disulfide bond. *J Immunol Methods* 142:223-30, 1991
20. Carlsson J, Drevin H, Axen R: Protein thiolation and reversible protein-protein conjugation. N- Succinimidyl 3-(2-pyridyldithio)propionate, a new heterobifunctional reagent. *Biochem J* 173:723-37, 1978
21. Press OW, Vitetta ES, Martin PJ: A simplified microassay for inhibition of protein synthesis in reticulocyte lysates by immunotoxins. *Immunol Lett* 14:37-41, 1986
22. Weil CS: Tables for convenient calculation of effective dose (LD₅₀ of ED₅₀) and intructions in their use. *Biometrics* :249-53, 1952
23. Hertler AA, Schlossman DM, Borowitz MJ, Poplack DG, Frankel AE: An immunotoxin for the treatment of T-acute lymphoblastic leukemic meningitis: studies in rhesus monkeys. *Cancer Immunol Immunother* 28:59-66, 1989
24. Amlot PL, Stone MJ, Cunningham D, Fay J, Newman J, Collins R, May R, McCarthy M, Richardson J, Ghetie V, Ramilo O, Thorpe PE, Uhr JW, Vitetta ES: A phase I study of an anti-CD22-deglycosylated ricin A chain immunotoxin in the treatment of B-cell lymphomas resistant to conventional therapy. *Blood* 82:2624-33, 1993
25. Ghetie V, Vitetta E: Immunotoxins in the therapy of cancer: from bench to clinic. *Pharmacol Ther* 63:209-34, 1994
26. Sausville EA, Headlee D, Stetler-Stevenson M, Jaffe ES, Solomon D, Figg WD, Herdt J, Kopp WC, Rager H, Steinberg SM, et al.: Continuous infusion of the anti-CD22 immunotoxin IgG-RFB4-SMPT-dgA in patients with B-cell lymphoma: a phase I study. *Blood* 85:3457-65, 1995

**A COMBINATION OF ANTI-CD₃ AND
ANTI-CD₇ RICIN A-IMMUNOTOXINS FOR THE
IN VIVO TREATMENT OF ACUTE
GRAFT-VERSUS-HOST DISEASE**

Ypke V.J.M. van Oosterhout

J. Liesbeth van Emst

Anton V.M.B. Schattenberg

Wil J.M. Tax

Dirk J. Ruiter

Hergen Spits

Fokke M. Nagengast

Roos Masereeuw

Sabine Evers

Theo de Witte

Frank W.M.B. Preijers



Blood 2000; 95: 3693-3701

ABSTRACT

This study evaluated the anti-graft versus host disease (GVHD) potential of a combination of immunotoxins (ITs), consisting of a murine CD3 (SPV-T3a) and CD7 (WT1) monoclonal antibody both conjugated to deglycosylated ricin A. *In vitro* efficacy data demonstrated that these IT act synergistically, resulting in an approximately 99% elimination of activated T cells at 10^{-8} M (about 1.8 μ g/ml). Because most natural killer (NK) cells are CD7⁺, NK activity was inhibited as well. Apart from the killing mediated by ricin A, binding of SPV-T3a by itself impaired *in vitro* cytotoxic T-cell cytotoxicity. Flow cytometric analysis revealed that this was due to both modulation of the CD3/T-cell receptor complex, and activation-induced cell death. These results warranted evaluation of the IT-combination in patients with refractory acute GVHD in an ongoing pilot study. So far, 4 patients have been treated with 3 to 4 infusions of 2 or 4 mg/m² IT-combination, administered intravenously at 48-hour intervals. The $T_{1/2}$ was 6.7 hours, and peak serum levels ranged from 258 to 3,210 ng/ml. Drug-associated side effects were restricted to limited edema, fever, and a modest rise of creatine kinase levels. One patient developed low-titer antibodies against ricin A. Infusions were associated with an immediate drop of circulating T cells, followed by a more gradual but continuing elimination of T/NK cells. One patient mounted an extensive CD8 T-cell response directly following treatment, not accompanied with aggravating GVHD. Two patients showed nearly complete remission of GVHD, despite unresponsiveness to the extensive pretreatment. These findings justify further investigation of the IT-combination for treatment of diseases mediated by T cells.

INTRODUCTION

Stem cell transplantation forms a widely accepted method for restoration of normal hematopoiesis of patients treated for a hematological malignancy or otherwise suffer-

ing from a defective hematopoietic or immunologic system. For a successful engraftment, a minimal number of donor T cells in the graft appears to be a prerequisite. The underlying mechanism by which these cells promote engraftment is not fully understood, but probably includes the creation of an immunologically tolerogenic environment by eliminating the remainder of the patient's immune system.

In treatment of a malignancy, the cotransplantation of donor T cells confers additional benefit because they contribute to the so-called graft-versus-leukemia (GVL) effect, which involves the elimination of residual malignant cells.¹ The basis of GVL forms the recognition by donor T cells of (minor or major) histocompatibility antigens expressed by the malignant recipient cells. Unfortunately, this antigenic disparity may also lead to graft versus host disease (GVHD), a major cause of morbidity and mortality after allogeneic stem cell transplantation.² GVHD is thought to be initiated by alloactivation of donor T cells resulting in the production of cytokines (interleukin [IL]-2 and interferon [INF]- γ) which in turn activate additional effector cells, like monocytes, macrophages, and natural killer (NK) cells, to produce inflammatory proteins (IL-1, tumor necrosis factor [TNF]- α , and IL-6).³ This may result in serious direct and indirect cytotoxicity to epithelial cells of skin, liver, and gut.

When prophylaxis has failed (typically cyclosporine, often combined with methotrexate), severe GVHD is usually treated with low-dose corticosteroids. If the reaction progresses, the dose may be increased (up to levels of 10-20 mg/kg/day) or, alternatively, polyclonal antithymocyte/antilymphocyte globulin (ATG/ALG) or experimental immunosuppressive drugs may be applied. An example of such an experimental reagent is Xomazyme-CD5 Plus, a murine CD5 monoclonal antibody (MoAb) conjugated to the A-chain of the phytolectin ricin. Especially in the initial reports, Xomazyme-CD5 Plus demonstrated substantial efficacy in treating steroid-resistant acute GVHD.^{4,5} In more recent comparative trials, Xomazyme-CD5 Plus was not more effective than high dose corticosteroids or ATG.^{6,7} Encouraged by its initial success, we developed a therapy based on the use of a combination of 2 anti-T-cell immunotoxins (IT): murine MoAb SPV-T3a (CD3) and WT1 (CD7), both conjugated to deglycosylated ricin A (dgA). In this article, we present the preclinical efficacy data and preliminary clinical results, both of which suggest that this IT-combination has the potential for helping to control severe acute GVHD.

MATERIALS AND METHODS

Immunotoxins

The "IT-combination" as referred to in this article, consists of a 1:1 mixture (w/w) of murine MoAb SPV-T3a (CD3)⁸ and WT1 (CD7)^{9,10} both conjugated to deglycosylated ricin A (dgA) (Inland, Austin, TX) using the SMPT-cross-linker (Pierce, Rockford, IL),

according to Ghetie *et al.*¹¹ The preparation and validation were supervised by the institutional Department of Clinical Pharmacy. The major characteristics of the IT are summarized in Table 1; details will be reported elsewhere (manuscript in preparation). Murine MoAb UPC-10 (IgG2a) and MOPC-141 (IgG2b) (both Sigma, St. Louis, MO) conjugated to dgA served as isotype-matched irrelevant controls for the *in vitro* experiments. Noteworthy, *in vitro* tissue screening unveiled a cross-reactivity of MoAb SPV-T3a with the basal epithelium of the human esophagus. To investigate potential *in vivo* consequences, the IT-combination was administered to two cynomolgus monkeys sharing the same cross reactivity. Subsequent esophagus biopsies revealed no toxicity towards the esophagus epithelium, nor could any localization of SPV-T3a-dgA be detected. Both monkeys demonstrated a transient rise of creatine kinase (CK)-levels upon administration of the IT-combination.

Table 1 IT characteristics

	SPV-T3a-dgA (CD3)	WT1-dgA (CD7)
MoAb	murine IgG2bκ	murine IgG2aλ
Composition (SDS-PAGE)	MoAb(dgA) _{≥3} 6%	MoAb(dgA) _{≥3} 0%
	MoAb(dgA) ₁₋₂ 88%	MoAb(dgA) ₁₋₂ 78%
	MoAb 6%	MoAb 23%
	dgA 0%	dgA 0%
Sterility	Sterile	Sterile
Endotoxins (EU/mg protein)	< 0.8	< 0.7
DNA content (pg/mg protein)	< 11	< 9
Protein A (ng/mg protein)	1.1 ± 0.3	2.7 ± 0.8
Cybacron Blue (nM)	< 10	< 10
A-chain activity (% native dgA)	84	91
Binding activity (% native MoAb)	70	90
Dissociation in human plasma at 37°C (% in 24h)	10	10
<i>In vitro</i> cross reactivity	Esophagus epithelium (++) Smooth muscle cells (+/-)	Kupfer cells (+/-)
LD ₅₀	25-45 µg/g mouse for the IT-combination	
Toxicity cynomolgus monkey	Reversible increase of CK-levels upon administration of human doses IT-combination (0.1-0.25 mg/kg)	

Peripheral blood mononuclear cells (PBMC)

PBMC were isolated from peripheral blood by Ficoll-Hypaque density centrifugation (Pharmacia, Uppsala, Sweden), and cultured in Iscove's medium (Flow, Irvine, Scotland) supplemented with 10% heat-inactivated Pooled Human Serum (PHS), penicillin (100 U/ml) and streptomycin (100 µg/ml). For generation of PHA-activated T cells, PBMC were preincubated with 40 µg/ml phytohaemagglutinin (PHA-HA15) (Murex, Dartford, UK) for 48 hours.

Flow cytometric quantification of in vitro cell kill

Nonactivated as well as PHA-activated PBMC ($10^6/\text{ml}$) were incubated in triplicate with varying concentrations of IT (10^{-13} - 10^{-8} M) at 37°C for 24 hours. Following treatment, cells were washed and cultured in the presence of PHA ($40\text{ }\mu\text{g}/\text{ml}$) for an additional four days to enable full exposure of IT toxicity. Subsequently, cells were labeled with propidium iodine (PI) (Molecular Probes, Junction City, OR) and calcein AM (Calc) (Molecular Probes) (both: $2\text{ }\mu\text{g}/\text{ml}$ for 30 minutes at room temperature), and analyzed on a flow cytometer (Coulter Epics Elite, Hialeah, FL). Cells being PI⁻ and Calc⁺ were referred to as viable cells. Prior to FCM analysis, a fixed amount of inert beads (Flow-Count fluorescent spheres, Coulter) was added to each sample ($1 \times 10^4/\text{ml}$) to enable the quantification of surviving cells.

Reduction of CTL toxicity by SPV-T3a

Cytotoxic T cell (CTL) toxicity was assayed *in vitro* using an EBNA3C12-reactive CTL clone. The CTL clone was incubated with MoAb SPV-T3a (10^{-8} M), or isotype matched control MoAb MOPC-141, at 37°C for 24 hours. Subsequently, cells were washed and cultured in culture medium for another 72 hours. Remaining cytolytic activity was assayed with an autologous Epstein-Barr virus (EBV)-transformed lymphoblastoid cell line (EBV-LCL), labeled with $100\text{ }\mu\text{Ci}$ ^{51}Cr (Amersham, Bucks, UK) at 37°C for 2 hours. Labeled EBV-LCL were plated in triplicate ($10^3/\text{well}$) in V-bottom microtiter plates (Greiner, Frickenhausen, Germany) and saturated with endogenous EBNA3C ($5\text{ }\mu\text{M}$ for 1 hour at 37°C) to enhance T-cell receptor (TCR)-mediated lysis. Subsequently, varying numbers of MoAb-treated CTL cells were added to each well in a final volume of $150\text{ }\mu\text{l}$ culture medium. Plates were centrifuged (50g , 1 minute) and further incubated at 37°C . After 4 hours, $100\text{ }\mu\text{l}$ supernatant was collected from each well and counted in a gamma counter. Specific lysis was expressed as percentage maximal lysis by detergent, both corrected for spontaneous ^{51}Cr -release.

Modulation of the CD3-antigen by SPV-T3a

Surface CD3 expression of SPV-T3a-treated CTL was determined by indirect fluorescence staining with a saturating amount of SPV-T3a ($10\text{ }\mu\text{g}/\text{ml}$, 4°C 30 minutes), followed by a fluorescein isothiocyanate (FITC)-conjugated F(ab')₂ goat-anti-mouse IgG (American Qualex International, La Mirada, CA). CD3-antigen still bound by SPV-T3a used for treatment was identified by staining with the FITC-conjugated antibody only. Expression was indicated as percentage relative to untreated control cells.

In vitro reduction of NK-activity.

The PBMC ($10^6/\text{ml}$) were incubated with 10^{-8}M MoAb or IT for 24 hours. Subsequently, cells were washed and cultured for 3 additional days without MoAb/IT (to enable full exposure of toxicity). PBMC were serially diluted in 96-well U-bottomed plates, and a fixed concentration of ^{51}Cr -labeled K562 cells was added ($10^4/\text{well}$) to yield effector/target ratios of 10:1 to 0.37:1 in a final volume of $150\text{ }\mu\text{l}$ culture medium. Plates were cen-

trifuged (50g, 1 minute) and further incubated at 37°C. After 4 hours, 100 µl supernatant was collected from each well and counted in a gamma counter. NK-activity was expressed as percentage maximum lysis by detergent, both corrected for spontaneous ⁵¹Cr-release. Recombinant IL-2 (500 U/ml) (Cetus, Emeryville, CA) was present during the entire assay to increase NK-activity.

Clinical pilot study

The first patients were treated in a single center, non-randomized, open-labeled, dose-escalating study (ongoing) with the aim of obtaining estimates of the safety and efficacy of the IT-combination administered to patients with life-threatening GVHD. Patients with acute GVHD were *eligible* if they had received second-line high-dose corticosteroid therapy (methylprednisolone 1000 mg/d) for at least 3 days without decrease in the severity of clinical symptoms. Patients were *not eligible* if they had evidence of intrapulmonary disease (which might aggravate the clinical severity of vascular leak syndrome [VLS]¹³), or had allergy or antibodies to mouse Ig or dgA. Before entering the trial, all patients gave informed consent in accordance with the institutional review board and ethics committee. For assessment of safety and responses, patients were evaluated daily during hospitalization, and weekly thereafter, with a physical examination and by obtaining serum chemistries, complete blood counts and leukocyte differential. In addition, blood samples were collected for determination of the pharmacokinetics and immunogenicity of the IT-combination. Toxicities were graded according to the National Cancer Institute Common Toxicity Criteria (NCI-CTC; version 2.0).

IT-combination administration

Patients were supposed to receive 4 doses of the IT-combination administered intravenously in 4-hour infusions at 48-hour intervals. Prior to therapy, an intravenous test dose of 200 µg IT-combination was administered to rule out anaphylactoid reactions. Immunosuppressive agents used for prophylaxis and initial treatment of GVHD were allowed to remain unchanged.

Pharmacokinetics

Plasma levels of intact IT were measured with an enzyme immunoassay. Affinity-purified rabbit antiricin antibody (Sigma) in carbonate buffer (pH 9.6) was adsorbed to 96-wells maxisorp plates (Nunc, Roskilde, Denmark), and residual binding sites were blocked with 4% bovine serum albumin (BSA). Patient plasma samples were serially diluted in 50% PHS in the adsorbed microtiter plates and incubated at 37°C for 1 hour. Subsequently, plates were washed and alkaline-phosphatase conjugated goat anti-mouse IgG2b or IgG2a (SBA, Birmingham, MI) was added at 37°C for 1 hour to bind captured SPV-T3a-dgA or WT1-dgA, respectively. Plates were washed, and the reaction was developed using p-nitrophenylphosphate. Optical densities were read at 450 nm, and levels of circulating IT were calculated from standard curves obtained with a known concentration of IT-combination diluted in pretreatment plasma. The detection limit

was 0.02 $\mu\text{g/ml}$ for both SPV-T3a-dgA and WT1-dgA. Plasma concentration versus time data were analyzed using the nonlinear least square regression program WinNonlin version 1.1 (Scientific Consulting, Apex, NC).

Measurement of HAMA and HARA

An enzyme immunoassay was used for detection of antibody responses to components of the IT-combination. Patient plasma samples were serially diluted (undiluted to 1:2,048) in phosphate-buffered saline with 1% human serum albumin in 96-wells microtiter plates (Nunc) containing either adsorbed SPV-T3a, WT1, or dgA. Bound human antibodies were probed using alkaline phosphatase-conjugated goat antihuman IgG/IgM (H+L) antibodies (Jackson, West Grove, PA), and the reaction was developed with p-nitrophenylphosphate. Optical densities were read at 450 nm, and the serum titer was expressed as the end-point dilution. Positive titers were considered to be at least 3 times background.

Flow Cytometry

Lymphocyte phenotyping was performed by multicolor flow cytometry using directly fluorescent-labeled MoAb according to a whole blood lysis method (FACS lysing, Becton Dickinson, San Jose, CA). T cells and NK cells were identified simultaneously with a mixture of fluorescence-labelled CD2 and CD5 MoAb (MT910-PE and DK23-PE, respectively) (DAKO, Copenhagen, Denmark). The percentage T/NK cells relative to all leukocytes was determined by costaining with a CD45-FITC MoAb (J33-FITC) (Immunotech, Marseille, France), and converted to absolute numbers based on the leukocyte count. B cells were quantified accordingly, using a PE-conjugated CD19 MoAb (HD37-PE) (DAKO).

Staging of GVHD and Definitions of Clinical Responses

Each organ system was staged grade 1 through 4 acute GVHD according to established criteria.¹⁴ Patients were also given an overall grade of acute GVHD (I through IV) based on severity of organ involvement.¹⁴ Responses to therapy were defined as follows. A complete response (CR) as the disappearance of symptoms in all organ systems. A partial response (PR) as the improvement of 1 organ or more, with no worsening in other organs. A mixed response (MR) as the improvement of 1 organ or more, with worsening in 1 other organ or more. Stable disease (SD) as no significant change in any organ system. Progressive disease (PD) as progression in 1 organ system or more without improvement in other organs.

RESULTS

In vitro elimination of T cells

T cells may become more susceptible to dgA-IT on activation.¹⁵ This would be beneficial for the treatment of immunological disorders. Ideally, such treatment inhibits ongoing or beginning immune responses without affecting the resting T cell pool with its ability to counter future hazardous infections. To address this issue, nonactivated and PHA-activated PBMC were treated with IT for 24 hours and analyzed by flow cytometry for the number of surviving cells (Figure 1). Without prior PHA activation, treatment with SPV-T3a-dgA and WT1-dgA, either alone or in combination, resulted in a 2.0- to 2.7-fold reduction of viable cells at 10^{-8} M (about 1.8 μ g/ml, the highest nontoxic concentration *in vitro*). After 48 hours PHA-activation, SPV-T3a-dgA, WT1-dgA, and the combination demonstrated an approximate 3-, 11-, and 34-fold increase in maximal killing capacity, respectively. The combination appeared very effective at concentrations above 10^{-9} M, resulting in the elimination of 99% of activated T cells at 10^{-8} M. Costaining with lineage-specific markers revealed no difference in vulnerability between CD4⁺ and CD8⁺ T-cell subsets (data not shown). Neither incubation with isotype-matched control IT, nor with MoAb SPV-T3a or WT1, significantly reduced the number of viable cells (< 1.15-fold reduction at 10^{-8} M; $n = 3$, data not shown).




Figure 1 Surviving T cells after treatment with IT

Nonactivated (A) and PHA-activated (B) T cells were eliminated by SPV-T3a-dgA (□) and WT1-dgA (○), applied individually and in combination (▲) (half a dose each). Nonactivated and PHA-activated PBMC were incubated with various concentrations IT for 24 hours at 37°C. Following treatment, cells were washed and cultured for an additional four days in the presence of PHA to enable full exposure of IT toxicity. Subsequently, cells were stained with viability markers and analyzed by flow cytometry for the number of viable T cells relative to the untreated control. Data represent the mean \pm SD as obtained with the PBMC of 3 healthy individuals.




Figure 2
Reduction of CTL cytotoxicity
by native MoAb SPV-T3a

An EBNA3C-reactive CTL clone was incubated with 10^{-8} M MoAb SPV-T3a (▲), isotype-matched irrelevant MoAb MOPC-141 (●) or culture medium (■) for 24 hours at 37°C. Subsequently, cells were washed and cultured for another 72 hours in culture medium. CTL cytotoxicity was then assayed by specific lysis of a ^{51}Cr -loaded autologous EBV-LCL, and expressed as percentage relative to maximum lysis by detergent. Data represent the mean \pm SD of 3 experiments performed in triplicate.

Immunosuppressive activity of native MoAb SPV-T3a

SPV-T3a may deliver an immunosuppressive effect, irrespective of its conjugation to dgA, by modulation of the T-cell-receptor/CD3-complex (TCR/CD3-complex) or by induction of Fas-mediated apoptosis according to the process described as activation-induced cell death (AICD).¹⁶ To mimic the alloactivation induced in a transplantation setting, a CTL clone reactive with minor histocompatibility antigen 'EBV-peptide EBNA3C' was stimulated with an autologous EBV-LCL. Incubation of this CTL with MoAb SPV-T3a for 24 hours, abrogated almost completely its cytolytic activity as measured in a ^{51}Cr -release assay (Figure 2). Flow cytometric analysis (Figure 3) revealed that this may be attributed to both (1) AICD of the CTL-clone (to 16% of the untreated control) and (2) modulation of the CD3 antigen (to 20% of the untreated control). In addition, about half of the non-modulated CD3 antigens was still occupied by SPV-T3a.




Figure 3
AICD and CD3-modulation
by native SPV-T3a

The EBNA3C-reactive CTL clone was incubated with 10^{-8} M MoAb SPV-T3a, or isotype-matched control MoAb MOPC-141, for 24 hours at 37°C. Subsequently, cells were washed and cultured for another 72 hours in culture medium. The number of surviving cells was then determined by viability staining and flow cytometric analysis, and expressed as percentage relative to the untreated control (A). Membrane expression of free and SPV-T3a-bound CD3-antigen was determined as described in "Material and Methods" and indicated as percentage relative to control cells (B). Data represent the mean \pm SD of 3 experiments performed in triplicate.




Figure 4
Effect of IT treatment on NK activity

PBMC were treated with 10^{-8} M WT1-dgA (■), SPV-T3a-dgA (▲) or without IT (□), for 24 hours at 37°C. After treatment, cells were washed and cultured for an additional 3 days in culture medium without IT. Subsequently, remaining NK activity was determined by specific lysis of ^{51}Cr -labeled K562 blasts and expressed as percentage relative to untreated cells. During the experiment, 50 U/ml recombinant IL-2 was added to the culture medium to increase NK activity. Data represent the mean \pm SD of 3 experiments performed in triplicate.

NK activity after IT treatment

Although initiated by CTL, GVHD is thought to be aggravated by less specific cytokine-stimulated “bystander cells” like NK cells.^{3,17,18} The capacity of the described MoAb or IT to reduce NK activity was assayed by inhibition of lysis of ^{51}Cr -loaded K562 cells. Figure 4 shows the NK activity of PBMC incubated with IT for 24 hours. As expected, treatment with SPV-T3a-dgA did not effect the NK-activity (NK cells being CD3⁺). In contrast, NK activity was almost completely abolished following 3 days after incubation with WT1-dgA. Similar results were obtained with cell line Daudi, which is predominantly vulnerable for lymphokine-activated killer cells (LAK cells; data not shown). Neither nonconjugated MoAb WT1, nor isotype-matched control IT UPC-10-dgA, impaired the lysis of K562 or Daudi cells (data not shown, n = 3).

Table 2 *Patient characteristics*

	Patient No.			
	1	2	3	4
Age /Sex	65/M	34/M	47/M	49/M
Primary disease ^a	MM	CML	MDS	MM
Primary treatment ^b	- M, P	- H, a-I	- none	- V, A, D
Transplant procedure ^c	- I, C, TBI, PSCT	- ATG, C, TBI, BMT	- I, C, TBI, BMT	- C, TBI, BMT, DLI
Staging of GVHD ^d	Liver: 3 Gut: 1 Skin: 4 Overall: IV	Liver: 0 Gut: 0 Skin: 4 Overall: III	Liver: 0 Gut: 2 Skin: 0 Overall: II	Liver: 4 Gut: 0 Skin: 0 Overall: IV
Previous GVHD	C, Pl, S	C, Pl, S	C, Pl, S	C, Pl, S
Medication ^e				
IT-combination	2, 2, 4 mg/m ²	2, 2, 4, 4 mg/m ²	4, 4, 4, 4, mg/m ²	4, 4, 4 mg/m ²

a MM indicates multiple myeloma; CML: chronic myeloid leukemia; MDS: myelodysplastic syndrome.
b M indicates melphalan; P: prednisone; H: Hydrea, a-I: α -interferon; V: vincristine; A: Adriamycine; D: dexamethason.
c I: indicates idarubicine; C: cyclophosphamide; TBI: total body irradiation; PSCT: allogeneic peripheral stem cell transplantation; ATG: antithymocyte globuline; BMT: allogeneic bone marrow transplantation; DLI: donor lymphocyte infusion.
d Each organ system was staged grade 1 through 4 aGVHD according to Glucksberg et al.¹⁴ Patients were also given an overall grade of aGVHD (I through IV) based on severity of organ involvement.¹⁴
e C indicates cyclosporine A; Pl: prednisolone; S: Solu-medrol (up to 1g daily for at least 3 days).

Pilot-study Participants

So far, 4 patients, all white men, have been enrolled in the study. The clinical features of these patients are summarized in Table 2. Patients 1 and 2 have been treated at the lowest dose level ($2 \times 2 \text{ mg/m}^2$ followed by $2 \times 4 \text{ mg/m}^2$) and patient 3 and 4 at the second dose level ($4 \times 4 \text{ mg/m}^2$). Due to their early deaths, patient 1 and 4 received only 3 of the 4 planned infusions.

Table 3 Pharmacokinetic parameters determined over the complete treatment course

Patient No. (dose, mg/m ²)	IT	T _{1/2} (h)	AUC (μg·h/ml)	Cl (L/h)	V _d (L)	C _{max} (ng/ml)
1 (2, 2, 4)	SPV-T3a-dgA	3.4 ± 0.6	5.6 ± 0.6	0.36 ± 0.04	1.8 ± 0.2	1,830
	WT1-dgA	5.5 ± 1.2	5.3 ± 0.8	0.37 ± 0.06	3.0 ± 0.3	1,380
	IT-Combination	4.0 ± 0.7	10.5 ± 1.3	0.19 ± 0.02	1.1 ± 0.1	3,210
2 (2, 2, 4, 4)	SPV-T3a-dgA	6.5 ± 1.0	3.4 ± 0.3	0.59 ± 0.06	5.6 ± 0.5	720
	WT1-dgA	6.7 ± 1.4	3.9 ± 0.5	0.5 ± 0.07	5.0 ± 0.6	930
	IT-Combination	6.6 ± 1.2	7.3 ± 0.8	0.28 ± 0.03	2.6 ± 0.3	1,650
3 (4, 4, 4, 4)	SPV-T3a-dgA	6.9 ± 0.8	9.5 ± 0.7	0.64 ± 0.05	6.4 ± 0.4	675
	WT1-dgA	9.1 ± 1.4	3.4 ± 0.3	0.58 ± 0.06	7.6 ± 0.6	545
	IT-Combination	6.9 ± 0.8	12.4 ± 0.9	0.32 ± 0.02	3.2 ± 0.2	1,220
4 (4, 4, 4)	SPV-T3a-dgA	9.2 ± 1.5	12.2 ± 1.1	0.32 ± 0.03	4.4 ± 0.4	910
	WT1-dgA	8.8 ± 2.7	7.5 ± 1.3	0.54 ± 0.09	6.8 ± 1.2	668
	IT-Combination	9.1 ± 1.9	19.7 ± 2.3	0.20 ± 0.02	2.7 ± 0.3	1,560

T_{1/2}, plasma half-life; AUC, area under the concentration versus time curve; Cl, clearance; V_d, volume of distribution; and C_{max}, maximum plasma concentration.

Pharmacokinetics

The plasma clearance curves best fitted a 1-compartment model with a constant rate infusion and a first-order elimination rate for both SPV-T3a-dgA and WT1-dgA individually and given in combination. The pharmacokinetic parameters determined over the entire courses are listed in Table 3. The mean T_{1/2} was 6.5 ± 2.4 hours for SPV-T3a-dgA, 7.5 ± 1.7 hours for WT1-dgA, and 6.7 ± 2.1 hours for the IT-combination. Peak plasma levels were attained directly following each infusion and decreased (nearly) to baseline level in about 48 hours (Figure 5). The maximum plasma concentration (C_{max}) for the IT-combination ranged from 258 ng/ml following the first dose of patient 2 (2 mg/m^2) to 3,210 ng/ml following the third dose of patient 1 (4 mg/m^2). The latter seemed to be an exception as the maximum peak plasma level of the other patients fell within the relative narrow range of 1,220 to 1,650 ng/ml (all attained after a 4 mg/m^2 dose). The 2-fold higher level of patient 1 may be explained by his aggravating multi-organ failure interfering with the IT plasma clearance. When comparing the separate infusions, the mean T_{1/2} of the IT-combination almost doubled over the complete treatment course (Table 4). This may be explained by a reduction of available target antigens, which act as an antigen sink especially during the first infusion(s).




Figure 5 Plasma clearance curves

IT-combination plasma clearance curves are shown from patient 1 (treated with 2, 2, and 4 mg/m²) (○); patient 2 (2, 2, 4, and 4 mg/m²) (□); patient 3 (4, 4, 4, and 4 mg/m²) (●); and patient 4 (4, 4, and 4 mg/m²) (×). IT-combination plasma concentrations were deduced by summation of the individual values as determined for SPV-T3a-dgA and WT1-dgA.

Safety

The adverse events noted during the first 6 weeks following initiation of therapy are listed in Table 5. Because of the advanced disease of the patients, the etiology of the observed clinical events was obscured by multiple medical complications and concomitant medications. Patient 1 and 4 died during therapy due to worsening of complications already existing before start of the treatment, patient 1 from progressive multiorgan failure, patient 4 to generalized aspergillosis in combination with a cytomegalovirus (CMV) infection. Those symptoms thought to be related to the IT-combination were mild and transient. Patient 1 demonstrated edema in his shoulder not associated with weight gain, potentially due to limited capillary leakage. Patient 3 experienced episodes of fever during and between the IT-combination infusions. This fever may have been caused by a viral infection as well (see biological responses). Patient 4 demonstrated a rise of plasma CK-levels (not accompanied by myalgia) to 280 U/l on day 9 of therapy, being 1.45 times the upper limit of the normal range for men.

Table 4 The $T_{1/2}$ (mean \pm SD) determined over the separate infusions

	Infusion No.			
	1 (n=4)	2 (n=4)	3 (n=4)	4 (n=2)
SPV-T3a-dgA	4.5 \pm 0.8	5.4 \pm 0.7	6.5 \pm 2.8	6.2 \pm 1.4
WT1-dgA	3.2 \pm 1.5	8.3 \pm 4.8	7.9 \pm 2.0	9.0 \pm 3.3
IT-Combination	3.8 \pm 1.3	5.7 \pm 1.5	7.1 \pm 2.2	7.1 \pm 0.4

Table 5 *Adverse Events observed during the first 6 weeks of therapy^a*

	Total no. Grades 3/4 adverse events	All adverse events likely related to IT-combination
Allergy/Immunology	0	1 ^b
Auditory/Hearing	0	0
Blood/bone marrow	0	0
Cardiovascular	0	1 ^c
Coagulation	1 ^d	0
Constitutional symptoms	2 ^e	1 ^b
Dermatology/skin	0	0
Endocrine	0	0
Gastrointestinal	0	0
Hemorrhage	0	0
Hepatic	1 ^f	0
Infection/febrile neutropenia	2 ^g	0
Lymphatics	0	0
Metabolic/Laboratory	0	0
Musculoskeletal	0	1 ^h
Neurology	1 ⁱ	0
Ocular/visual	0	0
Pain	0	0
Pulmonary	0	0
Renal/Genitorurinary	0	0
Sexual/reproductive	0	0
Syndromes	0	0

a CTC-NCI criteria (version 2.0), using the bone marrow transplantation-specific scaling when applicable.

b Episodes of fever during and between infusions

c Limited capillary leakage

d TTP

e 40% decrease Karnofsky (n=2)

f Transient rise of SGOT & SGPT (after day 24)

g Aspergillosis, CMV-infection

h Rise in CK-levels to 1.45 times upper normal value

i Transient aphasia (<1 hour), probably related to TTP

The increase was not accompanied by a rise in the heart muscle isomer creatine kinase MB (CK-MB). Patient 4 had aphasia of short duration (<1 hour) 1 day after the first infusion of the IT-combination. At that time he was taking cyclosporine and had hemolysis with high reticulocyte counts, progressive thrombocytopenia, and fragmented red cells in the blood smear. It was concluded that aphasia was associated with cyclosporin-induced thrombotic thrombocytopenic purpura (TTP) and was not caused by the IT-combination (aphasia has been reported as a side-effect of other immunotoxins¹⁹⁻²¹). Cyclosporine was withdrawn and aphasia did not recur despite the further administration of the IT-combination.

Immunogenicity

Patient plasma samples were measured frequently before, during and after the trial for human-anti-mouse and human-anti-ricin antibodies (HAMA and HARA, respectively). No antibodies against SPV-T3a nor WT1 were detected in any of the cases. Patient 4

demonstrated a weakly positive HARA titer, 8 times the pretreatment value, on day 8 (one day before his death).

Biological Responses

Biological responses were monitored by flow cytometric evaluation of circulating T cells and/or NK cells. The concentrations of NK cells were too low to enable a reliable quantification with NK-specific markers. Therefore, NK cells and T cells were identified simultaneously using a mixture of a fluorescent-labeled CD2 (binding T cells and NK cells) and a CD5 MoAb (binding T cells). Figure 6 shows the amount of circulating T/NK cells expressed as percentage relative to the concentration at start of therapy. Patient 2 could not be adequately monitored due to the low initial number of circulating T/NK cells (about $6 \times 10^7/l$). All 3 evaluable patients already demonstrated a remarkable reduction of T/NK cells during the first administration of IT-combination. Immediately after the 4 hours of infusion, the number of circulating T/NK cells of patient 1, 3, and 4 had dropped to 17%, 8% and 24% of the pretreatment level, respectively. For patient 1 and 4, this number gradually declined to 1 to 3 % after the third infusion. Patient 3, in contrast, showed signs of a rebound of NK/T cells between the infusions. After the last infusion, a progressive expansion of T/NK cells was observed resulting in




Figure 6 Number of T/NK cells after IT-combination treatment

Patient 1 (\blacktriangle) and 4 (\circ) received 3, and patient 3 (\blacksquare) received all of the 4 planned IT-combination doses, given as 4-hour infusions at 48-hour intervals. The arrow beneath the x axis indicates the start of the first infusion. T/NK cells were identified by being CD2⁺ and/or CD5⁺. Their number was expressed as percentage relative to the concentration at start of therapy (being 0.7 , 1.0 , and 0.2×10^9 cells/l for patient 1, 3, and 4, respectively). The inset figure represents the amount of T/NK cells of patient 3 as determined over a longer period.

a peak concentration of about 45 times the pretreatment value at day 14. Flow cytometric analysis with CD4/CD8 MoAb and a T-cell receptor V β -panel pointed out that these cells were virtually all CD8⁺ and oligoclonal of origin. At day 18, the number of T cells had decreased again to about 5 times the pretreatment level (probably by apoptosis as suggested by annexin staining, data not shown). These observations, combined with the episodes of fever patient 3 experienced during the first week, are suggestive for a T-cell response after a viral infection.

The slow B-cell repopulation following stem cell transplantation prevented a reliable quantification of circulating B cells in patients 1, 2, and 3. The concentration of peripheral blood B cells of patient 4 showed a normal biological fluctuation ($0.1\text{--}0.2 \times 10^9/\text{l}$), which was not affected by the IT-combination. Moreover, the complete blood counts and leukocyte differential of all patients revealed no toxicity towards any of the other hematopoietic lineages (data not shown), thereby illustrating the selectivity of treatment for T/NK cells.

Skin biopsy specimens from patient 2 clearly demonstrated the biological efficacy of the IT-combination treatment (Figure 7). The microscopic appearance before treatment is typical for severe acute GVHD, with scattered infiltration of lymphocytes and a severely affected epidermal basal layer. After treatment, no remaining signs of lymphocyte infiltration could be detected, and the basal epidermis had regained its natural well-organized configuration. The intestinal GVHD of patient 3 was confirmed by a biopsy from the colon performed at day -5 (grade 3). When repeated at day 31, biopsy demonstrated a marked reduction of lymphocyte infiltrates without further signs of active GVHD. In patient 4 only a post mortem liver sample could be taken. Microscopic analysis revealed extensive cholestasis. Notably, no infiltrating lymphocytes could be detected.

Clinical Responses

Patient 1 showed a slight but clear improvement of his skin, from grade 4 to 3, starting at day 4. His (abnormal) liver functions remained stable; his gastrointestinal tract could not be objectively monitored due to administration of morphine. After 7 days of therapy, he died as a consequence of multiorgan failure. Patient 2 showed a dramatic reduction of his grade 4 GVHD of the skin to grade 0/1, starting 4 days after the first infusion and lasting for about 1.5 month. He then developed a modest relapse, GVHD grade 2 of the skin, which responded well to standard therapy of relatively low-dose corticosteroids supplemented with UV-B radiation. Patient 2 died 8 months after the IT-combination administration due to generalized aspergillosis and toxoplasmic encephalitis. Patient 3 showed strong improvement of his intestinal GVHD, grade 2 to grade 0, within 7 days after initiation of therapy. Apart from a mild and controllable GVHD reaction of his skin, grade 1 to 2 responding on low-dose corticosteroids, he showed no further signs of active GVHD. Two months after initiation of therapy, patient




Figure 7 Skin biopsies specimens

Skin biopsies were obtained from patient 2 taken just before (A), and two weeks after (B) treatment with the IT-combination. The appearance of the skin before treatment is typical for severe GVHD. A scattered infiltration of lymphocytes can be observed, localized primarily around the blood vessels and the junctional region of dermis (lower side) and epidermis (upper side). The epidermal basal layer is destroyed and the epidermis is disrupted from the basal lamina. Following treatment, no remaining signs of lymphocyte infiltrations were detectable. The basal layer had regained its natural well-organized configuration (original magnification $\times 100$, hematoxylin-eosin)

3 died of a bacterial infection. Patient 4 showed a more or less stable GVHD of his liver during the 7 days of treatment. He died due to a generalized aspergillosis in combination with a CMV infection on day 8.

DISCUSSION

The *in vitro* efficacy data demonstrate that simultaneous application of SPV-T3a-dgA and WT1-dgA results in a synergistic cytotoxicity, leading to an approximately 99% elimination of activated T cells. A comparable synergism has been described for other IT combinations as well.^{15,22-28} The most obvious advantage above single IT-treatment is that fewer target cells will be negative for multiple antigens than for a single antigen. In addition, the cells that express substantial levels of multiple target antigens may be loaded with more IT molecules. When the respective IT follow a different intracellular routing, the chance to escape the cytotoxic activity of the IT may be further reduced. With respect to anti-T-cell IT, reports addressing the combination approach are thus far focused on *in vitro* applications, including the purging of bone marrow grafts.^{15,22-24} In this report, we state that the combination of SPV-T3a-dgA and WT1-dgA appears appropriate for the *in vivo* elimination of unwanted T cells as well. This particular combination affords important benefits that surpass the “common synergism” as observed with the combinations of anti-T cell IT described so far.

SPV-T3a was selected as CD3 MoAb based on its IgG2b-isotype which strongly reduces the risk of cytokine release syndrome.^{29,30} The presence of a CD3-IT provides instant immunosuppression independent of dgA-based cytotoxicity (Figures 2 and 3). This may be of vital importance *in vivo* during treatment of acute life-threatening situations such as refractory GVHD. The limited (AICD) as well as temporal nature (blocking and modulation of CD3/TCR) of the underlying mechanisms provide arguments for making a “real killer” of SPV-T3a by conjugating it to dgA. The inclusion of the CD7-IT is essential, apart from the above-mentioned synergism, because it makes NK cells a target for the IT-combination as well (Figure 4). Accumulating evidence points at a distinctive role of NK cells in the pathophysiology of GVHD.^{17,18}

The pioneer work of Vitetta and colleagues was of great value in designing the clinical pilot-study. They thoroughly studied the *in vivo* efficacy of RFB4-dgA (CD22) and HD37-dgA (CD19) in treating patients with non-Hodgkin Lymphoma in relation to different dose regimens.^{13,20,21,31-33} In the present study, four patients with refractory acute GVHD have been treated with individual doses of 2 or 4 mg/m² IT-combination, according to their intermittent bolus infusion protocol. The mean T_{1/2} of SPV-T3a-dgA and WT1-dgA resembled those of RFB4-dgA and RFB5-dgA (6.1-7.8 hours), the latter

being a CD25-IT for the treatment of Hodgkin lymphoma.³⁴ The blood clearance of these IT is more than twice as fast than reported for HD37-dgA ($T_{1/2}$ of 18.2 hours). As with RFB4-dgA and RFB5-dgA, the clearance appeared to be influenced by circulating target cells. In contrast to RFB4-dgA and RFB5-dgA, this did not result in large inter-patient variations. Instead, the $T_{1/2}$ of the IT-combination increased about 2-fold during the entire treatment course, probably due to (1) the rapid elimination of “MoAb-capturing” target cells (Figure 6), and/or (2) by a reduction of available binding sites per cell (modulation/occupation). The latter possibility especially holds true for CD3 (see also Figure 3b). The CD7-antigen, in contrast, demonstrates a rapid re-expression of free antigen, resulting in the restoration of pre-treatment expression levels within 48 hours (*in vitro* and *in vivo* observations, data not shown). Three patients suffered from severe GVHD that was more or less restricted to the skin, gastrointestinal tract, and liver, respectively. Their relative narrow-ranged $T_{1/2}$ and C_{max} demonstrates that the type of organ involved has no major impact on the pharmacokinetics. The single exception was the patient suffering from a generalized GVHD (affecting skin, liver and gastrointestinal tract) in combination with a severe kidney failure. His about 2-fold higher C_{max} may reflect a different IT handling due to his worsening multiorgan failure.

In three patients, C_{max} exceeded 1.2 to 1.6 times and in 1 patient C_{max} exceeded 3.5 times the “clinically save threshold level” as defined for RFB4-dgA and HD37-dgA (1 $\mu\text{g/ml}$).^{13,20,32} Drug-related toxicities appeared to be restricted to limited capillary leakage and a modest rise of CK-levels (both $n = 1$). This may be the result of the IT-combination encompassing two distinct IT, each having a different spectrum of side effects (resulting in a distribution and, thereby, reduction in severity of drug-related toxicities). This is in agreement with observations of Vallera *et al.*³⁵ and Onda *et al.*³⁶, who both reported on comparable IT (directed against the same target cell/antigen and containing the same toxin), that displayed different toxicity profiles when tested in the same mouse model. None of the 4 patients treated with the IT-combination developed detectable HAMA. Only 1 patient showed a positive HARA titer (8 times baseline level). Apart from the small population size, this may be attributed to the huge dosages of corticosteroids administered, and to incomplete reconstitution of the B cell compartment early after transplantation.

The design of the pilot study does not allow an estimation of long-term toxicities (*e.g.* increase in opportunistic infections or secondary malignancies). Three of the 4 patients treated so far eventually died of an infection. The IT-combination can not be excluded as a contributing factor to the underlying immunosuppression. However, a realistic long-term safety profile can only be obtained when the IT-combination is applied without extensive preceding and concomitant second-line medication. At least theoretically, based on its short $T_{1/2}$ and narrow specificity range, the IT-combination is expected to accomplish a relatively limited immunosuppression by only temporarily affecting the T/NK cell compartment. Moreover, the composition of the IT-combination is better

defined and expected to vary less than that of ATG/ALG³⁷⁻³⁹, which use is often associated with sustained T cell depletion.^{40,41} From this viewpoint, it appears encouraging that the third patient showed a strong oligoclonal T-cell expansion directly after treatment with the IT-combination. The presumed antiviral T-cell response was not associated with a relapse of GVHD. This suggests a strong immunogenicity of the respective antigen and/or some form of treatment selectivity towards activated cells. The latter is in agreement with the *in vitro* observations and might be explained by the fact that on activation T cells show an increased expression of the CD7 target-antigen^{9,10} and become vulnerable for CD3-triggered AICD.^{42,43}

All patients showed a response, ranging from the elimination of the majority of circulating T/NK cells, to the dramatic clinical responses as observed in the patients 2 and 3. The biphasic pattern of T/NK cell removal indicates that multiple mechanisms are involved, some of which have not been observed *in vitro*. As for the *in vitro* efficacy data, the IT-combination needs several days before it displays its full killing capacity. This is in accordance with observations of Beyers *et al.*⁵, who reported that it took 5 daily infusions of H65-ricin A (CD5) to achieve a more or less gradual reduction of T cells to 20% of the initial number. In contrast, the IT-combination induced *in vivo* a massive and rapid reduction of T/NK cells already during the first four hours of infusion. Comparable biological responses have been described for native murine MoAb⁴⁴⁻⁴⁶, including anti-CD3/TCR MoAb of IgG2b isotype.^{47,48} The underlying mechanism is not fully clear yet. Some indirect evidence is provided that MoAb SPV-T3a is responsible for this initial rapid T-cell reduction. Tax and colleagues (unpublished observation, W.J.M.T.) demonstrated that nonconjugated MoAb WT1 did not influence the number of circulating T cells when administered *in vivo* to renal patients for the treatment of graft rejection. In addition, flow cytometric analysis of blood samples of the first patient demonstrated that the minor population of CD3⁺CD7⁺ cells (mostly NK cells) was reduced at a much slower rate (up to 2 days) than the CD3⁺CD7⁺ T cells (data not shown). In general, reduction of lymphocytes by native murine MoAb is often transient, a few days or less. In the present study, the initial rapid elimination was followed by a more gradual but sustained reduction of T/NK cells. This suggests a dgA-based elimination of target cells, which is supported by the *in vitro* killing capacity of the IT-combination observed at concentrations obtained *in vivo*.

It is impossible from the clinical data to determine the exact contribution of SPV-T3a-dgA and WT1-dgA (or their MoAb-moieties) to the observed biological and clinical responses. Unfortunately, no animal models are available that allow an efficacy evaluation of the individual components (SPV-T3a does not bind monkey-CD3). The high tolerability of the IT-combination argues against the clinical testing of theoretically less effective individual components (being single IT or MoAb). Instead, further studies should be focussed on elucidating the full potency of the IT-combination in its present form. Apart from dose optimization and application in an earlier phase of GVHD, this

may include the treatment of alternative indications that potentially benefit from the *in vivo* elimination of T cells.

The authors are aware of the limited nature of the clinical data obtained so far. However, the major finding that emerged from these data is that the IT-combination concept might work. Substantial biological as well as clinical responses have been observed in the absence of acute severe toxicities. This is especially meaningful considering the extensive pretreatment of the enrolled patients. Ideally, further studies will point out that the IT-combination forms a safe and effective tool for helping control certain diseases mediated by T cells.

ACKNOWLEDGMENTS

We are grateful to Dr. Harry Dolstra who generously provided the EBNA3C-reactive CTL-cell clone, to Dr. Elly van de Wiel-van Kemenade for helpful advice throughout the project, and to Arie Pennings, Cathy Maass, Mary Leenders and Jacky Ross for technical assistance.


REFERENCES

1. Horowitz MM, Gale RP, Sondel PM, Goldman JM, Kersey J, Kolb HJ, Rimm AA, Ringden O, Rozman C, Speck B, et al.: Graft-versus-leukemia reactions after bone marrow transplantation. *Blood* 75:555-62, 1990
2. Ferrara JL, Deeg HJ: Graft-versus-host disease. *N Engl J Med* 324:667-74, 1991
3. Antin JH, Ferrara JL: Cytokine dysregulation and acute graft-versus-host disease. *Blood* 80:2964-8, 1992
4. Kernan NA, Byers V, Scannon PJ, Mischak RP, Brochstein J, Flomenberg N, Dupont B, O'Reilly RJ: Treatment of steroid-resistant acute graft-vs-host disease by in vivo administration of an anti-T-cell ricin A chain immunotoxin. *Jama* 259:3154-7, 1988
5. Byers VS, Henslee PJ, Kernan NA, Blazar BR, Gingrich R, Phillips GL, LeMaistre CF, Gilliland G, Antin JH, Martin P, et al.: Use of an anti-pan T-lymphocyte ricin A chain immunotoxin in steroid-resistant acute graft-versus-host disease. *Blood* 75:1426-32, 1990
6. Hings IM, Severson R, Filipovich AH, Blazar BR, Kersey JH, Ramsay NK, McGlave PB, Weisdorf DJ: Treatment of moderate and severe acute GVHD after allogeneic bone marrow transplantation. *Transplantation* 58:437-42, 1994
7. Martin PJ, Nelson BJ, Appelbaum FR, Anasetti C, Deeg HJ, Hansen JA, McDonald GB, Nash RA, Sullivan KM, Witherspoon RP, Scannon PJ, Friedmann N, Storb R: Evaluation of a CD5-specific immunotoxin for treatment of acute graft- versus-host disease after allogeneic marrow transplantation. *Blood* 88:824-30, 1996
8. Spits H, Keizer G, Borst J, Terhorst C, Hekman A, de Vries JE: Characterization of monoclonal antibodies against cell surface molecules associated with cytotoxic activity of natural and activated killer cells and cloned CTL lines. *Hybridoma* 2:423-37, 1983
9. Tax WJ, Greaves MF, Willems HM, Leeuwenberg HF, Capel PJ, Koene RA: WT1: a monoclonal antibody reactive with T-ALL but not with other leukemias. *Hamatol Bluttransfus* 28:139-41, 1983
10. Tax WJ, Tidman N, Janossy G, Trejdosiewicz L, Willems R, Leeuwenberg J, De Witte TJ, Capel PJ, Koene RA: Monoclonal antibody (WT 1) directed against a T cell surface glycoprotein: characteristics and immunosuppressive activity. *Clin Exp Immunol* 55:427-36, 1984
11. Ghetie V, Thorpe P, Ghetie MA, Knowles P, Uhr JW, Vitetta ES: The GLP large scale preparation of immunotoxins containing deglycosylated ricin A chain and a hindered disulfide bond. *J Immunol Methods* 142:223-30, 1991
12. Burrows SR, Misko IS, Sculley TB, Schmidt C, Moss DJ: An Epstein-Barr virus-specific cytotoxic T-cell epitope present on A- and B-type transformants. *J Virol* 64:3974-6, 1990
13. Amlot PL, Stone MJ, Cunningham D, Fay J, Newman J, Collins R, May R, McCarthy M, Richardson J, Ghetie V, et al.: A phase I study of an anti-CD22-deglycosylated ricin A chain immunotoxin in the treatment of B-cell lymphomas resistant to conventional therapy. *Blood* 82:2624-33, 1993
14. Glucksberg H, Storb R, Fefer A, Buckner CD, Neiman PE, Clift RA, Lerner KG, Thomas ED: Clinical manifestations of graft-versus-host disease in human recipients of marrow from HL-A-matched sibling donors. *Transplantation* 18:295-304, 1974
15. Preijers FW, Tax WJ, Wessels JM, Capel PJ, De Witte T, Haanen C: Different susceptibilities of normal T cells and T cell lines to immunotoxins. *Scand J Immunol* 27:533-40, 1988
16. Kabelitz D, Pohl T, Pechhold K: Activation-induced cell death (apoptosis) of mature peripheral T lymphocytes. *Immunol Today* 14:38-9, 1993
17. Rhoades JL, Cibull ML, Thompson JS, Henslee-Downey PJ, Jennings CD, Sinn HP, Brown SA, Eichhorn TR, Cave ML, Jezek DA: Role of natural killer cells in the pathogenesis of human acute graft- versus-host disease. *Transplantation* 56:113-20, 1993

18. Hill GR, Krenger W, Ferrara JL: The role of cytokines in acute graft-versus-host disease. *Cytokines Cell Mol Ther* 3:257-66, 1997
19. Frankel AE, Laver JH, Willingham MC, Burns LJ, Kersey JH, Vallera DA: Therapy of patients with T-cell lymphomas and leukemias using an anti-CD7 monoclonal antibody-ricin A chain immunotoxin. *Leuk Lymphoma* 26:287-98, 1997
20. Stone MJ, Sausville EA, Fay JW, Headlee D, Collins RH, Figg WD, Stetler-Stevenson M, Jain V, Jaffe ES, Solomon D, Lush RM, Senderowicz A, Ghetie V, Schindler J, Uhr JW, Vitetta ES: A phase I study of bolus versus continuous infusion of the anti-CD19 immunotoxin, IgG-HD37-dgA, in patients with B-cell lymphoma. *Blood* 88:1188-97, 1996
21. Vitetta ES, Stone M, Amlot P, Fay J, May R, Till M, Newman J, Clark P, Collins R, Cunningham D, et al.: Phase I immunotoxin trial in patients with B-cell lymphoma. *Cancer Res* 51:4052-8, 1991
22. Vallera DA, Ash RC, Zanjani ED, Kersey JH, LeBien TW, Beverley PC, Neville DM, Jr., Youle RJ: Anti-T-cell reagents for human bone marrow transplantation: ricin linked to three monoclonal antibodies. *Science* 222:512-5, 1983
23. Uckun FM, Azemove SM, Myers DE, Vallera DA: Anti-CD2 (T, p50) intact ricin immunotoxins for GVHD-prophylaxis in allogeneic bone marrow transplantation. *Leuk Res* 10:145-53, 1986
24. Katz FE, Janossy G, Cumber A, Ross W, Blacklock HA, Tax W, Thorpe PE: Elimination of T cells from human peripheral blood and bone marrow using a cocktail of three anti-T cell immunotoxins. *Br J Haematol* 67:407-11, 1987
25. Crews JR, Maier LA, Yu YH, Hester S, O'Briant K, Leslie DS, DeSombre K, George SL, Boyer CM, Argon Y, et al.: A combination of two immunotoxins exerts synergistic cytotoxic activity against human breast-cancer cell lines. *Int J Cancer* 51:772-9, 1992
26. Ghetie MA, Tucker K, Richardson J, Uhr JW, Vitetta ES: The antitumor activity of an anti-CD22 immunotoxin in SCID mice with disseminated Daudi lymphoma is enhanced by either an anti-CD19 antibody or an anti-CD19 immunotoxin. *Blood* 80:2315-20, 1992
27. Engert A, Gottstein C, Bohlen H, Winkler U, Schon G, Manske O, Schnell R, Diehl V, Thorpe P: Cocktails of ricin A-chain immunotoxins against different antigens on Hodgkin and Sternberg-Reed cells have superior anti-tumor effects against H-RS cells in vitro and solid Hodgkin tumors in mice. *Int J Cancer* 63:304-9, 1995
28. Flavell DJ, Noss A, Pulford KA, Ling N, Flavell SU: Systemic therapy with 3BIT, a triple combination cocktail of anti-CD19, -CD22, and -CD38-saporin immunotoxins, is curative of human B-cell lymphoma in severe combined immunodeficient mice. *Cancer Res* 57:4824-9, 1997
29. Holtrop S, Rijke-Schilder GP, Koene RA, Tax WJ: A polymorphic Fc receptor for mouse IgG2b on human B cells and monocytes. *Immunology* 74:613-20, 1991
30. Frenken LA, Koene RA, Tax WJ: The role of antibody isotype in IFN-gamma and IL-2 production during anti-CD3-induced T cell proliferation. *Transplantation* 51:881-7, 1991
31. Conry RM, Khazaeli MB, Saleh MN, Ghetie V, Vitetta ES, Liu T, LoBuglio AF: Phase I trial of an anti-CD19 deglycosylated ricin A chain immunotoxin in non-Hodgkin's lymphoma: effect of an intensive schedule of administration. *J Immunother Emphasis Tumor Immunol* 18:231-41, 1995
32. Sausville EA, Headlee D, Stetler-Stevenson M, Jaffe ES, Solomon D, Figg WD, Herdt J, Kopp WC, Rager H, Steinberg SM, et al.: Continuous infusion of the anti-CD22 immunotoxin IgG-RFB4-SMPT-dgA in patients with B-cell lymphoma: a phase I study. *Blood* 85:3457-65, 1995
33. Senderowicz AM, Vitetta E, Headlee D, Ghetie V, Uhr JW, Figg WD, Lush RM, Stetler-Stevenson M, Kershaw G, Kingma DW, Jaffe ES, Sausville EA: Complete sustained response of a refractory, post-transplantation, large B-cell lymphoma to an anti-CD22 immunotoxin. *Ann Intern Med* 126:882-5, 1997

34. Engert A, Diehl V, Schnell R, Radszuhn A, Hatwig MT, Drillich S, Schon G, Bohlen H, Tesch H, Hansmann ML, Barth S, Schindler J, Ghetie V, Uhr J, Vitetta E: A phase-I study of an anti-CD25 ricin A-chain immunotoxin (RFT5-SMPT- dgA) in patients with refractory Hodgkin's lymphoma. *Blood* 89:403-10, 1997
35. Valleria DA, Carroll SF, Snover DC, Carlson GJ, Blazar BR: Toxicity and efficacy of anti-T-cell ricin toxin A chain immunotoxins in a murine model of established graft-versus-host disease induced across the major histocompatibility barrier. *Blood* 77:182-94, 1991
36. Onda M, Kreitman RJ, Vasmatazis G, Lee B, Pastan I: Reduction of the nonspecific animal toxicity of anti-Tac(Fv)-PE38 by mutations in the framework regions of the Fv which lower the isoelectric point. *J Immunol* 163:6072-7, 1999
37. Greco B, Bielory L, Stephany D, Hsu SM, Gascon P, Nienhuis A, Young N: Antithymocyte globulin reacts with many normal human cell types. *Blood* 62:1047-54, 1983
38. Raefsky EL, Gascon P, Gratwohl A, Speck B, Young NS: Biological and immunological characterization of ATG and ALG. *Blood* 68:712-9, 1986
39. Rebollato LM, Gross U, Verbanac KM, Thomas JM: A comprehensive definition of the major antibody specificities in polyclonal rabbit antithymocyte globulin. *Transplantation* 57:685-94, 1994
40. Bell L, Girardin C, Sharma A, Goodyer P, Mazer B: Lymphocyte subsets during and after rabbit anti-thymocyte globulin induction in pediatric renal transplantation: sustained T cell depletion. *Transplant Proc* 29:6S-9S, 1997
41. Muller TF, Grebe SO, Neumann MC, Heymanns J, Radsak K, Sprenger H, Lange H: Persistent long-term changes in lymphocyte subsets induced by polyclonal antibodies [see comments]. *Transplantation* 64:1432-7, 1997
42. Wesselborg S, Janssen O, Kabelitz D: Induction of activation-driven death (apoptosis) in activated but not resting peripheral blood T cells. *J Immunol* 150:4338-45, 1993
43. Owen-Schaub LB, Yonehara S, Crump WLD, Grimm EA: DNA fragmentation and cell death is selectively triggered in activated human lymphocytes by Fas antigen engagement. *Cell Immunol* 140:197-205, 1992
44. Chatenoud L, Baudrihay MF, Kreis H, Goldstein G, Schindler J, Bach JF: Human in vivo antigenic modulation induced by the anti-T cell OKT3 monoclonal antibody. *Eur J Immunol* 12:979-82, 1982
45. Bertram JH, Gill PS, Levine AM, Boquiren D, Hoffman FM, Meyer P, Mitchell MS: Monoclonal antibody Tr01 in T cell malignancies: a clinical, pharmacokinetic, and immunologic correlation. *Blood* 68:752-61, 1986
46. Parlevliet KJ, ten Berge IJ, Yong SL, Surachno J, Wilmink JM, Schellekens PT: In vivo effects of IgA and IgG2a anti-CD3 isotype switch variants. *J Clin Invest* 93:2519-25, 1994
47. Anasetti C, Martin PJ, Storb R, Appelbaum FR, Beatty PG, Davis J, Doney K, Hill HF, Stewart P, Sullivan KM, et al.: Treatment of acute graft-versus-host disease with a nonmitogenic anti- CD3 monoclonal antibody. *Transplantation* 54:844-51, 1992
48. Land W, Hillebrand G, Illner WD, Abendroth D, Hancke E, Schleibner S, Hammer C, Racenberg J: First clinical experience with a new TCR/CD3-monoclonal antibody (BMA 031) in kidney transplant patients [letter]. *Transpl Int* 1:116-7, 1988

SUMMARY AND DISCUSSION



SUMMARY AND DISCUSSION

This thesis addresses the use of ITs in relation to hematological disorders. **Chapter 1** provides a general overview of the mechanisms of action and current status of ITs, whereas **Chapter 2** focuses on the potential applications in the field of hematology. The subsequent **Chapters 3 to 5** describe part of the institutional fundamental research performed to improve this form of therapy. In **Chapters 6 and 7**, the rationale as well as the 'in house' preparation and first clinical results of an IT-combination for GVHD are addressed. Hereinafter, **Chapters 3 to 7** are discussed in more detail followed by a general perspective regarding the current and future status of ITs.

Chapter 3 describes a comparison of the internalization and cytotoxicity of an anti-CD3 ricin A chain IT (WT32-RTA). The study is based on an improved method to determine the cellular uptake of radiolabeled MoAb/IT. Initially, internalization studies were solely relying on cell-associated radioactivity. The activity removable by acid or protease treatment formed a measure for membrane bound MoAb/IT, whereas non-removable activity represented the amount of internalized molecules. Though very useful in studying several IT-related phenomena, this method had a major shortcoming. Intracellularly processed (read: degraded) and subsequently exocytosed MoAb/IT was not taken into account. With regard to IT, however, these molecules could very well have contributed to the intoxication of the cell by delivering the toxin before being processed. When studying internalization, it is therefore important to measure the amount of processed and shed MoAb/IT-fragments in the culture supernatant as well. This was hampered, initially, by the presence of other low weight radioactive material in the starting material. Removal of this material was cumbersome, involving dialysis or repetitive dilution/concentration steps of the radioactive starting material. Some investigators circumvented this problem by preloading target cells with radiolabeled MoAb/IT at 4°C, and subsequently washing out small radioactive particles and unbound MoAb/IT. The actual internalization studies could then be performed at 37°C, with intracellularly

processed and shed MoAb/IT now being readily detectable as the low weight radioactive material emerging in the supernatant. A limitation of this strategy, however, was the absence of unbound MoAb/IT surrounding the target cells. Normally, this excess of MoAb/IT influences the MoAb-antigen binding equilibrium, and forms a 'sink' for the binding of re-expressed antigen. With the introduction of dispensable concentrators, the extensive removal of low weight radioactive material came within practical reach (to <0.3% of the total radioactivity). Consequently, internalization studies could be performed in the continuous presence of unbound MoAb/IT and more accurately reflected the therapeutic situation.

When thus applied to study the internalization of the CD3 MoAb/IT with leukemia T-cell line Jurkat, an ongoing biphasic cellular uptake of MoAb/IT could be demonstrated. This internalization was characterized by a rapid initial uptake, followed by slower but progressive internalization driven by the constant re-expression of CD3-antigen. Addition of cytotoxicity enhancers NH_4Cl or monensin retarded intercellular degradation, resulting in an increased number of intracellularly present IT/MoAb. This should increase the possibility of cytoplasmatic deliverance of the toxin-moiety and, thereby, lower the amount of internalized IT necessary for intoxication of the cell. These observations closely matched the kinetics of cytotoxicity as determined in a protein synthesis inhibition assay. Both prolongation of the incubation time and addition of NH_4Cl or monensin clearly increased the efficacy of the CD3-IT towards Jurkat cells. An interesting observation was that the IT-induced cytotoxicity was characterized by a lag period due to a delay in either the internalization or the subsequent processing of membrane bound IT. Translated to the *ex vivo* purging of bone marrow, the above results argue for a 24 h IT-incubation in the presence of either NH_4Cl or monensin to ensure an optimal purging efficacy, also to cells with low antigen density, without causing harm to normal progenitor cells.

Looking at a future role of internalization studies, they may be of help in determining optimal dose schedules for *in vivo* treatment. They could provide a rationale for selecting either a continuous or bolus infusion protocol, depending on their effect on internalization. When considering multiple bolus infusions, internalization studies may help to identify the optimal interval. Increasing the time between infusions, for instance, may render target cells more vulnerable for the subsequent dose by allowing a more complete re-expression of antigen.

Another example of the relevance of internalization studies is described in **Chapter 4**. This study addresses the striking discrepancy in cytotoxicity of two anti-CD19 ricin A-ITs differing in isotype only (IgG1 and IgG2a). Internalization experiments with both anti-CD19 MoAb, as well as with their F(ab')_2 fragments, revealed that the presence of the IgG1 Fc-part enhanced the cellular uptake. The involvement of $\text{Fc}\gamma\text{RII}$ (CD32) in this process was demonstrated by a decreased cytotoxicity of the IgG1-IT (and not IgG2a-

IT) in the presence of FcγRII-blocking MoAb. These results suggest that the IgG1 Fc-part of membrane bound anti-CD19 IT interacts with FcγRII, which supports the cellular uptake of the antigen-IT complex analogous to the mechanism described by Kurlander. Additional experiments identified functional FcγRIIa as the responsible FcγR-isoform. The tremendous increase in potency as compared to the relative modest rise in internalization (~100-fold and 3-fold, respectively), suggests that the enhanced uptake is accompanied by a different intracellular routing which favors ricin A toxicity. Alternatively, the increased number of intracellular molecules may exceed a critical threshold above which the cell is no longer capable of efficiently inactivating the IT. The relevance of this phenomenon for other ITs has to be determined yet. Additional experiments with an IgG1 anti-CD22 IT revealed no effect of FcγRII-blocking on cytotoxicity. It should be noted, though, that the intracellular routing of anti-CD22 ITs is more effective than that of anti-CD19 ITs (as indicated by a superior toxicity despite similar or lower internalization rates). Any favorable effect of the IgG1-isotype might therefore only marginally influence the already high efficacy of anti-CD22 ITs. Nevertheless, the above study forms a clear example of the potential importance of the MoAb isotype on IT efficacy. This will be even more apparent regarding *in vivo* applications, considering the potential interactions with the patient's immune system. As such, these data provide an argument against the routine exclusion of MoAb Fc-fragments from recombinant fusion toxins. Though being technically more convenient, Fv-based ITs lack the (potential) benefit of Fc-mediated effector mechanisms contributing to, or even synergizing, the toxin-based elimination of target cells.

Prior to clinical use, ITs have to be extensively screened for activity against target and non-target cells. **Chapter 5** describes a sensitive flow cytometric (FCM) method capable of determining IT-induced cell kill in freshly derived patient-samples. This method is based on the FCM determination of surviving target-cells following staining with viability markers propidium iodine (PI) and calceine (Calc). The exact quantification of surviving cells was enabled by admixture of a known concentration of inert beads prior to FCM-analysis. An important finding using this method was that IT-based inhibition of protein synthesis is not directly accompanied by the physical disruption of the cell membrane. A post-treatment period of several days was necessary to enable full exposure of the IT-induced cell kill, comparable to that as determined with clonogenic assays.

As demonstrated in Chapter 5, the described method enables the evaluation of target cell depletion in a heterogeneous population as blood or BM. The use of a cell-specific, fluorescence-labeled, MoAb is most straightforward, enabling live/death discrimination and target-cell identification in one analysis. For this approach to be useful, the target-cell needs to express a specific antigen besides the target-antigen already used for therapy. The latter is inappropriate as this antigen might be modulated or covered by the therapeutic MoAb/IT. Moreover, residual malignant cells may be enriched for cells

with low levels of target-antigen expression. When no appropriate ‘identification-antigen’ is present, target cells may be identified by PCR-amplification of a cell specific DNA-sequence. This requires the pre-sorting of the viable fraction to prevent the DNA of IT-eliminated cells from contributing to the PCR-signal. In practice, triple-sorted viable cells should contain less than one in 10^6 erroneously sorted non-viable cells, matching the detection limit of a sensitive PCR-assay.

The flowcytometric analysis of cell kill has now been applied in various studies evaluating the efficacy of varying IT towards different cell types, including freshly derived heterogeneous samples.^{1,2} These studies confirmed the relevance of directly measuring the number of surviving cells, and suggest this assay may prove a helpful tool in optimizing IT-therapy. As such, it forms a valuable addition to the conventional clonogenic and protein synthesis inhibition assays.

The **first part of Chapter 7** describes the experiments that provided the rationale for the development of an IT-based therapy for aGVHD. These experiments pointed out that anti-CD3 and anti-CD7 ricin A-IT (SPV-T3a-dgA and WT1-dgA, respectively), act synergistically in killing T cells. The most obvious advantage of using a combination of ITs is that fewer target cells will be negative for multiple antigens than for a single antigen. Moreover, cells expressing both target-antigens may be loaded with more IT molecules. Apart from this ‘common synergism’, the particular combination of SPV-T3a-dgA and WT1-dgA affords important additional benefits. The presence of a CD3-IT provides instant immunosuppression, which relies on the binding of the CD3/T-cell-receptor complex (CD3/TCR) and is independent of dgA-based toxicity. The underlying mechanisms, modulation of the CD3/TCR complex and activation induced cell death (AICD), may be of vital importance *in vivo* during treatment of acute life-threatening situations such as refractory GVHD. The anti-CD3 MoAb SPV-T3a was selected based on its ‘non-T-cell-activating’ IgG2b-isotype which strongly reduces the risk of cytokine release syndrome (associated with the use of ‘activating’ CD3 MoAb). Inclusion of the CD7-IT is essential, apart from the above mentioned ‘common synergism’, since it makes NK cells (largely CD3⁺/CD7⁺) a target for the IT-combination as well. Accumulating evidence points to a distinctive role of NK cells in the pathophysiology of aGVHD. Altogether, these results formed the base for the clinical evaluation of the IT-combination in patients with refractory aGVHD.

The conversion of laboratory experiments to the early clinical studies asks for a revolutionary change of the production process. Apart from the scale up of equipment and materials, the handling, purity and activity of the final product have to meet stricter requirements. It would be most straightforward to outsource such productions to authorized contract manufacturers with relevant experience. Unfortunately, the required financing often forms a major obstacle. Even early clinical studies are generally considered ‘too less fundamental’ to qualify entirely for regular scientific granting. Potential

pharmaceutical partners, on the other hand, often prefer clear signs of clinical efficacy before taking the obvious risk of investing in experimental drugs. To overcome this impasse, and supported by a grant of the government (technology foundation STW), it was decided to perform most of the production and validation 'in house'. **Chapter 6** describes this production of clinical grade material as performed at the Central Hematology Laboratory under supervision of the department of Clinical Pharmacy, and with the assistance, amongst others, of the departments of Pathology, Virology and the Central Animal Laboratory. To ensure the highest possible safety and functionality of the final product, previously a draft quality was formulated based on the FDA-document 'Points to consider in the manufacture and testing of monoclonal antibody products for human use'. Reports of the production of comparable ITs already tested in FDA-approved trials formed an additional guideline. The actual production provided sufficient material for the planned clinical pilot study. Yields were comparable to those previously reported in literature (>25% MoAb-recovery). The specifications of the final products mostly matched those as prescribed in the draft-quality. The few minor deviations, regarding the percentage unconjugated WT1 and the DNA-content of both WT1-dgA and SPV-T3a-dgA, were, retrospectively, considered to be within an acceptable range. Animal experiments revealed a murine LD₅₀ (lethal dose for half of the animals) equal, or higher, than that of similar ITs already in clinical use. The major concern formed the cross-reactivity of SPV-T3a with the human esophagus, observed during the *in vitro* screening of a human tissue panel. To evaluate the potential clinical relevance of this observation, human doses IT-combination were administered to two cynomolgus monkeys (sharing the *in vitro* cross-reactivity). Neither toxicity nor binding of SPV-T3a-dgA to the monkey esophagus could be detected. Notably, therapeutic serum concentrations (>1.8 µg/ml) were obtained for several hours in both monkeys without acute irreversible toxicities occurring. The main side effect was a transient rise of CK-levels, indicative for dgA-associated muscle injury. Altogether, the above data justified the evaluation of the produced material in a clinical pilot-study for the treatment of life-threatening aGVHD. As such, the 'in house' production can be considered an essential stepping stone towards the future large-scale productions required for the actual registration studies.

So far five patients have been treated with three to four infusions of 2 or 4 mg/m² IT-combination, administered intravenously at 48-hour intervals. The **second part of Chapter 7** describes the results obtained with the first four patients. The T_{1/2} was ~7 hours, with peak serum levels ranging from 258 to 3,210 ng/ml. The infusions were well tolerated. Drug-associated side effects were restricted to limited edema, fever, and a modest rise of creatine kinase levels, suggesting the maximal tolerated individual dose lies well above 4 mg/m². Remarkably, the IT-combination did not provoke vascular leakage, or other serious toxicities, while the peak serum concentrations in all patients either reached (n = 1) or exceeded (n = 4) the 'clinically save threshold' as defined for

comparable IT (being 1 µg/ml). An explanation for this may be that administration of two distinct ITs may result in a distribution and, thereby, dilution of individual side effects. This is in accordance with observations of others who reported on almost identical ITs (directed against the same target cell/antigen and containing the same toxin), which displayed different toxicity profiles when tested in the same mouse model. Moreover, the concomitant administration of high dose corticosteroids may have positively affected the maximal tolerable dose. Notably, only one of the patients treated so far developed low titer antibodies against ricin A. This may again be attributed to the high dose corticosteroids. Alternative explanations are the incomplete reconstitution of the B cell compartment early post-transplantation, and the successful depletion of accessory T cells. Biological responses could be noted in all five patients, as characterized by a rapid and extensive drop of circulating T/NK cells. The biphasic pattern of this depletion suggests the involvement of multiple mechanisms. The rapid depletion of T cells, already observed directly after infusion, appears to be MoAb-related. While the subsequent more gradual but sustained elimination of T/NK cells points to a ricin A-based mechanism. Noteworthy, one patient mounted an extensive CD8 T-cell response directly following treatment (indicative for an anti-viral immune response), not accompanied with aggravating GVHD. The other patient whose peripheral T cells could be reliably monitored for a longer period, demonstrated a gradual reconstitution of circulating T cells starting two weeks after initiation of therapy. The accompanying shift in the TCR-Vβ repertoire, suggested these were newly generated T cells. (patient No. 5, unpublished observations). Two of the five patients already died during the first treatment week, due to underlying diseases and pre-existing organ impairment. The three other patients demonstrated extensive clinical responses, despite unresponsiveness to the intensive pretreatment.

Though limited in nature, these preliminary data are suggestive for a clear therapeutic window where, in contrast, hardly one existed for the other dgA-based IT tested so far. This may be explained by the combination approach, which, on one hand, decreases the overall toxicity, and, on the other hand, increases effectiveness by appealing on multiple, synergistic, effector mechanisms. Notwithstanding the clear biological and clinical responses, all five patients eventually died due to transplantation related complications (mostly infections). These results are inherent to the type of study, only permitting the enrollment of patients with very advanced and therapy-resistant disease (not responding to cyclosporin plus 1g corticosteroids daily). The absence of severe side effects, now justifies the treatment of a limited number of patients with less advanced aGVHD. This enables the evaluation of the IT-combination in a more appropriate clinical setting. Moreover, it should predict the suitability of the IT-cocktail for alternative indications that may benefit from the temporary removal of T/NK cells. Most obvious is the treatment of graft-rejection (either stem cells or solid organs). In addition, animal studies clearly demonstrated that anti-T cell IT may have a role in creating the prop-

er immunologic environment for non-myeloablative stem cell transplantation, achieving tolerance induction, or even treating certain T-cell mediated auto-immune disorders.

General perspective

The introduction of a new therapeutic strategy often creates high expectations regarding its use for treating life-threatening diseases. These do not always reflect the inevitable hurdles generally accompanying the development cycle of new drugs. As such, ITs form no exception. Based on the initial reports, describing complete cures of lethally tumor-xenografted mice, they seemed the ultimate representation of Paul Ehrlich's magic bullet concept. The subsequent translation of ITs to the clinic, however, revealed major obstacles preventing instant therapeutic success in man. Several of the problems encountered resembled those previously seen in MoAb trials. Obvious examples form the occurrence of humoral responses, in this case also directed against the toxin-moiety, and the heterogeneous expression of target-antigens. Moreover, as with MoAbs, the Phase I/II testing in patients with advanced bulky disease is not considered optimal to expose the full potential of this form of treatment. Most importantly, though, the clinical studies revealed serious toxicities, not previously detected to the same extent in the test animals. Especially for ricin-based ITs, the occurrence of severe vascular leakage dramatically narrowed the therapeutic window. As a consequence, safely attainable serum concentrations appeared to be several-fold lower than those previously determined to be effective in mice.³ Altogether, these early clinical results tempered the initial hype, leaving only a limited number of old-time believers still trying to make the concept work. Basically, their approach was to transfer ITs back to appropriate animal models, and to solve the individual obstacles with the help of new exciting (recombinant-DNA) techniques coming available.⁴⁻⁶ As it looks now, there are some serious indications that their persistence will be rewarded.

With respect to reduction of the immunogenicity, various strategies are currently being explored. One is based on the conjugation of non-immunogenic human toxin equivalents to either human antibodies, growth factors or cytokines. This approach already resulted in the construction of several 'fully human' ITs with proven *in vitro* activity⁷⁻¹². Another strategy focuses on making the highly potent conventional toxins less immunogenic by genetically altering the most immunogenic determinants, or by pegylation of the molecule.^{13,14} Combined with routinely produced human MoAbs, these may render 'Stealth' ITs being (almost) invisible for the patient's immune system. A third approach is based on transient repressing immune reactions by the co-administration of immunomodulating agents.^{15,16} In order to anticipate on heterogeneous antigen expression, the simultaneous use of multiple ITs have been addressed. Several animal studies convincingly demonstrated the synergistic efficacy of a cocktail of ITs.¹⁷⁻¹⁹ Additionally, the first clinical results obtained with such combinations are now being

reported.^{2,20} Noteworthy, the IT-combination described in Chapters four and five underscored the potential significance of MoAb-based effector mechanisms contributing to the overall IT efficacy.² The superiority of the concomitant use of ITs with conventional chemo/radiation therapy has been clearly demonstrated in several animal experiments.²¹⁻²³ These encouraging results should facilitate the clinical evaluation of ITs in the more appropriate adjuvant setting. With regard to the dose limiting toxicities, the most important finding is probably the unraveling of the cause of vascular leakage. Baluna *et al.* of the group of Vitetta identified a sequence motif in the ricin A chain, present in other toxins and the VLS-inducing cytokine IL-2 as well, which is responsible for VLS by binding to and damaging endothelial cells.^{6,24-27} Future studies should point whether deletions or mutations in this sequence, or the use of nondamaging blocking peptides, may increase the therapeutic index of ITs prepared with the conventional plant or bacterial toxins.

Altogether, these recent developments will undoubtedly result in the production of next generation ITs with strongly improved *in vivo* characteristics. It appears that ITs are still very much alive and ready to fulfill at least some of their clinical promises in the foreseeable future. As such, the recent FDA-approval of a recombinant IT for the treatment of a refractory hematological malignancy (Ontak, DAB389IL-2), might be regarded as an important and indicative turning point.

REFERENCES

1. van Horssen PJ, van Oosterhout YV, Evers S, Backus HH, van Oijen MG, Bongaerts R, de Witte T, Preijers FW: Influence of cytotoxicity enhancers in combination with human serum on the activity of CD22-recombinant ricin A against B cell lines, chronic and acute lymphocytic leukemia cells. *Leukemia* 13:241-9, 1999
2. van Oosterhout YV, van Ernst L, Schattenberg AV, Tax WJ, Ruiter DJ, Spits H, Nagengast FM, Masereeuw R, Evers S, de Witte T, Preijers FW: A combination of anti-CD3 and anti-CD7 ricin A-immunotoxins for the *in vivo* treatment of acute graft-versus-host disease. *Blood* 95:3693-3701, 2000
3. Sausville EA: Use of immunotoxins in the treatment of lymphoma. *Cancer J Sci Am* 6:135-8, 2000
4. Valleria DA: Immunotoxins: will their clinical promise be fulfilled? [editorial]. *Blood* 83:309-17, 1994
5. Vitetta ES, Thorpe PE, Uhr JW: Immunotoxins: magic bullets or misguided missiles? *Immunol Today* 14:252-9, 1993
6. Vitetta ES: Immunotoxins and vascular leak syndrome. *Cancer J Sci Am* 6 Suppl 3:S218-24, 2000
7. Rybak SM, Hoogenboom HR, Meade HM, Raus JC, Schwartz D, Youle RJ: Humanization of immunotoxins. *Proc Natl Acad Sci U S A* 89:3165-9, 1992
8. Zewe M, Rybak SM, Dubel S, Coy JF, Welschof M, Newton DL, Little M: Cloning and cytotoxicity of a human pancreatic RNase immunofusion. *Immunotechnology* 3:127-36, 1997
9. Newton DL, Pollock D, DiTullio P, Echelard Y, Harvey M, Wilburn B, Williams J, Hoogenboom HR, Raus JC, Meade HM, Rybak SM: Antitransferrin receptor antibody-RNase fusion protein expressed in the mammary gland of transgenic mice. *J Immunol Methods* 231:159-67, 1999
10. Gho YS, Chae CB: Luteinizing hormone releasing hormone-RNase A conjugates specifically inhibit the proliferation of LHRH-receptor-positive human prostate and breast tumor cells. *Mol Cells* 9:31-6, 1999
11. Psarras K, Ueda M, Tanabe M, Kitajima M, Aiso S, Komatsu S, Seno M: Targeting activated lymphocytes with an entirely human immunotoxin analogue: human pancreatic RNase1-HUMAN IL-2 fusion. *Cytokine* 12:786-90, 2000
12. Linardou H, Epenetos AA, Deonarain MP: A recombinant cytotoxic chimera based on mammalian deoxyribonuclease-I. *Int J Cancer* 86:561-9, 2000
13. Roscoe DM, Pai LH, Pastan I: Identification of epitopes on a mutant form of *Pseudomonas* exotoxin using serum from humans treated with *Pseudomonas* exotoxin containing immunotoxins. *Eur J Immunol* 27:1459-68, 1997
14. Tsutsumi Y, Onda M, Nagata S, Lee B, Kreitman RJ, Pastan I: Site-specific chemical modification with polyethylene glycol of recombinant immunotoxin anti-Tac(Fv)-PE38 (LMB-2) improves antitumor activity and reduces animal toxicity and immunogenicity. *Proc Natl Acad Sci U S A* 97:8548-53, 2000
15. Gelber EE, Vitetta ES: Effect of immunosuppressive agents on the immunogenicity and efficacy of an immunotoxin in mice. *Clin Cancer Res* 4:1297-304, 1998
16. Siegall CB, Haggerty HG, Warner GL, Chace D, Mixan B, Linsley PS, Davidson T: Prevention of immunotoxin-induced immunogenicity by coadministration with CTLA4Ig enhances antitumor efficacy. *J Immunol* 159:5168-73, 1997
17. Ghetie MA, Tucker K, Richardson J, Uhr JW, Vitetta ES: The antitumor activity of an anti-CD22 immunotoxin in SCID mice with disseminated Daudi lymphoma is enhanced by either an anti-CD19 antibody or an anti-CD19 immunotoxin. *Blood* 80:2315-20, 1992
18. Flavell DJ, Noss A, Pulford KA, Ling N, Flavell SU: Systemic therapy with 3BIT, a triple combination cocktail of anti-CD19, -CD22, and -CD38-saporin immunotoxins, is curative of human B-cell lymphoma in severe combined immunodeficient mice. *Cancer Res* 57:4824-9, 1997

19. Engert A, Gottstein C, Bohlen H, Winkler U, Schon G, Manske O, Schnell R, Diehl V, Thorpe P: Cocktails of ricin A-chain immunotoxins against different antigens on Hodgkin and Sternberg-Reed cells have superior anti-tumor effects against H-RS cells *in vitro* and solid Hodgkin tumors in mice. *Int J Cancer* 63:304-9, 1995
20. Messmann RA, Vitetta ES, Headlee D, Senderowicz AM, Figg WD, Schindler J, Michiel DF, Creekmore S, Steinberg SM, Kohler D, Jaffe ES, Stetler-Stevenson M, Chen H, Ghetie V, Sausville EA: A phase I study of combination therapy with immunotoxins IgG-HD37- deglycosylated ricin A chain (dgA) and IgG-RFB4-dgA (Combotox) in patients with refractory CD19(+), CD22(+) B cell lymphoma. *Clin Cancer Res* 6:1302-13, 2000
21. Ghetie MA, Tucker K, Richardson J, Uhr JW, Vitetta ES: Eradication of minimal disease in severe combined immunodeficient mice with disseminated Daudi lymphoma using chemotherapy and an immunotoxin cocktail. *Blood* 84:702-7, 1994
22. Ghetie MA, Podar EM, Gordon BE, Pantazis P, Uhr JW, Vitetta ES: Combination immunotoxin treatment and chemotherapy in SCID mice with advanced, disseminated Daudi lymphoma. *Int J Cancer* 68:93-6, 1996
23. Wei BR, Ghetie MA, Vitetta ES: The combined use of an immunotoxin and a radioimmunoconjugate to treat disseminated human B-cell lymphoma in immunodeficient mice. *Clin Cancer Res* 6:631-42, 2000
24. Baluna R, Vitetta ES: Vascular leak syndrome: a side effect of immunotherapy. *Immunopharmacology* 37:117-32, 1997
25. Baluna R, Rizo J, Gordon BE, Ghetie V, Vitetta ES: Evidence for a structural motif in toxins and interleukin-2 that may be responsible for binding to endothelial cells and initiating vascular leak syndrome. *Proc Natl Acad Sci U S A* 96:3957-62, 1999
26. Baluna R, Vitetta ES: An *in vivo* model to study immunotoxin-induced vascular leak in human tissue. *J Immunother* 22:41-7, 1999
27. Baluna R, Coleman E, Jones C, Ghetie V, Vitetta ES: The effect of a monoclonal antibody coupled to ricin A chain-derived peptides on endothelial cells *in vitro*: insights into toxin-mediated vascular damage. *Exp Cell Res* 258:417-24, 2000

SAMENVATTING

Immunotoxines (ITs) zijn hybride moleculen bestaande uit een zogenaamde monoclonale antistof (MoAb) gekoppeld aan een effectieve gifstof (toxine). De gebruikte MoAbs hebben de eigenschap dat ze zeer sterk, en met een hoge mate van specificiteit, binden aan een eiwit (antigeen) dat voorkomt op de buitenkant van de te vernietigen cel. Eenmaal in de bloedbaan ingespoten, zullen de ITs zich aan deze cellen kunnen binden. Het IT-antigeen complex wordt vervolgens door de cel opgenomen (geïnternaliseerd), waarna het toxine zijn werk kan doen en de cel zal uitschakelen (zie Figuur 1, Hoofdstuk 1). Daar de gebruikte toxines op eigen kracht de cel niet binnenkomen, treffen ITs in principe alleen die cellen waartegen het MoAb gericht is.

Het UMC St Radboud introduceerde 15 jaar geleden als een van de eerste research centra het IT onderzoek in Nederland. De eerste klinische toepassing betrof de zuivering van autologe beenmergtransplantaten van patiënten met een kwaadaardige bloedziekte. Doel van deze behandeling was de eventueel aanwezige kankercellen in het beenmergtransplantaat op te ruimen om daarmee de kans op terugkeer van de kwaadaardige ziekte te verkleinen. De min of meer routinematige toepassing van deze behandeling kan beschouwd worden als het startpunt van dit proefschrift. Deze toepassing leidde op zichzelf tot additioneel onderzoek om de mate van opzuivering van het beenmerg te verbeteren, alsook om deze op een betrouwbare manier te monitoren. Internationaal leidde het enthousiasme over de mogelijke therapeutische toepassingen onder tussen tot diverse (meest Amerikaanse) studies waarbij ITs aan patiënten werden toegediend voor de bestrijding van diffuse en solide tumoren en uiteenlopende immuunziekten. Aangemoedigd door de behaalde resultaten, speciaal op het gebied van beenmerg- en lymfeklierandoeningen, en ingegeven door de klinische behoefte, besloot de afdeling Hematologie een op IT gebaseerde therapie te ontwikkelen voor omgekeerde afstoting na beenmergtransplantatie (graft-versus-host disease: GVHD). Dit proefschrift beschrijft een deel van het IT onderzoek zoals dat gedurende de laatste tien jaar in het UMC St Radboud is uitgevoerd. Na een algemene inleiding (**Hoofdstuk 1**) en de positionering binnen de hematologie (**Hoofdstuk 2**) volgt een beschrijving van het fundamentele laboratorium onderzoek (**Hoofdstukken 3 en 5**) en de opzet en eerste klinische resultaten van een pilot-studie voor de behandeling van GVHD (**Hoofdstukken 6 en 7**). In het hierop volgende worden Hoofdstukken 3 tot en met 7 in meer detail besproken, gevolgd door een algemene beschouwing over de huidige en toekomstige rol van ITs.

Hoofdstuk 3 beschrijft een verbeterde methode om de cellulaire opname te bepalen van radioactief gelabelde MoAbs of daarvan afgeleide ITs. Het bijzondere aan deze methode is dat ze bij de bepaling van de totale internalisatie rekening houdt met moleculen die intracellulair zijn afgebroken en vervolgens door de cel zijn uitgescheiden. Dit

is met name van belang voor ITs, waarbij dergelijke moleculen kunnen bijdragen aan de intoxicatie van een cel door het toxine af te leveren alvorens te worden afgebroken.

Bovenstaande methode werd gebruikt om de relatie tussen internalisatie en effectiviteit van een anti-CD3 IT (WT32-RTA) te bestuderen. Met behulp van leukemische T-cellijn Jurkat, kon een voortdurende cellulaire opname van het anti-CD3 MoAb/IT worden gedemonstreerd. Deze werd gekenmerkt door een snelle initiële opname gedurende de eerste 12 uur, gevolgd door een langzamere maar voortdurende internalisatie gedreven door constante reëxpressie van het CD3-antigeen. Toevoeging van NH_4Cl of monensine vertraagde de intracellulaire afbraak, wat resulteerde in een toegenomen hoeveelheid intracellulair aanwezig IT/MoAb. Theoretisch leidt dit tot een verhoogde kans op afgifte van het toxine, en daarmee een verlaging van het aantal geïnternaliseerde moleculen benodigd voor de intoxicatie van de cel. Deze observaties kwamen sterk overeen met de daadwerkelijke effectiviteit van het anti-CD3 IT zoals bepaald in een eiwitsynthese inhibitie assay. Zowel een verlenging van de incubatietijd, alsook de toevoeging van NH_4Cl of monensine, leidde tot een verhoogde cytotoxiciteit ten aanzien van Jurkat cellen. Een interessante observatie hierbij was dat de maximale eiwitsynthese remming iets leek na te lopen, ten gevolge van een vertraging in de internalisatie ofwel daaropvolgende verwerking van membraangebonden IT. Vertaald naar de zuivering van beenmerg, pleiten bovenstaande resultaten voor een 24-uurs incubatie met IT in de aanwezigheid van NH_4Cl of monensine. Dit zou resulteren in een optimale effectiviteit, ook ten aanzien van cellen met lage antigene dichtheid, zonder schade te berokkenen aan normale hematopoietische progenitorcellen.

Voor wat betreft de toekomst, kunnen internalisatiestudies van pas komen bij de bepaling van het optimale doseringsschema voor klinische studies. Zij zouden een onderbouwing kunnen leveren voor protocollen met een continue dan wel meerdere kortdurende infusies, afhankelijk van het effect op de internalisatie. In het geval van kortdurende infusies kunnen zij bovendien van nut zijn voor de bepaling van het optimale interval tussen twee opeenvolgende infusies. Verlenging van dit interval, bijvoorbeeld, kan leiden tot een meer volledige reëxpressie van het doelwit-antigeen, en daarmee, een verhoging van de gevoeligheid voor een volgende dosis.

De relevantie van internalisatie studies wordt onderstreept in **Hoofdstuk 4**. Deze studie behandelt het opmerkelijke verschil in effectiviteit tussen twee anti-CD19 ITs die alleen verschillen in MoAb-isotype (IgG1 en IgG2a). Internalisatie studies met zowel intact IgG als F(ab')_2 fragmenten, brachten aan het licht dat de aanwezigheid van het anti-CD19 IgG1-Fc gedeelte resulteert in een verhoogde cellulaire opname. De rol van $\text{Fc}\gamma\text{RII}$ (CD32) in deze kon worden aangetoond door een verlaagde effec-

tiviteit van het anti-CD19 IgG1-IT in aanwezigheid van FcγRII-blokkerende MoAb. Deze resultaten wijzen op een interactie van membraangebonden anti-CD19 IgG1-MoAb met FcγRII, resulterende in een verhoogde internalisatie van het antigeen-IT complex, naar analogie van het zogenaamde Kurlander-effect. Met behulp van additionele experimenten kon (functioneel) FcγRIIa vervolgens geïdentificeerd worden als het verantwoordelijke receptor-isotype. De relatief bescheiden toename in internalisatie gaat gepaard met een onevenredige verhoging van de cytotoxiciteit (ongeveer 3-voudig en 100-voudig, respectievelijk). Dit wijst erop dat de toename in internalisatie gepaard gaat met een afwijkende intracellulaire route gunstig voor de effectiviteit van het toxine, ofwel dat de hoeveelheid intracellulair IT een kritische grens overschrijdt waarboven ze niet meer effectief geïnactiveerd kan worden. De relevantie van dit fenomeen voor andere ITs is nog niet duidelijk. Additionele experimenten met een anti-CD22 IgG1-IT lieten geen positief effect zien van de blokkade van FcγRII op de cytotoxiciteit. Echter, de intracellulaire routing van anti-CD22 IT is veel effectiever dan die van anti-CD19 IT (gegeven de superieure toxiciteit bij gelijke of lagere internalisatie). Enige positieve effecten van het IgG1-isotype hebben daarom waarschijnlijk slechts een marginaal effect op de reeds hoge effectiviteit van het CD22-IT.

Bovenstaande *in vitro* studie geeft een duidelijk voorbeeld van het potentiële belang van het MoAb Fc-gedeelte op de uiteindelijke IT effectiviteit. Bij de daadwerkelijke klinische toepassing zal dit nog duidelijker naar voren komen, denkend aan de mogelijke interacties met het humane immuun systeem. Als zodanig vormen bovenstaande data een argument tegen het routinematig uitsluiten van Fc-fragmenten bij zogenaamde 'recombinante fusietoxines'. Deze missen de eventuele bijdrage van Fc-gebaseerde effector mechanismen aan de toxine-gebaseerde opruiming van ongewenste cellen.

Voor de ontwikkeling van een optimale IT therapie is het van belang het aantal ongewenste cellen te bepalen dat ontsnapt aan verschillende behandelingschema's. In **Hoofdstuk 5** wordt een gevoelige flowcytometrische methode beschreven om de IT-geïnduceerde celdood in verse patiëntenmonsters op een adequate wijze te kunnen monitoren. Deze methode is gebaseerd op de flowcytometrische identificatie van dode en levende cellen na kleuring met viabiliteitsmarkers. De exacte kwantificering van overlevende cellen wordt mogelijk gemaakt door bijmenging van een bekende hoeveelheid inerte bolletjes voorafgaande aan de analyse. Een belangrijke bevinding bij het gebruik van deze methode was de totale omvang van de IT-geïnduceerde celdood pas na enkele dagen aangetoond kan worden.

Zoals gedemonstreerd in Hoofdstuk 5, kan met de beschreven methode de opruiming van ongewenste cellen in een heterogene populatie als bloed of beenmerg worden

bestudeerd. Het meest eenvoudige is het om voor de identificatie van deze cellen gebruik te maken van een fluorescerend celspecifiek MoAb. Dit vereist echter de expressie van een 'identificatie antigeen' anders dan het therapeutisch gebruikte 'doelwit antigeen'. Deze laatste kan gemoduleerd zijn of bezet met het therapeutische MoAb/IT. Daarnaast kunnen de overblijvende cellen verrijkt zijn voor cellen met een lage expressie van het doelwit antigeen. Als alternatief kan gekozen worden voor de PCR-amplificatie van een celspecifieke DNA-sequentie. Pre-sorting van levende cellen moet in dit geval voorkomen dat het DNA van dode maar nog aanwezige cellen ten onrechte bijdraagt aan het PCR signaal. In de praktijk leidt een 'triple-sort' tot minder dan één op de miljoen foutief gesorteerde dode cellen, wat overeenkomt met de detectielimiet van een gevoelige PCR assay.

De flowcytometrische bepaling van IT-geïnduceerde celdood is inmiddels toegepast in uiteenlopende studies, ondermeer voor de monitoring van IT-effectiviteit in heterogene, vers geïsoleerde, bloed- of beenmergmonsters van patiënten. Deze studies suggereren dat deze methode zeer bruikbaar is voor de optimalisering van IT-therapie, en een waardevolle aanvulling vormt op de conventionele assays.

Het **eerste deel van Hoofdstuk 7** beschrijft de experimenten die de onderbouwing leverden voor de ontwikkeling van een IT-gebaseerde therapie voor aGVHD. Deze experimenten wezen uit dat een anti-CD3 en anti-CD7 IT (SPV-T3a-dgA en WT1-dgA, respectievelijk), een synergistische werking vertonen ten aanzien van de eliminatie van T cellen. Een algemeen voordeel van het gebruik van een combinatie van ITs is dat minder cellen negatief zullen zijn voor twee dan voor één antigeen. Bovendien kunnen cellen die beide antigenen tot expressie brengen beladen worden met meer IT moleculen. Afgezien van dit 'gewone' synergisme, kent de combinatie SPV-T3a-dgA en WT1-dgA nog een aantal extra voordelen. Het CD3-IT levert additionele immunosuppressie door de binding van het CD3/T-cell-receptor (CD3/TCR) complex. De onderliggende oorzaken, modulatie van het CD3/TCR complex en activatie geïnduceerde celdood (AICD), kunnen van vitaal belang zijn tijdens de behandeling van acute levensbedreigende situaties zoals therapieresistente aGVHD. Het anti-CD3 MoAb (SPV-T3a) is geselecteerd op basis van zijn 'niet-activerende' IgG2b-isotype, welke het risico op 'cytokine release syndrome' sterk vermindert. De toevoeging van het CD7-IT is essentieel, afgezien van het bovengenoemde 'gewone' synergisme, omdat het ook NK cellen (grotendeels CD3⁺ CD7⁺) elimineert. Er zijn steeds meer aanwijzingen voor de rol van NK cellen in de pathofysiologie van aGVHD. Bovenstaande bevindingen vormden de wetenschappelijke basis voor de klinische evaluatie van de IT-combinatie in patiënten met therapieresistente GVHD.

De vertaling van laboratoriumexperimenten naar (vroeg) klinische studies vraagt een grondige aanpassing van het productieproces. Afgezien van de opschaling van materialen en grondstoffen, moeten de bewerking, zuiverheid en activiteit van het eindproduct aan veel strengere eisen voldoen. **Hoofdstuk 6** beschrijft de productie zoals uitgevoerd op het Centraal Hematologisch Laboratorium onder supervisie van de afdeling Klinische Farmacie, en met de ondersteuning van, onder andere, de afdelingen Pathologie, Virologie en het Centraal Dierenlaboratorium. Om een zo hoog mogelijke veiligheid en functionaliteit van het eindproduct te garanderen, werd van te voren een ontwerpqualiteit geformuleerd gebaseerd op het FDA-document 'Aandachtspunten in de vervaardiging en het testen van monoclonale antilichaam producten voor humaan gebruik'. Beschrijvingen van de productie en validatie van vergelijkbare IT die reeds getest waren in FDA-goedgekeurde trials vormden een additionele richtlijn. De 'in huis' productie leverde voldoende materiaal op voor de voorgenomen pilot-studie. De specificaties van het eindproduct kwamen grotendeels overeen met de ontwerpqualiteit. De enkele uitzonderingen, aangaande het percentage ongebonden WT₁ en de DNA-hoeveelheden van WT₁-dgA en SPV-T3a-dgA, vielen, retrospectief, binnen aanvaardbare grenzen. De LD₅₀ (lethale dosis voor 50% van de dieren) voor muizen, kwam overeen, of was zelfs hoger, dan die van vergelijkbare IT die reeds in de kliniek werden toegepast. De voornaamste zorg betrof de kruisreactiviteit van SPV-T3a met humaan oesophagus epitheel, waargenomen tijdens de *in vitro* screening van een humaan weefselpanel. Ter bestudering van de klinische relevantie van deze bevinding, werden humane doses IT-combinatie toegediend aan een tweetal cynomolgus apen. In deze studies kon geen toxiciteit noch binding van het SPV-T3a-dgA aan de oesophagus worden aangetoond. Therapeutische serumconcentraties (>1.8 µg/ml) werden in beide apen bereikt gedurende meerdere uren, zonder dat er onomkeerbare acute toxiciteiten optraden. De belangrijkste bijwerking was een tijdelijke verhoging van creatine kinase (CK)-waarden, wijzend op spieraafbraak door het toxine.

Bovenstaande data rechtvaardigden de evaluatie van het geproduceerde materiaal in een klinische pilot-studie voor de bestrijding van levensbedreigende aGVHD. De onderliggende 'in huis' productie kan wat dat betreft gezien worden als een essentiële stap richting de toekomstige grootschalige productie voor de daadwerkelijke registratiestudies.

Tot nog toe zijn vijf patiënten behandeld met drie of vier infusies van 2 of 4 mg/m² IT-combinatie, toegediend middels een vier uur durend intraveneus infuus met tussenpozen van 48 uur. De resultaten zoals behaald met de eerst vier patiënten staan beschreven in het **tweede deel van Hoofdstuk 7**. De gemiddelde serum halfwaardetijd was ongeveer 7 uur, met piek serumwaarden variërend tussen 258 tot 3.210 ng/ml.

Ondanks dat piek serumconcentraties werden behaald die gelijk ($n = 1$) of boven ($n = 4$) de klinisch veilige grens lagen zoals gedefinieerd voor vergelijkbare IT ($1 \mu\text{g/ml}$), werden geen ernstige toxiciteiten waargenomen. De bijeffecten waren beperkt tot een geringe vochtretentie ($n = 1$), koorts ($n = 2$) en een lichte stijging van CK waarden ($n = 1$), wijzend op een maximaal tolereerbare doses van 4 mg/m^2 of meer. Dit opmerkelijk gunstige toxiciteitsprofiel is waarschijnlijk terug te voeren op de toediening van twee verschillende ITs, welke mogelijk resulteert in een distributie en daarmee verdunning van individuele bijeffecten. Daarnaast kan de gelijktijdige toediening van hoge doses corticosteroïden de maximaal tolereerbare dosis positief beïnvloed hebben. Slechts één van de tot nog toe behandelde patiënten ontwikkelde een geringe antilichaamrespons tegen ricine A. Afgezien van de hoge dosering corticosteroïden, kan dit verklaard worden door een onvolledige reconstitutie van het B-cel compartiment en/of de succesvolle eliminatie van accessoire T-cellen door de IT-combinatie.

Alle vijf patiënten vertoonden een duidelijke biologische respons, gekenmerkt door een gefaseerde daling van circulerende T/NK cellen. De initiële snelle depletie van T-cellen, reeds gedurende de infusies, lijkt MoAb-gerelateerd, terwijl de daaropvolgende meer geleidelijke daling van T/NK cellen wijst op een ricine A gebaseerd mechanisme. Een patiënt ontwikkelde direct na de laatste infusie een sterke oligoclonale expansie van CD8-positieve T-cellen, niet gepaard gaande met een terugkeer van de GVHD (indicatief voor een anti-virale immuun reactie). De andere patiënt wiens perifere T-cellen voor een langere tijd gevolgd konden worden, liet twee weken na start van de behandeling een geleidelijke terugkeer van circulerende (waarschijnlijk naïeve) T-cellen zien. Twee van de vijf patiënten overleden voor toediening van het laatste infusie ten gevolge van reeds bestaande complicaties. De drie overige patiënten vertoonden een duidelijke klinische respons, ondanks dat hun ziekte niet gereageerd had op de voorafgaande behandeling.

Bovenstaande gegevens wijzen op een duidelijke 'therapeutische window'. Dit kan verklaard worden door de combinatie-benadering die gelijktijdig de toxiciteit vermindert en de effectiviteit verhoogt door een beroep te doen op meerdere, synergistische, effector mechanismen. Niettegenstaande de duidelijke biologische en klinische responsen, zijn alle vijf patiënten uiteindelijk overleden aan transplantatie gerelateerde complicaties (meest infecties). Dit is inherent aan het type studie, welke alleen de behandeling van patiënten met vergevorderde, therapieresistente, ziekte toestaat. De bemoedigende eerste klinische resultaten rechtvaardigen nu de evaluatie bij patiënten met minder vergevorderde GVHD. Daarnaast kunnen andere indicaties overwogen worden die baat zouden hebben bij de tijdelijke opruiming van T- en NK-cellen. Meest voor de hand liggend is de behandeling van afstoting na transplantatie (van stamcellen of solide orga-

nen). Afgezien hiervan hebben dierstudies overtuigend aangetoond dat anti-T-cel ITs de juiste immunologische omgeving kunnen creëren voor niet-myeloablatieve stamcel-transplantatie, het realiseren van tolerantie-inductie, of zelfs de behandeling van bepaalde T-cel gemedieerde autoimmuun-ziekten.

Algemene beschouwing

De introductie van nieuwe therapeutische strategieën roept dikwijls hooggespannen verwachtingen op ten aanzien van de behandeling van levensbedreigende ziekten. Deze worden vaak gevolgd door de onvermijdelijke tegenslagen die veelal gepaard gaan met de ontwikkeling van nieuwe medicijnen. ITs vormen hierop geen uitzondering. Volgend op de eerste publicaties, die de genezing beschreven van muizen getransplanteerd met levensbedreigende humane tumoren, werden ITs beschouwd als immunologische 'magic bullets'. Gedurende de daaropvolgende vertaalslag naar de kliniek, stuitte men echter op belangrijke obstakels die onmiddellijke therapeutisch successen in de weg stonden. Een aantal van deze obstakels was reeds waargenomen bij de vroege MoAb-studies. Duidelijke voorbeelden vormen het optreden van antilichaam-reacties, in dit geval tevens gericht tegen het toxine, en de heterogene expressie van doelwit-antigenen. Afgezien hiervan, lijkt de opzet van Fase I/II studies, met de inclusie van patiënten met therapie-resistente uitgebreide ziekte, niet optimaal voor de evaluatie van de potentie van MoAbs of ITs. De belangrijkste tegenvaller was echter het optreden van ernstige bijwerkingen welke voorheen niet in die mate bij proefdieren waren waargenomen. In het geval van ricine-gebaseerde ITs betrof dit met name ernstige 'vaatlekkage syndroom'. De hieruit volgende, relatief lage, maximaal tolereerbare doses resulteerden in maximale serumconcentraties die beduidend lager lagen dan de therapeutische concentraties waargenomen in de oorspronkelijke muizenmodellen. Alles bij elkaar temperden deze resultaten de oorspronkelijke hype, wat als gevolg had dat slechts een relatief klein groep van onderzoekers van het eerste uur overbleef om het concept nieuw leven in te blazen. De ITs werden hiertoe teruggebracht in het diermodel om daar de verschillende obstakels één voor één aan te pakken met de beschikbaar gekomen nieuwe revolutionaire technieken (o.a. recombinant DNA-technieken). Zoals het zich nu laat aanzien, lijkt dit doorzettingsvermogen beloond te worden.

Wat betreft de immunogeniciteit, worden momenteel verschillende strategieën toegepast. Een van deze is gebaseerd op de identificatie van humane toxine-equivalenten (b.v. humane RNases). Koppeling van dergelijke stoffen aan antilichamen, groeifactoren of cytokines van humane origine, resulteert in principe in hybride moleculen met lage of afwezige immunogeniciteit. Deze benadering heeft reeds geresulteert in de constructie van verscheidene 'volledig humane' ITs met bewezen *in vitro* activiteit. Een

andere strategie is gericht op de verlaging van de immunogeniciteit van de uiterst potente conventionele toxines. Dit kan ofwel door de genetische modificatie van de meest immunogene determinanten, ofwel door deze chemisch af te schermen met polyethyleen glycol ('PEGylation'). In combinatie met humane MoAb, zou het gebruik van dergelijke gemodificeerde toxines resulteren in zogenaamde 'Stealth ITs' welke nagenoeg onzichtbaar zijn voor het immuunsysteem van de patiënt. Als alternatief voor bovenstaande strategieën, bestaat er ook nog de mogelijkheid van een tijdelijke onderdrukking van immuunreacties door een gedoseerd gebruik van immuun-onderdrukkende medicijnen.

De aanpak van heterogene antigene expressie wordt gezocht in de gelijktijdige toediening van meerdere ITs gericht tegen verschillende antigenen. Het synergistische effect van dergelijke combinaties is inmiddels in verscheidene dierstudies op overtuigende wijze aangetoond. Op dit moment verschijnen publicaties van de eerste klinische resultaten verkregen met dergelijke combinaties. De IT-combinatie beschreven in Hoofdstukken 5 en 6 onderschrijft bovendien het belang van MoAb-gebaseerde effector mechanismen op de algehele effectiviteit van IT.

Verscheidene dierexperimenten hebben inmiddels overduidelijk het nut aangetoond van het gebruik van ITs als aanvulling op de conventionele chemotherapie of bestraling. Deze bemoedigende resultaten brengen de klinische evaluatie van ITs in de meer geëigende adjuvant setting dichterbij.

Wat betreft de dosis limiterende bijwerkingen, is de meest in het oog springende ontwikkeling waarschijnlijk de opheldering van de oorzaak van vaatlekkage syndroom (VLS). In de A-keten van ricine is een aminozuurketen volgorde (motief) geïdentificeerd die ook aanwezig is in de andere conventionele plantaardige en bacteriële toxines alsook in het eveneens vaatlekkage inducerende Interleukine-2 (IL-2). Dit motief blijkt verantwoordelijk voor VLS door de binding en destructie van endotheliale cellen. Naar verwachting zal de verwijdering of mutatie van dit motief, ofwel het gebruik van niet-destructieve blokkerende peptides, de therapeutische index van conventionele ITs sterk verhogen.

Het leidt geen twijfel dat de bovenstaande ontwikkelingen zullen resulteren in de productie van nieuwe ITs met sterk verbeterde *in vivo* karakteristieken. Alles wijst er dan ook op dat ITs levensvatbaarder zijn dan ooit, en binnen afzienbare tijd tenminste een deel van hun oorspronkelijke beloften in zullen lossen. De FDA-goedkeuring van een recombinant IT (Ontak, DAB389IL-2) voor de behandeling van een therapieresistente hematologische maligniteit, kan dan ook beschouwd worden als een belangrijke mijlpaal.

LIST OF ABBREVIATIONS

AICD	activation-induced cell death
ATG	antihymocyte globuline
AUC	area under the concentration versus time curve
BMT	bone marrow transplantation
BSA	bovine serum albumin
Calc	calcein AM
CD	cluster of differentiation
CK	creatine kinase
C _{max}	maximum plasma concentration
CsA	cyclosporin A
CTL	cytotoxic T lymphocyte
dgA	deglycosylated ricin A-chain
DLI	donor lymphocyte infusion
DLT	dose limiting toxicity
DT	diphtheria toxin
DTT	dithiothreitol
ELISA	enzym-linked immunosorbent assay
ER	endoplasmatic reticulum
Fab	antigen binding fragment of monoclonal antibody
Fv	variable fragment of Fab
FCM	flow cytometry
FCS	fetal calf serum
FITC	fluorescein isothiocyanate
FS	forward scatter
aGVHD	acute graft-versus-host disease
HAMA	human-anti-mouse antibodies
HARA	human-anti-ricin antibodies
HLA	human leucocyte-associated antigen
HRP	horseradish peroxidase
ID ₅₀	50% inhibiting dose
IFN	interferon
IL	interleukin

Ig	Immunoglobulin
IT	immunotoxin
kDa	kilo Dalton
LD ₅₀	50% lethal dose
MHC	major histocompatibility complex
MoAb	monoclonal antibody
MTD	maximal tolerable dose
NHL	non-Hodgkin's lymphoma
NK cell	natural killer cell
NO	nitric oxide
PBL	peripheral blood lymphocytes
PBS	phosphate buffered saline
PE	<i>Pseudomonas</i> exotoxin
pH	hydrogen ion strength
PHA	phytohaemagglutinin
PI	propidium iodide
RAS	right angle scatter
RIP	ribosome inhibitory protein
SDS-PAGE	sodium dodecyl sulphate-polyacrylamide gel electrophoresis
SMPT	4-succinimidylloxycarbonyl- α -methyl- α -(2-pyridyldithio)toluene
SPDP	<i>N</i> -succinimidyl-3-(2-pyridyldithio)propionate
SD	standard deviation
STC	stem cell transplantation
TCR	T cell receptor
TNF	tumor necrosis factor
T _{1/2}	plasma half-life
TTP	thrombotic thrombocytopenic purpura
VH	heavy chain variable domain
VL	light chain variable domain
VLS	vascular leak syndrome
WBC	white blood cell count

LIST OF PUBLICATIONS

- 1: Van Oosterhout YVJM, Preijers FWMB, Wessels HMC, De Witte T: Cytotoxicity of CD3-ricin A chain immunotoxins in relation to cellular uptake and degradation kinetics. *Cancer Res* 52:5921-5925, 1992
- 2: Van Oosterhout YVJM, Van den Herik-Oudijk IE, Wessels HMC, De Witte T, Van de Winkel JGJ, Preijers FWMB: Effect of isotype on internalization and cytotoxicity of CD19-ricin A immunotoxins. *Cancer Res* 54:3527-3532, 1994
- 3: Van Oosterhout YVJM, Preijers FWMB, Meijerink JPP, Pennings AHM, De Witte T: A quantitative flow cytometric method for the determination of immunotoxin-induced cell kill in marrow grafts. *Prog Clin Biol Res* 389:89-95, 1994
- 4: Van Horssen PJ, Van Oosterhout YVJM, De Witte T, Preijers FWMB: Cytotoxic potency of CD22-ricin A depends on intracellular routing rather than on the number of internalized molecules. *Scand J Immunol* 41:563-569, 1995
- 5: Van Horssen PJ, Preijers FW, Van Oosterhout YVJM, De Witte T: Highly potent CD22-recombinant ricin A results in complete cure of disseminated malignant B-cell xenografts in SCID mice but fails to cure solid xenografts in nude mice. *Int J Cancer* 68:378-383, 1996
- 6: Van Horssen PJ, Van Oosterhout YVJM, Evers S, Backus HHJ, Van Oijen MGCT, Bongaerts R, De Witte T, Preijers FWMB: Influence of cytotoxicity enhancers in combination with human serum on the activity of CD22-recombinant ricin A against B cell lines, chronic and acute lymphocytic leukemia cells. *Leukemia* 13:241-249, 1999

7. Van Oosterhout YVJM, Van Ernst JE, Schattenberg AVMB, Tax WJ, Ruiter DJ, Spits H, Nagengast FM, Masereeuw R, Evers S, De Witte T, Preijers FWMB: A combination of anti-CD3 and anti-CD7 ricin A-immunotoxins for the in vivo treatment of acute graft versus host disease. *Blood* 95:3693-3701, 2000
8. Van Horssen PJ, Preijers FWMB, Van Oosterhout YVJM, Eling WMC, De Witte T: Relationship of the CD22 immuntoxin dose and the tumor establishment in a SCID mice model. *Leukemia Lymphoma* 39; 591-599, 2000
9. Van Oosterhout YVJM, Van Ernst JE, Bakker HH, Preijers FWMB, Schattenberg AVMB, Ruiter DJ, Evers S, Koopman JP, De Witte T: production, in an academic setting, of anti-CD3 and anti-CD7 ricin A-immunotoxins for a clinical pilot study. *Int J Pharm*, in press

DANKWOORD

De verschijningsdata van de in dit proefschrift opgenomen artikelen geven al aan dat de promotie geen vluiggertje geweest is. Het aantal mensen dat ik dank verschuldigd ben is met de jaren meegegroeid. Zonder anderen daarbij tekort te willen doen, wil ik enkele van hen graag met name noemen.

Theo, jij hebt als promotor een cruciale rol gespeeld. Ook toen bij andere het geloof in de ‘magic bullets’ tanende was, bleef jij enthousiast over de mogelijke toepassingen van immunttoxines. Ik ben je dan ook zeer dankbaar voor je beslissing om, min of meer tegen de stroom in, het IT-combinatie concept in de praktijk te brengen. Je enthousiasme en doelgerichte aanpak heeft het project door menig moeilijk moment gesleept. *Frank*, jij hebt mij wegwijs gemaakt in de wereld van immunotoxines. Het in dit proefschrift beschreven onderzoek steunt op de stevige grondvesten die jij als een van de internationale pioniers op dit gebied hebt gelegd. Bedankt ook voor je nauwgezette correcties van de manuscripten die je onder ogen kreeg. Ondanks, of misschien wel dankzij, dat we niet altijd op een lijn zaten bleven onze discussies zeer levendig en vormden vaak een drijfveer om ons (heel vaak ook on-)gelijk te halen op het lab. *Ton*, jij vormde als ‘chef de clinique’ de cruciale schakel naar de patiënten. Het door jouw uitgestraalde vertrouwen en onuitputtelijke enthousiasme werkte als een katalysator op het onderzoek. Ik wil je ook bedanken voor je altijd snelle en adequate correcties. *Hans*, jou ben ik dankbaar voor de reeds in een vroeg stadium toegezegde ondersteuning vanuit het laboratorium. Zonder deze hadden we de risicovolle pilot-studie niet van de grond gekregen.

Liesbeth, jij stond mee aan de basis van het IT-combinatie idee en hebt het leeuwendeel van de laboratoriumwerkzaamheden voor je rekening genomen. Je handigheid en accuratesse bij de uitvoering van experimenten deden me al snel beseffen dat ik praktisch op het lab eigenlijk nog maar weinig te zoeken had. Afgezien van de gezelligheid en onze vruchtbare ‘hak op de tak’ besprekingen, wil ik je bedanken voor je inzet, doorzettingsvermogen en inventiviteit waarmee je het onderzoek mee hebt vormgegeven. Nooit klopte ik vergeefs bij je aan. *Sabine*, jij maakte indruk door de voortvarende wijze waarop je nieuwe testen introduceerde en door je inzet bij de evaluatie van patiënten monsters. Ook jij heel erg bedankt voor de fijne samenwerking. Op het lab waren er verder natuurlijk *Arie Pennings*, *Peter Linssen*, *Elly Geestman* en de andere collega's die ik wil bedanken voor hun praktische ondersteuning en gezellige momenten. Ook de studenten *Sonja*, *Zeban*, *Carolien*, en *Arnold* ben ik zeer dankbaar voor hun inbreng bij het beschreven onderzoek. *Peter Donnelley* wil ik noemen voor zijn (vaak spoed) correcties van Engelse teksten.

Peter van Horssen, als mede immuntoxine-promovendus heb je de ‘ups en downs’ van dichtbij meegemaakt. Ik denk met plezier terug aan onze fijne samenwerking en wil je bijzonder bedanken voor de steun en adviezen gedurende het onderzoek. *PeWe, Toon, Jules*, niet te geloven dat wij met z’n vieren dat kleine kamertje gedeeld hebben. Jullie naambordjes hangen vandaag de dag nog steeds aan de deur, als een soort van eerbetoon aan de (enige echte) mannen die ‘de Barak’ gekend heeft. Jullie heel erg bedankt voor de gezelligheid en onderlinge support. Ook de latere bewoners *Maria* en *Isabelle* wil ik bedanken voor de gezelligheid.

Een van de leuke ervaringen bij de uitvoering van het beschreven onderzoek was de enthousiaste en actieve ondersteuning door andere afdelingen. In eerste instantie was daar Klinische Farmacie. Met op twee A-4tjes samengevat de karakteristieken van de IT-combinatie, verschenen we bij *Hans Bakker* en *Chiel Hekster* met het verzoek om zo snel mogelijk met de klinische studie te kunnen beginnen. Een goed jaar later kwamen we weer bij ze vandaan. De A4-tjes waren aangezwollen tot twee 5 cm-dikke ordners vol met beschrijvingen van de op het medicijn uitgevoerde testen. Deze omvangrijke evaluatie en vastlegging was het resultaat van een nauwe en prettige samenwerking tussen de afdelingen Hematologie en Klinische Farmacie. *Hans, Chiel* en *collega's* bedankt voor jullie inspanningen en buitengewoon constructieve houding in deze. Een ander treffend voorbeeld wordt gevormd door *Dirk Ruiter, Cathy Maass* en *collega's* van de afdeling Pathologie. *Professor Ruiter* besteedde menig uur in de namiddag aan het nauwgezet screenen van humane weefsels op ongewenste kruisreactiviteit. Heel veel dank hiervoor. Daarnaast wil ik bedanken *Jos van der Logt* en *Frans Heessen* voor de uitvoering van de biosafety testen, *Albert Peters* en *collega's* voor de prettige en professionele ondersteuning bij de dierexperimenten, *Fokke Nagengast* voor de endoscopie bij de apen en niet te vergeten *Roos Masereeuw* voor de farmacokinetiek analyses. De Verpleging, *Toine, Lucia* en *collega's*, wil ik bedanken voor het mee opzetten en de uitvoering van de monitoring van patiënten. Door jullie toedoen is deze zeer accuraat en zo min mogelijk belastend verlopen. Ik heb heel veel respect voor de professionele wijze waarop jullie de meestal zeer zieke patiënten begeleiden en verzorgen. *Quirine* wil ik bedanken voor de snelle en accurate verwerking van de patiënten gegevens. *Wil Tax* en *Hergen Spits*, jullie bedankt voor het ter beschikking stellen van de hybridoma's en de wetenschappelijke adviezen gedurende het onderzoek. Ook de inbreng van *Jan van de Winkel* mag niet onvermeld blijven. Jan, afgezien van de samenwerking voor de CD19 studie, wil ik je bedanken voor je luisterend oor en je waardevolle adviezen. Ik heb bewondering voor de ongekennde wijze waarop je mensen weet te motiveren.

Los van de (klinisch-)wetenschappelijke kant was er ook de kant van octrooien, -subsidies, adviseurs en contracten. Op dit gebied ben ik dank verschuldigd aan *Wilma*

Derksen (Farma Research), *Emile Douma* en *Gerrit den Hartog* (Ernst en Young), *Theone Kon* (Amersham Pharmacia Biotech), *Jan de Koning*, *Tosca* en *Lotte* (CSC), *Hein van de Pasch* (Stichting Gelderkennis), *Jan Prins* (Stork), *Johan Renes* en *collega's* (Verenigde) en *Eljo Schuite* (Sigma-Tau Ethifarma). Jullie allen bedankt voor de prettige samenwerking en praktische ondersteuning. Ook de bondgenoten bij het 'Immunotoko avontuur', *Joost Ballemans*, *Hans Bolluijt*, *Gerard Hoskens*, *Frits Smedts* en *Rolph Wouters* wil ik bedanken voor hun belangeloze inzet. Vanzelfsprekend ben ik ook veel dank verschuldigd aan de subsidiegever NWO. De direct betrokkenen, *Nico Boots* en *Werner Most*, wil ik complimenteren voor de constructieve en prettige wijze waarop ze hun projecten begeleiden.

Ik wil eindigen met het bedanken van familie, vrienden en kennissen voor hun belangstelling en steun. *Geert-Jan* en *Marcel*, jullie bedankt voor de ontelbare adviezen en morele support. *Ansel*, *Twan* en *Timo* ook jullie heel erg bedankt voor jullie meeleven en meedenken. *Pa* en *Ma*, jullie ben ik heel veel dank verschuldigd voor alle ondersteuning door de jaren heen. Zonder jullie hand en spandiensten was er van promoveren weinig terechtgekomen (ik hoop met dit proefschrift het 'middagje Pietje Huisman' eindelijk weggepoetst te hebben). Ik ben jullie en *Hermien* ook heel dankbaar voor die ontelbare keren dat jullie bij ons hebben opgepast en het werk gewoon door kon gaan. En bovenal was er natuurlijk *Joyce*. Ik ben heel blij dat je er altijd voor mij was, zonder jou was dit nooit gelukt. Ik heb je jaren voor (de gek) gehouden dat de promotie er bijna aan zat te komen, dat we alleen 'even die klinische studie moesten doen'. Bedankt voor je begrip en de ruimte die je me gaf om mijn droom na te jagen in plaats van 'snel' te promoveren en een 'normale' baan te zoeken. Ik moet niet zelden als een Don Quichote zijn overgekomen. *Jelle*, *Youp* en *Merel*, jullie bedankt voor jullie opgewekte en stimulerende aanwezigheid.

CURRICULUM VITAE

De schrijver van dit proefschrift werd geboren op 30 mei 1964 te Nijmegen. In 1984 behaalde hij het Atheneum-B diploma aan het Elshof College te Nijmegen, waarna hij begon met de studie Biologie aan de Katholieke Universiteit Nijmegen. In het kader van deze studie werden stages gelopen bij de afdeling Parasitologie van het bedrijf Intervet te Boxmeer (Dr. A. Vermeulen), de afdeling Moleculaire Biologie van de Katholieke Universiteit Nijmegen (Prof. Dr. R.N.H. Konings†) en de afdeling Hematologie (Prof. Dr. C. Haanen, later Prof. Dr. T. de Witte) van het UMC St Radboud te Nijmegen. In september 1989 behaalde hij het doctoraal examen Biologie met als specialisatie 'Toegepaste Richting'. Vanaf die tijd is hij werkzaam op het Centraal Hematologisch Laboratorium (Dr. J. Wessels, later Dr. E. Mensink) van het UMC St Radboud, als wetenschappelijk onderzoeker op het door Dr. F. Preijers geïnitieerde Immunotoxine onderzoek. Gedurende dit onderzoek is het plan opgevat om de veelbelovende laboratorium resultaten te vertalen in een experimenteel medicijn, de zogenaamde immuno-toxine-combinatie (IT-combinatie), voor patiënten met levensbedreigende omgekeerde afstotingsziekte (graft-versus-host disease: GVHD). Middels een subsidie van de Stichting Technische Wetenschappen (Technologiestichting STW) van de Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), en met een actieve ondersteuning door diverse afdelingen binnen het UMC St Radboud, heeft de schrijver een pilot-studie opgezet om de IT-combinatie op beperkte schaal in de kliniek te testen. De 'in huis' productie en daaropvolgende klinische toepassing vormen als zodanig de bekroning van het in dit proefschrift beschreven onderzoek. Op basis van de eerste hoopgevende resultaten is besloten de IT-combinatie verder bedrijfsmatig te ontwikkelen, waartoe in 1998 het aan het UMC St Radboud gelieerde 'Immunotoko B.V.' is opgericht. Dit initiatief is ondersteund door de NWO die de schrijver een subsidie heeft toegekend in het kader van het Stimuleringsprogramma voor Innovatief Geneesmiddelenonderzoek en Ondernemerschap in Nederland (STIGON). Met deze support gaat hij, als bestuurder van Immunotoko B.V., de komende periode trachten de IT-combinatie verder te ontwikkelen richting registratie als een 'Orphan Drug' tegen GVHD en afstoting van transplantaten.

De schrijver is getrouwd met Joyce Burgers en de trotse vader van Jelle, Youp en Merel.

